


213



GENERAL HEADQUARTERS
SUPREME COMMANDER for the ALLIED POWERS
PUBLIC HEALTH and WELFARE SECTION



Public Health and Welfare
in
Japan

Annual Summary — 1949
Volume II
(Part One)

1950

W2

MS9

P9 P

1949

v.2, pt.1

0021

Public Health and Welfare in Japan - 1949

VOLUME II

PROVISIONAL SUMMARY OF HEALTH AND WELFARE STATISTICS - 1949

-- Foreword --

The Provisional Summary of Health and Welfare Statistics for 1949 supplements the tabulations published in the 1948 Summary plus thirteen additional historical tabulations previously not available. This volume is in two parts.

Public Health and Welfare in Japan - 1949

VOLUME II

PROVISIONAL SUMMARY OF HEALTH AND WELFARE STATISTICS - 1949

Index to Contents

	<u>Page</u>
Introduction	3
Population and Factors Affecting Birth and Death Rate Trends . .	4
Population	4
Factors Affecting Birth and Death Rate Trends	4
Births	5
Number and Rates	5
Seasonal Rates	6
Mortality and Morbidity	6
Deaths According to Age	7
Deaths From Selected Causes	7
Tuberculosis (All Forms)	7
Intracranial Lesions of Vascular Origin	8
Diarrhea, Enteritis and Ulceration of Intestines	9
Senility	10
Cancer and Other Malignant Tumors	10
Pneumonia (All Forms)	11
Diseases of the Heart	11
Congenital Debility	11
Accidental Deaths	12
Nephritis (All Forms)	12
Bronchitis	13
Ulcer of Stomach or Duodenum	13
Ill-defined and Unknown Causes	13
Suicide	14
Premature Birth	14
Measles	15
Simple Meningitis	15
Whooping Cough	16
Dysentery	16
Congenital Malformations	17
Pleurisy Not Specified as Tuberculous	17
Beriberi	18
Syphilis	18
Diseases of Pregnancy, Childbirth and Puerperium . . .	19
Appendicitis	19
Tetanus	19
Diabetes Mellitus	20
Homicide	20
Diphtheria	20
Japanese "B" Encephalitis	21
Injury at Birth	22
Typhoid Fever	22
Poliomyelitis	22
Cerebrospinal Meningitis	23

Public Health and Welfare in Japan - 1949

	<u>Page</u>
Deaths From Selected Causes (Continued)	
Influenza	23
Leprosy	24
Paratyphoid Fever	24
Scarlet Fever	25
Malaria	25
Rabies	25
Typhus Fever and Typhus-like Diseases	26
Glanders	26
Smallpox	26
Yellow Fever	26
Plague	26
Cholera	27
Anthrax	27
Chancroid	27
Gonorrhea	27
Lymphogranuloma Venereum	27
Trachoma	28
Puerperal Infection	28
Dengue Fever	28
Infant Mortality.	28
Infant Deaths From Selected Causes	29
Congenital Debility	29
Pneumonia (All Forms)	30
Diarrhea, Enteritis and Ulceration of Intestines	30
Premature Birth	30
Bronchitis	31
Other Diseases Peculiar to the First Year of Life	31
Congenital Malformations	31
Whooping Cough	32
Measles	32
Beriberi	33
Ill-defined and Unknown Causes	33
Meningitis, Not Due to Meningococcus	33
Accidental Deaths	34
Convulsions	34
Tuberculosis (All Forms)	34
Syphilis	35
Injury at Birth	35
Non-Puerperal Septicemia and Purulent Infection	35
Tetanus	36
Erysipelas	36
Diphtheria	36
Influenza	37
Dysentery	37
Cerebrospinal (Meningococcus) Meningitis	37
Japanese "B" Encephalitis	38
Scarlet Fever	38
Typhus Fever and Typhus-Like Diseases	38
Typhoid Fever	38
Paratyphoid Fever	39
Cholera	39
Anthrax	39

Public Health and Welfare in Japan - 1949

	<u>Page</u>
Infant Deaths From Selected Causes (Continued)	
Leprosy	39
Glanders	39
Malaria	39
Yellow Fever	40
Rabies	40
Stillbirths	40
Marriages	41
Diverces	41
Hospital Strength	42
Number of Hospitals	42
Bed Capacity	42
Total Patient Load	43
In-Patient Load	44
Out-Patient Load	45
Bed Occupancy	46
Venereal Disease Clinic Activities	48
Nutrition Surveys	50
School Lunch Program	50
Public Assistance Program	51
Non-Nationals	51
Births, Deaths and Infant Deaths to Japanese Nationals Outside of Japan	51

Public Health and Welfare in Japan - 1949

Index to Charts

Chart

- A-1 Birth and Death Rates: Japan, 1875-1949
- A-2 Birth Rates by Month: Japan, 1947-1949 and Median Rate 1935-1941
- A-3 Prefectural Vital Statistics: 1949 Calendar Year (Birth Rate and Death Rate)
- A-4 Birth and Death Rates by Month: Japan, 1939-1949
- A-5 Death Rates by Month: Japan, 1947-1949 and Median Rate 1935-1941
- A-6 Death Rates for Ten Leading Causes of Death in 1949: Japan, 1923-1943 and 1947-1949
- A-7 Prefectural Morbidity Statistics: 1949 Calendar Year (Tuberculosis, Dysentery and Malaria)
- A-8 Prefectural Morbidity Statistics: 1949 Calendar Year (Influenza, Smallpox and Measles)
- A-9 Prefectural Morbidity Statistics: 1949 Calendar Year (Scarlet Fever, Whooping Cough, Diphtheria)
- A-10 Prefectural Morbidity Statistics: 1949 Calendar Year (Japanese "B" Encephalitis, Typhus Fever and Pneumonia)
- A-11 Prefectural Morbidity Statistics: 1949 Calendar Year (Typhoid Fever, Paratyphoid Fever and Epidemic Meningitis)
- A-12 Infant Death and Stillbirth Rates: Japan, 1899-1949
- A-13 Infant Death Rates by Month: Japan, 1947-1949 and Median Rate 1938-1942
- A-14 Prefectural Vital Statistics: 1949 Calendar Year (Infant Death Rate and Stillbirth Rate)
- A-15 Death Rates for Eight Selected Causes of Infant Deaths in 1949: Japan, 1923-1943 and 1947-1949
- A-16 Infant Death and Stillbirth Rates by Month: Japan, 1939-1949
- A-17 Stillbirth Rates by Month: Japan, 1947-1949 and Median Rate 1935-1941
- A-18 Marriage and Divorce Rates by Month: Japan, 1939-1949
- A-19 Marriage Rates by Month: Japan, 1947-1949 and Median Rate 1932-1938

Public Health and Welfare in Japan - 1949

Chart

- A-20 Prefectural Vital Statistics: 1949 Calendar Year
(Marriage Rate and Divorce Rate)
- A-21 Marriage and Divorce Rates: Japan, 1883-1949
- A-22 Divorce Rates by Month: Japan, 1947-1949 and Median Rate
1932-1938
- A-23 Epidemiological Case Card (Cholera, Dysentery, Ekiri,
Typhoid, Paratyphoid)
- A-24 Epidemiological Case Card (Smallpox, Typhus Fever, Scarlet
Fever, Diphtheria, Cerebrospinal Meningitis, Plague,
Japanese "B" Encephalitis)
- A-25 Epidemiological Case Card (Tuberculosis)
- A-26 Morbidity Schedule (Communicable Diseases)
- A-27 Morbidity Schedule (Tuberculosis)
- A-28 V.D. Case Reporting Card and Morbidity Schedule

Public Health and Welfare in Japan - 1949

Index to Tables

Table

- 1 Population by Prefecture: Japan, 1949
- 2 Population by Age: Japan, 1948-1949
- 3 Population, Live Births, Deaths, Infant Deaths, Stillbirths,
Marriages and Divorces: Japan, 1947-1949
- 4 Live Birth, Death, Infant Death, Stillbirth, Marriage and
Divorce Rates: Japan, 1947-1949
- 5 Live Births by Month: Japan, 1947-1949
- 6 Live Birth Rates by Month: Japan, 1947-1949
- 7 Live Births by Prefecture: Japan, 1947-1949
- 8 Live Birth Rates by Prefecture: Japan, 1947-1949
- 9 Deaths by Month: Japan, 1948-1949
- 10 Death Rates by Month: Japan, 1948-1949
- 11 Deaths by Prefecture: Japan, 1948-1949
- 12 Death Rates by Prefecture: Japan, 1948-1949
- 13 Deaths from Selected Causes: Japan, 1948-1949
- 14 Death Rates for Selected Causes: Japan, 1948-1949
- 15 Deaths from Selected Causes by Month: Japan, 1920-1949
- 16 Death Rates for Selected Causes by Month: Japan, 1920-1949
- 17 Deaths from Ten Selected Causes by Prefecture: Japan,
1920-1949
- 18 Death Rates for Ten Selected Causes by Prefecture: Japan,
1920-1949
- 19 Deaths from the Ten Leading Causes of Death: Japan, 1948-
1949
- 20 Death Rates for the Ten Leading Causes of Death: Japan,
1948-1949
- 21 Rank Order of the Ten Leading Causes of Death: Japan,
1948-1949
- 22 Deaths by Age: Japan, 1948
- 23 Death Rates by Age: Japan, 1948

Public Health and Welfare in Japan - 1949

Table

- 24 Deaths from Tuberculosis (All Forms) by Age: Japan, 1948-1949
- 25 Death Rates for Tuberculosis (All Forms) By Age: Japan, 1948-1949
- 26 Live Births, Maternal Deaths, and Maternal Death Rates: Japan, 1947-1949
- 27 Cases of Selected Communicable Diseases: Japan, 1948-1949
- 28 Case Rates for Selected Communicable Diseases: Japan, 1948-1949
- 29 Cases of Selected Communicable Diseases by Month: Japan, 1948-1949
- 30 Case Rates for Selected Communicable Diseases by Month: Japan, 1948-1949
- 31 Cases of Selected Communicable Diseases by Prefecture: Japan, 1920-1949
- 32 Case Rates for Selected Communicable Diseases by Prefecture: Japan, 1920-1949
- 33 Natural Increase in Population: Japan, 1947-1949
- 34 Infant Deaths by Month: Japan, 1947-1949
- 35 Infant Death Rates by Month: Japan, 1947-1949
- 36 Infant Deaths by Prefecture: Japan, 1947-1949
- 37 Infant Death Rates by Prefecture: Japan, 1947-1949
- 38 Infant Deaths from Selected Causes: Japan, 1947-1949
- 39 Infant Death Rates for Selected Causes: Japan, 1947-1949
- 40 Infant Deaths from the Ten Leading Causes of Infant Deaths in 1949: Japan, 1947-1949
- 41 Infant Death Rates for the Ten Leading Causes of Infant Deaths in 1949: Japan, 1947-1949
- 42 Stillbirths by Month: Japan, 1947-1949
- 43 Stillbirth Rates by Month: Japan, 1947-1949
- 44 Stillbirths by Prefecture: Japan, 1947-1949
- 45 Stillbirth Rates by Prefecture: Japan, 1947-1949

Public Health and Welfare in Japan - 1949

Table

- 46 Marriages by Month: Japan, 1947-1949
- 47 Marriage Rates by Month: Japan, 1947-1949
- 48 Marriages by Prefecture: Japan, 1947-1949
- 49 Marriage Rates by Prefecture: Japan, 1947-1949
- 50 Divorces by Month: Japan, 1947-1949
- 51 Divorce Rates by Month: Japan, 1947-1949
- 52 Divorces by Prefecture: Japan, 1947-1949
- 53 Divorce Rates by Prefecture: Japan, 1947-1949
- 54 Venereal Disease Clinic Activities, Japan, 1947-1949
- 55 Consumption of Total and Rationed Food per Capita per Day
in Calories: Tokyo, Other Large Cities and Rural Areas,
1947-1949
- 56 Food Consumption in Grams per Capita per Day: Tokyo, Other
Large Cities and Rural Areas, 1947-1949
- 57 Nutritive Value of Food Consumed Daily: Tokyo, Other Large
Cities and Rural Areas, 1947-1949
- 58 Average Body Height and Weight: Japan, 1947-1949
- 59 School Lunch Program by Month: Japan, 1946-1949
- 60 Public Assistance Program by Month: Japan, 1946-1949
- 61 Live Births, Deaths, Infant Deaths, Stillbirths, Marriages
and Divorces by Prefecture: Japan, 1949
- 62 Live Birth, Death, Infant Death, Stillbirth, Marriage and
Divorce Rates by Prefecture: Japan, 1949
- 63 Live Births by Month by Prefecture: Japan, 1949
- 64 Live Birth Rates by Month by Prefecture: Japan, 1949
- 65 Deaths by Month by Prefecture: Japan, 1949
- 66 Death Rates by Month by Prefecture: Japan, 1949
- 67 Deaths from Selected Causes by Month: Japan, 1949
- 68 Death Rates for Selected Causes by Month: Japan, 1949
- 69 Deaths from Selected Causes by Prefecture: Japan, 1949

Public Health and Welfare in Japan - 1949

Table

- 70 Death Rates for Selected Causes by Prefecture: Japan, 1949
- 71 Deaths from Selected Causes by Month by Prefecture, Japan, 1949
- 72 Death Rates for Selected Causes by Month by Prefecture: Japan, 1949
- 73 Maternal Deaths and Death Rates by Month: Japan, 1949
- 74 Maternal Deaths and Death Rates by Prefecture: Japan, 1949
- 75 Maternal Death Rates by Month by Prefecture: Japan, 1949
(Rates per 1,000 Live Births Each Month)
- 76 Communicable Disease Cases and Deaths and Communicable Disease Case and Death Rates: Japan, 1949 (Rates per 100,000 Population per Annum)
- 77 Communicable Disease Cases by Month: Japan, 1949
- 78 Communicable Disease Case Rates by Month: Japan, 1949
(Rates per 100,000 Population per Annum)
- 79 Communicable Disease Deaths by Month: Japan, 1949
- 80 Communicable Disease Death Rates by Month: Japan, 1949
(Rates per 100,000 Population per Annum)
- 81 Communicable Disease Cases and Deaths and Communicable Disease Case and Death Rates by Prefecture: Japan, 1949
(Rates per 100,000 Population per Annum)
- 82 Communicable Disease Cases by Month by Prefecture: Japan, 1949
- 83 Communicable Disease Case Rates by Month by Prefecture: Japan, 1949 (Rates per 100,000 Population per Annum)
- 84 Infant Deaths by Month by Prefecture: Japan, 1949
- 85 Infant Death Rates by Month by Prefecture: Japan, 1949
(Rates per 1,000 Live Births Each Month)
- 86 Infant Deaths from Selected Causes by Month: Japan, 1949
- 87 Infant Death Rates for Selected Causes by Month: Japan, 1949 (Rates per 1,000 Live Births Each Month)
- 88 Infant Deaths from Selected Causes by Prefecture: Japan, 1949

Public Health and Welfare in Japan - 1949

Table

- 89 Infant Death Rates for Selected Causes by Prefecture: Japan, 1949 (Rates per 1,000 Live Births)
- 90 Infant Deaths from Selected Causes by Month by Prefecture: Japan, 1949
- 91 Infant Death Rates for Selected Causes by Month by Prefecture: Japan, 1949 (Rates per 1,000 Live Births Each Month)
- 92 Stillbirths by Month by Prefecture: Japan, 1949
- 93 Stillbirth Rates by Month by Prefecture: Japan, 1949 (Rates per 1,000 Live Births Each Month)
- 94 Marriages by Month by Prefecture: Japan, 1949
- 95 Marriage Rates by Month by Prefecture: Japan, 1949 (Rates per 1,000 Population per Annum)
- 96 Divorces by Month by Prefecture: Japan, 1949
- 97 Divorce Rates by Month by Prefecture: Japan, 1949 (Rates per 1,000 Population per Annum)
- 98 Hospital Strength by Kind of Hospital and by Prefecture: Japan, 1949
- 99 Total Case Load of Venereal Disease Clinics by Month: Japan, 1949
- 100 Patients Not Previously Treated by Any Clinic or Physician Admitted to Venereal Disease Clinics by Month: Japan, 1949
- 101 Cases Discontinuing Treatment Without Permission at Venereal Disease Clinics and Number of Visits for Delinquent Cases by Month: Japan, 1949
- 102 Contacts Reported at Venereal Disease Clinics by Month: Japan, 1949
- 103 Visits for Contact Tracing and Number of Contacts Found by Venereal Disease Clinics, by Month: Japan, 1949
- 104 Consumption of Total and Rationed Food in Calories per Capita per Day Each Quarterly Nutrition Survey: All Japan, Tokyo, Other Cities and Rural Areas, 1949
- 105 Nutrients in Grams and Calories per Capita per Day Each Quarterly Nutrition Survey: All Japan, Tokyo, Other Cities, and Rural Areas, 1949

Public Health and Welfare in Japan - 1949

Table

- 106 Food Consumption in Grams per Capita per Day Each Quarterly Nutrition Survey: All Japan, Tokyo, Other Cities and Rural Areas, 1949
- 107 Nutritive Value of Food Consumed According to Quarterly Nutrition Surveys: All Japan, Tokyo, Other Cities and Rural Areas, 1949
- 108 Number of Persons and Percent of Total Surveyed Showing Nutritional Deficiency Symptoms According to Nutrition Surveys: All Japan, Tokyo, Other Cities and Rural Areas, 1949
- 109 Number of Persons Found Underweight and Overweight Measured from the Standard Weight by Age Groups: All Japan, Tokyo, Other Cities and Rural Areas, 1949
- 110 Percent of Persons Found Underweight and Overweight Measured from the Standard Weight by Age Groups: All Japan, Tokyo, Other Cities and Rural Areas, 1949
- 111 Average Weight Deviation in Kilograms for Persons Above and Below the 10 percent Level Measured from the Standard Weight by Age Groups: All Japan, Tokyo, Other Cities and Rural Areas, 1949
- 112 Live Births, Deaths, Infant Deaths, Stillbirths, Marriages and Divorces of Non-Japanese Nationals in Japan by Month: 1949
- 113 Live Births, Deaths and Infant Deaths of Japanese Nationals Outside Japan by Month, 1949

Public Health and Welfare in Japan - 1949

VOLUME II

PROVISIONAL SUMMARY OF HEALTH AND WELFARE STATISTICS FOR 1949 AND HISTORICAL DATA IN JAPAN

INTRODUCTION

All data for 1949 in this report are provisional. Historical tabulations included in the report for 1948 have not been repeated, but final data when available or provisional data for 1947, 1948 and 1949 are shown. Reference must be made to both reports for a complete historical series.

Among the new historical tables included in this report are the following: Deaths and Death Rates from Selected Causes By Month, 1920-1949 (Tables 15 and 16); Deaths and Death Rates From Ten Selected Causes By Prefecture, 1920-1949 (Tables 17 and 18); Cases and Case Rates For Selected Communicable Diseases By Prefecture, 1920-1949 (Tables 31 and 32); Venereal Disease Clinic Activities, 1947-1949 (Table 54); Nutrition Surveys, 1947-1949 (Tables 55-58); School Lunch Program, 1946-1949 (Table 59); and Public Assistance, 1946-1949 (Table 60).

Other new tables limited to 1949 include the following: Hospital Strength by Kind of Hospital and by Prefecture (Table 98); Total Case Load of Venereal Disease Clinics By Month (Table 99); Patients Not Previously Treated By Any Clinic or Physician Admitted to Venereal Disease Clinics By Month (Table 100); Cases Discontinuing Treatment Without Permission at Venereal Disease Clinics and Number of Visits for Delinquent Cases by Month (Table 101); Contacts Reported At Venereal Disease Clinics By Month (Table 102); Visits Per Contact Tracing and Number of Contacts Found by Venereal Disease Clinics, By Month (Table 103); Consumption of Total and Rationed Food in Calories Per Capita Per Day, Each Quarterly Nutrition Survey: All Japan, Tokyo, Other Cities and Rural Areas (Table 104); Nutrients in Grams and Calories Per Capita Per Day, Each Quarterly Nutrition Survey: All Japan, Tokyo, Other Cities and Rural Areas (Table 105); Food Consumption in Grams Per Capita Per Day, Each Quarterly Nutrition Survey: All Japan, Tokyo, Other Cities and Rural Areas (Table 106); Nutritive Value of Food Consumed According to Quarterly Nutrition Surveys: All Japan, Tokyo, Other Cities and Rural Areas (Table 107); Number of Persons and Percent of Total Surveyed Showing Nutritional Deficiency Symptoms According to Nutrition Surveys: All Japan, Tokyo, Other Cities and Rural Areas (Table 108); Number of Persons Found Underweight and Overweight Measured from the Standard Weight by Age Groups: All Japan, Tokyo, Other Cities and Rural Areas (Table 109); Percent of Persons Found Underweight and Overweight Measured From the Standard Weight by Age Groups: All Japan, Tokyo, Other Cities and Rural Areas (Table 110); Average Weight Deviation in Kilograms for Persons Above and Below the Ten Percent Level Measured from the Standard Weight by Age Groups: All Japan, Tokyo, Other Cities and Rural Areas (Table 111).

All tabulations of births, deaths, stillbirths, marriages and divorces are by place of occurrence. Data refer to Japanese Nationals only and events which occurred within Japan proper excepting that

Public Health and Welfare in Japan - 1949

Tables 112 and 113 show the number of vital events reported by months for non-Japanese residents in Japan proper and Japanese Nationals outside of it.

Rates are shown by geographic areas referred to as "shi" and "gun". These terms cannot be defined with great exactness, but they constitute a Japanese concept of urban and rural areas. The word "shi" refers to places having a population of 30,000 or more; "gun" to all other places including towns and villages. A "gun" would represent that part of a county exclusive of all "shi" within its borders. The distinction between towns (machi) and villages (mura) is even less definite, although the former would contain more stores, temples and schools than the latter.

All monthly rates, and rates based on data for part of a year, are estimated on an annual basis. Rates for specific causes of death or morbidity are calculated per 100,000 population estimated as of 1 July; infant death rates and stillbirth rates, per 1,000 live births occurring during the same period.

Morbidity data for 1920-1936 refer to the civilian population in Japan and Okinawa. Data subsequent to 1937 are for Japan proper only. The civilian population includes both Japanese Nationals and foreigners, except members of the Occupation Forces.

Monthly death statistics 1947-1949 were based on calendar months, whereas monthly case reports were based on combinations of data for four or five week periods. However, both monthly mortality and morbidity rates were calculated on an annual basis.

POPULATION AND FACTORS AFFECTING BIRTH AND DEATH RATE TRENDS

Population

The estimated population as of 1 July 1949 was 82,200,000 (Ref. Table 1). Tokyo led the list with an estimated population of 5,551,800.

Populations by age are shown in Table 2 by five-year age groups. Beginning with 10,577,000 at 0-4 years the number decreased regularly to 2,709,000 at 55-59 years. For 60 years and over it was 6,291,000.

Populations estimated as of July 1, 1949 which were used for the computation of "shi" and "gun" rates, were as follows: "shi" (28,402,000) and "gun" (53,798,000). For July 1, 1948 they were as follows: "shi" (27,711,000) and "gun" (52,489,000).

Factors Affecting Birth and Death Rate Trends

There were 2,723,873 births and 950,719 deaths in 1949. The excess of births over deaths was 1,773,154 and the natural rate of increase was 21.6 per 1,000 population (Ref. Table 33). It was

slightly lower than the rate (21.9) in the preceding year, which was the highest ever recorded. The birth rate decreased from 33.8 in 1948 to 33.1 in 1949, a reduction of 0.7 per 1,000 population. The death rate over the same period declined 0.3 per 1,000 population. The death rate may be expected to decrease slightly in 1950 but the birth rate is expected to decline markedly. Further decreases in the birth rate are expected to occur in 1951 while the death rate may remain about the same as in 1950. A much greater reduction in the rate of natural increase is expected in 1950 because the abnormally high marriage rate in 1948 (12.0) fell sharply to 10.3 in 1949 and is expected to fall further in 1950.

During the next twenty-five years the birth rate may decline to less than 20 per 1,000 population by 1970 and possibly less than 10 per 1,000 population in the next fifty years. On the other hand, the age specific death rates now being recorded are not expected to continue the same trend in the future. The 1949 rate is already considerably below the rate that would be expected on the basis of its secular trend. As the birth rate continues to decrease in the future, the death rate can be expected to turn upward as the population becomes older in its age composition. It is impossible to say how soon the death rate will do so, but it might well be before 1960 and possibly between 1955 and 1960.

Studies originally made on factors affecting birth and death rates during the summer of 1947 indicate that the estimated population in 1960 is unlikely to be less than 90 million and might be as high as 93 million. Correspondingly, estimates for 1970 may be 95 to 100 million and for 1980, 98 to 106 million.

BIRTHS

Number and Rates

Live births totalled 2,723,873 in 1949. The crude provisional birth rate was 33.1 per 1,000 population which represents a small reduction below the figure (33.8) in 1948. As previously stated in the discussion of the factors affecting birth and death rate trends, the birth rates recorded during the 3-year period 1947-1949 have been abnormally high. It would seem most unlikely that the birth rate would fall enough during 1950 to be in agreement with the expected figure (about 26) according to the trend established 1920-1937. Considering the decline recorded in the marriage rate for 1949 and probable further decrease in 1950, it is quite possible that the rate would approximate the trend figure in 1951. (Ref. Chart A-1 and Tables 3 and 4)

The birth rate for all "shi" combined was 32.7 per 1,000 population in 1949 and for all "gun" 33.4. (Ref. Tables 63 and 64)

Prefectural rates ranged from 27.5 in Nara to 41.4 in Aomori. Other prefectures having rates well above the national average include Hokkaido (40.3), Nagasaki (38.5), Miyazaki (37.6), Iwate (37.5) and

Public Health and Welfare in Japan - 1949

Fukuoka (37.0). Those having rates well below the average include Nagano (27.8), Shiga (28.8), Kyoto (28.9), Wakayama (29.0) and Mie (29.3). (Ref. Tables 7, 8, 61-64)

Seasonal Rates

The monthly distribution of the birth rates followed the usual seasonal cycle, moving from a high point (46.7) in January to a low point (28.1) in June and then rising to a secondary peak (32.8) in September, after which it decreased to 29.3. (Ref. Chart A-2 and Tables 5 and 6) The September peak occurred somewhat earlier than the high point usually recorded in November. In this respect it followed the 1946 and 1947 series. The December rate (29.3) was the highest recorded for this month since 1925. The increase, although less than 5% above the 1948 figure, may be attributed to an educational program to change the practice of registering births which actually occurred in December as though they had occurred in January. With the abolition of the Japanese method of counting age on 1 January 1950, the principal reason for delaying the registration of December births was eliminated. It is believed that the increase in the December rate reflects this fact. There may be some people who will still make December registrations falsely in January in the belief that the occurrence of a birth in January is considered a lucky event, but the number of such persons is not believed to be large. With the closing of the business year and preparations for the long New Year Holiday period, there is a greater tendency to delay registrations of births near the end of December than at other times in the year.

In past years monthly rates for December and January computed on the basis of the number of births have been subject to some error. The December rates were too high and the January rates too low. This would affect computation of the infant death rates and also the maternal death rates.

MORTALITY AND MORBIDITY

Deaths totalled 950,719 in 1949. The crude death rate was 11.6 per 1,000 population, the lowest of record. During the period 1900 to 1925 the annual rate was 20 or above. It was still a little above 20 at the beginning of the 1926-1950 period. By 1941 it had fallen to 15.7, after which it increased gradually to 17.4 in 1944 and then suddenly rose to 29.2 in 1945. It then returned to 17.6 in 1946, continuing downward in 1947 (14.6) and 1948 (11.9).

The death rate (11.6) recorded in 1949 is well below the trend established 1925-1937. While it may decrease still further, the reduction is expected to be slower in the future. With the aging of the population, the rate is expected to take an upward trend, possibly beginning between 1955 and 1960. (Ref. Chart A-1 and Table 4)

The death rate for all "shi" was 11.0 per 1,000 population, and for all "gun" 11.9. Prefectural death rates (Ref. Chart A-3 and

Public Health and Welfare in Japan - 1949

Tables 11 and 12) ranged from 9.7 in Kanagawa to 14.0 in Toyama. Among other prefectures having low rates were Tokyo (9.8), Shizuoka (10.3), Nagano (10.7), Hyogo (10.7) and Yamanashi (10.7). Those having high rates include Iwate (13.9), Aomori (13.5), Ishikawa (13.5), Oita (13.3) and Fukui (13.2).

The monthly distribution of death rates (Ref. Chart A-4 and Tables 9, 10, 65 and 66) in 1949 was similar to that recorded in the preceding year except that rates for January and February were somewhat lower in 1949. The tendency to form a summer peak was very slight, unlike earlier years (Ref. Chart A-5). According to data available since 1920, new low records were made in January, February, March, June, July, August and September.

Deaths According to Age

The distribution of Rates by Age forms the typical j-shaped curve. In every age group (Ref. Tables 22 and 23) the death rates established an all-time low record in 1948. The rate under 4 years (24.1) was less than half the 1929 figure (52.1). Appreciable reductions were made in many of the specific age groups. Data were not available for 1949 except for tuberculosis which was obtained by special tabulation.

Deaths from Selected Causes

Death rates for the following eleven causes were lowest of record according to data recorded in 1920 and thereafter: tuberculosis, diarrhea and enteritis (all ages), congenital debility, nephritis, ill-defined and unknown causes, simple meningitis, pleurisy not specified as tuberculous, beriberi, diphtheria, typhoid fever and paratyphoid fever. Death rates for tetanus, influenza, cerebrospinal meningitis, scarlet fever, leprosy, malaria, smallpox and diseases of pregnancy, childbirth and puerperium equalled the lowest rate previously recorded. There were no deaths recorded for plague, cholera, anthrax and yellow fever. Death rates for senility, dysentery, rabies, intracranial lesions of vascular origin and pneumonia (all forms) were the second lowest of record.

The death rate for congenital malformations was the highest of record. Injury at birth and homicide equalled the highest rate previously recorded. The rate for suicide was the highest since 1937. The rate for cancer and other malignant tumors was the fourth highest of record and the highest since 1942.

Tuberculosis, All Forms (Int. List No. 13-22)

The leading cause of death in 1949 was tuberculosis, which was responsible for 138,764 deaths and a death rate of 168.8 per 100,000 population, the lowest of record. About every seventh death from all causes in 1949 was caused by tuberculosis (Ref. Chart A-6 and Tables 19, 20 and 21).

The death rate from tuberculosis for all "shi" combined was 213.5 per 100,000 population and for all "gun" 145.2.

Public Health and Welfare in Japan - 1949

Prefectural death rates (Ref. Tables 18 and 70) from this disease ranged from 92.0 in Yamanashi Prefecture to 249.6 in Hokkaido. These prefectures held the same relative position in 1948. Other prefectures having rates well above the national average include Aomori (217.1), Kyoto (214.1), Tokyo (209.5), Iwate (206.4), Fukuoka (203.8) and Osaka (201.5). Those having rates much lower than average include Ibaraki (110.5), Tochigi (116.0), Gumma (121.7), Nagano (122.9) and Shizuoka (129.9).

The percent distribution of deaths from tuberculosis in 1949 according to age was as follows: under 5 years (5.2%), 5-9 years (2.3%), 10-14 years (2.0%), 15-19 years (9.1%), 20-24 years (18.8%), 25-29 years (15.5%), 30-34 years (11.2%), 35-39 years (9.1%), 40-44 years (6.8%), 45-49 years (5.6%), 50-54 years (4.7%), 55-59 years (3.8%), 60 years and over (5.9%). (Ref. Table 24)

According to age the death rate for children under 5 years was 67.8 per 100,000 population following which it decreased to 31.7 at 10-14 years. The rate rose rapidly to a peak of 357.2 at 25-29 years after which it decreased to 190.0 at 45-49 years. It rose again to 192.1 at 50-54 years and 195.5 at 55-59 years and then decreased sharply to 130.3 at 60 years and over. (Ref. Charts A-25, A-27 and Table 25)

Cases totalled 469,504 in 1949 (Ref. Chart A-7 and Tables 76 and 81). The case rate was 571.2 per 100,000 population. Comparison can only be made with data recorded in 1948 because tuberculosis was not made a reportable disease until 1947. The first full year for which data were available was 1948, in which 378,851 cases were reported, the rate being 475.0. It will be observed that the case rate increased 20% due to improvement in the case finding program.

The monthly death rate from this disease rose from 159.5 in January to a peak of 186.3 in May, following which it decreased to 146.7 in December. Correspondingly the case rate rose from 390.7 in January to a peak of 681.2 in June, after which it decreased to 484.6 in December.

Intracranial Lesions of Vascular Origin (Int. List No. 83)

The second leading cause of death was intracranial lesions of vascular origin, which caused 100,650 deaths in 1949 or 10.6% of deaths from all causes. The death rate was 122.4 and although it increased above the 1948 figure (117.6) it was the second lowest of record. Data are not available for 1944-1946. With a single exception (fourth in 1947) it has held second place in importance as a cause of mortality since 1939. (Ref. Chart A-6 and Tables 19, 20 and 21)

For all "shi" combined the death rate was 99.9 and for all "gun" 134.4.

The seasonal distribution of rates showed a tendency for the rates to be lower during the summer months and highest in the last and first quarters. (Ref. Table 16) Prefectural rates (Ref. Tables 70 and 72) ranged from 84.3 in Osaka Prefecture to 188.2 in Akita. Other prefectures having death rates well above the national average include

Public Health and Welfare in Japan - 1949

Iwate (181.5), Yamagata (172.6), Ibaraki (159.8), Niigata (159.7), Chiba (152.3) and Nagano (151.2). Those having rates well below the average include Hokkaido (89.1), Tokyo (94.4), Kyoto (94.7), Hyogo (96.2) and Nagasaki (103.5).

Diarrhea, Enteritis and Ulceration of the Intestines (International List No. 119-120)

There were 72,186 deaths from this cause with a death rate of 87.8 per 100,000 population, the third highest of all causes in 1949 and in each of the two preceding years. It held forth place in importance from 1938-1943. Approximately eight of each 100 deaths in 1949 resulted from this cause. The 1949 rate was the lowest of record and substantially below the figure (103.8) in the preceding year. In 1936, it was almost twice (172.6) and in 1923, over three times (283.6) as great.

The death rate for all "shi" combined was 67.4 and for all "gun" 98.6. The rate for rural areas was approximately 46% greater than for the urban. In 1948, it was 55% higher.

Prefectural death rates ranged from 40.4 per 100,000 population in Tokyo-to to 171.7 in Toyama. Other prefectures having rates well above the national average include Fukui (144.6), Aomori (144.9), Shiga (134.6), Akita (125.8) and Yamagata (125.2). Those having rates well below the average include Kanagawa (46.1), Chiba (64.8), Nagano (65.8), Kyoto (70.1) and Aichi (71.9).

Death rates were markedly higher in the summer months reaching 137.6 in August and 133.6 in July. They were approximately double the rates in January and December which were 69.1 and 69.3, respectively. (Ref. Table 16)

Deaths from diarrhea, enteritis and ulceration of intestines under two years (Int. List No. 119) totalled 41,822 from this cause with a death rate of 50.9 per 100,000 population, the lowest of record.

The rate for all "shi" combined was 42.2 and for all "gun" 55.4.

Prefectural rates (Ref. Tables 69 and 70) ranged from 26.6 in Kanagawa to 114.2 in Aomori. Other prefectures having rates well above the average include Toyama (100.5), Iwate (90.5), Akita (89.0), Ishikawa (82.2) and Yamagata (78.5). Those having rates well below the average also include Tokyo-to (27.7), Kochi (28.6), Nagano (28.6), Gumma (32.9) and Okayama (32.9).

In July the death rate was 81.3 forming a seasonal peak with rates in January and December being 40.5 and 43.2 respectively.

Deaths from diarrhea, enteritis and ulceration of intestines two years and over (Int. List No. 120) totalled 30,364 from this cause with a death rate of 36.9 per 100,000 population, the lowest of record.

The rate for all "shi" combined was 25.1 and for all "gun" 43.2.

Public Health and Welfare in Japan - 1949

Prefectural rates ranged from 12.7 in Tokyo to 71.3 in Toyama. Other prefectures having rates well above the national average include Fukui (70.1), Shiga (67.1), Tochigi (56.9), Tokushima (54.9) and Niigata (54.6). Among those having rates well below average were Kanagawa (19.5), Miyagi (22.2), Fukuoka (27.1), Hokkaido (28.0) and Iwate (28.8).

The seasonal distribution of rates showed a peak in August of 60.6. The January and December rates were 28.6 and 26.1 respectively.

Senility (Int. List No. 162)

Senility, the fourth leading cause of death was responsible for 66,699 deaths and a death rate of 81.1. Seven of each 100 deaths from all causes can be attributed to senility. Although the death rate increased a little above the 1948 figure (79.9), it was the second lowest of record. Senility prior to 1948 held fifth position of importance for 24 years.

For all "shi" combined the rate was 57.6 and for all "gun" 93.6.

Prefectural rates (Ref. Tables 18 and 70) ranged from 46.8 in Tokyo to 132.8 in Kochi. Other prefectures having rates well above the national average include Tottori (120.2), Fukui (119.1), Ehime (117.0), Oita (117.0) and Okayama (116.6). Others having rates well below average include Hokkaido (55.1), Osaka (56.4), Kanagawa (68.9), Toyama (64.9) and Miyagi (65.4).

Monthly death rates were highest in the first quarter and lowest during the summer months. (Ref. Table 16).

Cancer and Other Malignant Tumors (Int. List No. 45-5b)

The fifth leading cause of death was cancer (Ref. Table 19) which caused 58,890 deaths. A little more than six out of each 100 deaths from all causes were from this disease. The death rate was 71.6, continuing the uneventful trend in the rate for at least a quarter of a century. It held the same relative position of importance in 1948. Historically it increased from tenth position in 1931 to fifth place in 1949. Cancer has advanced further in its position of importance than any other cause of death since 1920.

The rate for all "shi" combined was 78.7 and for all "gun" 67.9.

Prefectural rates (Ref. Tables 18 and 70) ranged from 47.9 in Iwate to 99.4 in Nara. Other prefectures having rates well above the national average include Wakayama (85.5), Ishikawa (83.4), Niigata (83.1), Tottori (83.1) and Chiba (82.2). Those having rates much lower than average include Aomori (53.2), Kagoshima (55.1), Akita (57.4), Shizuoka (57.5) and Hokkaido (59.3).

As usual, the death rates in the third quarter were somewhat higher than those for the rest of the year and those in the first quarter were lowest. (Ref. Table 16)

Public Health and Welfare in Japan - 1949

Pneumonia, All Forms (Int. List No. 107-109)

The sixth leading cause of death was pneumonia (Ref. Chart A-7 and Tables 13 and 19), which was responsible for 56,636 deaths or 6% of deaths from all causes. The death rate was 68.9 per 100,000 population, the second lowest of record.

For all "shi" combined the death rate was 68.4 and for all "gun" 69.2.

Prefectural rates ranged from 50.5 in Tottori to 103.1 in Iwate. Other prefectures having rates well above the national average include Aomori (92.0), Fukushima (86.3), Gumma (81.4), Toyama (79.6) and Hokkaido (79.5). Those having rates well below the average include Okayama (51.8), Kyoto (54.1), Kochi (54.3) and Hiroshima (55.0).

Pneumonia case reports are available for the second year since the disease was made reportable. The case rates ranged from 80.2 per 100,000 population in Chiba to 479.6 in Toyama. In 1948 Chiba and Toyama also had the lowest and highest case rates from pneumonia. (Ref. Table 32)

Death rates were definitely higher during the first quarter with 108.1 recorded for February. The August rate was exceptionally low with 29.0. (Ref. Table 16)

Diseases of the Heart (Int. List No. 90-95)

The seventh leading cause of death was diseases of the heart which caused 52,825 deaths. Out of the total number of deaths from all causes, approximately 1 out of 20 resulted from this cause. This is the third consecutive year in which this cause of death has held seventh place in importance. For 10 years (1934-1943) it held ninth position. The death rate was 64.3 per 100,000 population, the highest recorded since 1939. Data are not available for 1944-1946.

For all "shi" combined the death rate was 58.9 and for all "gun" 67.1.

Prefectural death rates from this cause ranged from 51.4 in Kochi to 80.2 in Shimane. Other prefectures having rates well above the national average include Ibaraki (77.8), Tokushima (76.6), Nagano (75.3), Gumma (75.0), and Chiba (74.9). Those having rates well below average include Aomori (53.0), Kanagawa (53.2), Fukuoka (54.4), Hokkaido (54.6) and Tokyo (56.4).

The seasonal distribution of rates shows the March figure (74.9) to be highest, followed by January with 73.2. Rates for months in the third quarter were the lowest with 52.7 for August (Ref. Tables 67 and 68).

Congenital Debility (Int. List No. 158)

Congenital debility, the eighth leading cause of death was responsible for 37,285 deaths and the death rate (45.4 per 100,000 population) was the lowest of record. Approximately one out of every 25

Public Health and Welfare in Japan - 1949

deaths can be attributed to this cause. Although congenital debility advanced from ninth place in importance in 1948, in previous years it was frequently in sixth place.

For all "shi" combined the rate was 35.1 and for all "gun" 50.8.

Prefectural rates ranged from 23.0 in Kanagawa to 89.2 in Aomori. Other prefectures having rates well above the national average include Akita (86.7), Iwate (70.7), Saga (70.1), Fukui (65.9) and Yamagata (65.6). Those having rates well below average include Tokyo (24.8), Nagano (30.1), Yamanashi (31.7), Wakayama (34.6) and Hyogo (35.9).

Rates were highest during the first quarter and lowest in the summer months. September had the lowest rate (29.6). (Ref. Table 16)

Accidental Deaths (Int. List No. 169-195)

The ninth leading cause of death (Ref. Tables 19 and 20) was responsible for 34,498 deaths, which was 3.6% of the total number of deaths from all causes. The death rate was 42.0 per 100,000 population and lower than that recorded in each of the two preceding years. However, in earlier years it has often been less.

The death rate for all "shi" combined was 40.7 and for all "gun" 42.6.

Prefectural rates ranged from 29.4 in Shiga to 57.6 in Yamaguchi. Other prefectures having rates well above the national average include Fukuoka (57.3), Fukui (57.0), Saga (55.7), Toyama (55.7) and Iwate (55.2). Those having rates well below average include Saitama (30.6), Kyoto (31.3), Nara (31.7), Chiba (32.8) and Wakayama (33.7).

August had the peak death rate (63.9), approximately twice the rates recorded for January (32.3) and February (31.5). Rates for July and September were 61.9 and 46.9 respectively.

Nephritis. All Forms (Int. List No. 130-132)

The tenth leading cause of death, nephritis, was responsible for 33,859 deaths representing 3.6% of the total number of deaths from all causes. The death rate was 41.2, the lowest of record. In 1939, the rate (82.2) was approximately twice this figure. The maximum recorded rate was 107.5 in 1923.

The death rate for all "shi" was 37.1 and for all "gun" 43.3.

Prefectural rates ranged from 26.6 in Wakayama to 61.6 in Tottori. Other prefectures having rates well above the national average include Saga (60.7), Kagoshima (57.8), Miyazaki (57.7), Oita (51.8) and Ibaraki (51.3). Those having rates well below the average include Hokkaido (27.1), Aichi (30.6), Aomori (30.7) and Kyoto (30.8).

Rates during the summer months were somewhat lower than those recorded in the first and last quarters. (Ref. Table 16)

Bronchitis (Int. List No. 106)

Bronchitis caused 25,695 deaths in 1949 with a death rate of 31.3 per 100,000 population. The rate was a little less than the figure (32.3) recorded in the preceding year, although not as low as in 1941 (30.1) and 1942 (30.4) which were low points of record.

The death rate for all "shi" combined was 21.4 and for all "gun" 36.5. (Ref. Table 70)

Prefectural rates ranged from 15.3 in Tokyo-to to 49.3 in Toyama. Other prefectures having rates well above the national average include Niigata (46.5), Iwate (45.2), Oita (45.0), Ishikawa (44.0) and Mie (43.7). Those having rates well below the average include Kanagawa (16.5), Osaka (19.5), Kyoto (19.8), Miyagi (23.4), Fukuoka (25.6) and Shizuoka (26.1).

Rates were lowest as usual in the third quarter with the rate for August (16.3) being the lowest. The highest rate was recorded for February (48.5).

Ulcer of Stomach or Duodenum (Int. List No. 117)

There were 19,521 deaths from this cause. The death rate was 23.7 per 100,000 population. In 1920, the rate was 11.2. The secular trend was upward and by 1943 the rate had risen to 23.0. Data are not available for the period 1944-1946. The highest point (33.2) was reached in 1947 following which it decreased to 25.8 in 1948.

The death rate for all "shi" combined was 22.7 and for all "gun" 24.3.

Prefectural rates ranged from 14.6 in Hokkaido to 33.8 in Oita. Other prefectures having rates well above the national average include Shimane (32.5), Ishikawa (30.5), Yamaguchi (30.3), Tottori (29.6) and Kumamoto (29.2). Those having rates well below the average include Aomori (17.5), Miyagi (18.6), Akita (18.2), Ehime (19.7) and Aichi (20.0).

Monthly death rates ranged from 19.4 in August to 27.6 in December. (Ref. Table 16)

Ill-Defined and Unknown Causes (Int. List No. 200a and c)

Deaths from this cause totalled 16,900 in 1949, the death rate being 20.6 per 100,000 population, the lowest of record. In 1943 it was 42.6. The secular trend has been downward for many years. During the 6-year period between 1943 and 1949, the rate decreased about 50%. It is of interest to note that during the 21-year period 1921-1942 the rate decreased less than 50%.

In 1949, a little less than 18 medical certifications of death out of each 1,000 deaths registered were incomplete or contained unacceptable information thus preventing proper statistical classification of the cause of death. In 1948 the ratio was a little more than

Public Health and Welfare in Japan - 1949

18. The extent of the improvement in the quality of medical certifications is shown by the fact that in 1920 the ratio was approximately 50 per 1,000 registrations. Efforts have been made through lectures to physicians, regional conferences and the distribution of literature, to obtain properly prepared medical certifications.

It is not possible to construct a continuous historical rate series based on a combination of International List Numbers 199 and 200. Sometimes deaths from "sudden death" have been included and at other times excluded. For that reason the historical series (Ref. Tables 15-18) contained in this report are based on International List Numbers 200a and c. However, data have been included (Ref. Tables 69-72) for 1949 by prefecture for sudden death and ill-defined and unknown causes (199-200).

According to that tabulation the death rate for all "shi" combined was 20.0 and for all "gun" 21.0. The prefectural rates ranged from 12.7 in Tottori to 32.7 in Fukui. Other prefectures having rates well above the national average include Kagawa (31.6), Aomori (30.5), Nara (30.5), Hokkaido (29.4) and Wakayama (26.7). Those having rates well below the average include Kyoto (15.7), Miyazaki (16.1), Aichi (16.3), Shimane (16.3) and Gumma (16.6).

Monthly death rates ranged from 16.5 in September to 24.6 in March. Rates recorded for the summer months were lower than those for the rest of the year.

Suicide (Int. List No. 163-164)

There were 14,211 suicides in 1949 with a death rate of 17.3 per 100,000 population, the highest of record since 1937. Data are not available for the period 1944-1946. Between 1920 and 1932 the trend was gradually upward. Then it turned downward reaching an all-time low record of 11.8 in 1943. In 1947, the first full year for which data were available after the termination of the war, the rate was 15.7. It rose slightly to 15.9 in 1948.

The death rate for all "shi" combined was 18.7 and for all "gun" 16.5.

Prefectural rates ranged from 10.5 in Miyagi to 26.2 in Kyoto and Wakayama. Other prefectures having rates well above the national average include Shiga (24.7), Nagano (24.2), Gifu (21.8), Hyogo (21.4) and Nara (21.3). Those having rates well below the average include Aomori (10.7), Nagasaki (11.3), Yamanashi (12.2), Fukushima (12.3), Ibaraki (12.4) and Kagoshima (12.4).

The peak month for death rates from suicides in 1949 was in May (22.8) with a secondary peak in July (20.0). The same general seasonal pattern was observed in 1948.

Premature Birth (Int. List No. 159)

There were 13,667 deaths from premature birth resulting in a death rate of 16.6 per 100,000 population. This represents a reduction in the rate for the second successive year. In 1947 and 1948,

Public Health and Welfare in Japan - 1949

the rates were respectively 18.7 and 17.1. However, the 1949 rate is still approximately double any rate recorded since 1932 when such data first became available, with the single exception of 1936 (9.0). The trend from 1936 to 1943 was gradually downward.

The death rate for all "shi" combined was 15.0 and for all "gun" 17.5.

Prefectural death rates ranged from 9.3 in Kanagawa to 28.1 in Iwate. Other prefectures having rates well above the national average include Toyama (26.2), Akita (24.9), Ibaraki (24.7), Chiba (23.9), and Yamagata (22.7). Those having rates well below the average include Yamaguchi (9.7), Yamanashi (11.8), Tokyo (12.6), Hokkaido (12.6) and Shizuoka (13.5).

The distribution of death rates by months was not according to usual experience because the lowest rate (13.8) was recorded in October when usually it has been in May or June. (Ref. Table 16)

Measles (Int. List No. 35)

Deaths from measles totalled 12,440 in 1949 with a death rate of 15.1 per 100,000 population. Although the rate was more than twice the 1948 figure (7.0), it is lower than might have been expected as the upper year of a 2-year cycle.

The death rate for all "shi" combined was 16.3 and for all "gun" 14.5.

Prefectural death rates ranged from 2.4 in Kochi to 28.1 in Hokkaido. Other prefectures having rates well above average include Nara (24.8), Osaka (24.4), Fukuoka (24.2), Niigata (23.0), Shiga (21.0) and Toyama (20.6). Those having rates well below the average include Kagawa (2.8), Okayama (3.2), Totterri (4.9), Tokushima (5.3) and Tochigi (6.1).

There were 165,308 cases of measles in 1949 and the case rate was 201.1 per 100,000 population. This disease was made reportable on 20 January 1947. In 1948 the case rate was 68.6.

The distribution of case rates by month in 1949 showed the highest rate to be in May (575.8). Rates for April and June were 366.0 and 511.1 respectively.

Prefectural case rates ranged from 47.6 in Kochi to 627.1 in Shimane. (Ref. Table 81) Other prefectures having case rates well above the national average include Shiga (609.6), Toyama (578.6), Fukuoka (433.1), Saga (382.6) and Kyoto (344.4). Those having rates well below the average include Tochigi (55.0), Okayama (71.4), Tokushima (77.5), Iwate (80.8), Kagawa (82.4) and Chiba (82.8). (Ref. Chart A-8)

Simple Meningitis (Int. List No. 81a)

Simple meningitis caused 9,108 deaths. The death rate was 11.1 per 100,000 population, the lowest of record and less than one-tenth of the 1924 rate (116.2).

Public Health and Welfare in Japan - 1949

The death rate for all "shi" combined was 10.3 and for all "gun" 11.5. (Ref. Table 70)

Prefectural rates ranged from 5.9 in Tokyo to 17.1 in Gifu. Other prefectures having rates well above the national average include Ishikawa (16.5), Akita (16.2), Toyama (15.0), Oita (14.6), Hokkaido (14.4) and Tokushima (14.0). Those having rates well below the average include Yamaguchi (6.5), Nagano (7.1), Tochigi (8.2). The rate was 8.6 in Ibaraki, Tottori and Wakayama prefectures.

The monthly distribution of death rates was unlike the usual record of previous years in which the rate for August or July was generally higher than that for other months. In 1949 the maximum rate was for April (13.6). (Ref. Table 16).

Whooping Cough (Int. List No. 9)

Deaths from whooping cough totalled 9,147, the rate of 11.1 per 100,000 population being almost double the figure (5.9) for the preceding year.

The death rate for all "shi" combined was 12.0 and for all "gun" 10.7.

Prefectural death rates ranged from 3.1 in Wakayama to 21.5 in Yamagata. Other prefectures having rates well above the national average include Aomori (18.9), Gumma (18.0), Miyagi (17.7), Saitama (15.7), Hokkaido (15.4) and Ehime (15.4). Those having rates well below the average include Kochi (3.6), Miyazaki (4.3), Tokushima (4.5) and Kumamoto (5.4). It was 5.7 in Kagoshima, Okayama and Shimane Prefectures. The seasonal distribution of death rates shows that the rate in March (7.6) increased to a peak in August (17.0) and then decreased to 8.7 in November.

There were 126,827 cases and the case rate was 154.3 per 100,000 population, more than twice that (66.2) recorded in the preceding year. This disease was made reportable on 20 January 1947. Prefectural case rates ranged from 41.1 in Kochi to 336.7 in Shiga. Other prefectures having rates well above the national average include Toyama (328.5), Ehime (287.4), Kagawa (261.2), Hokkaido (239.0) and Saitama (238.9). Those having rates well below the average include Kagoshima (53.4), Chiba (62.2), Oita (63.1), Miyazaki (63.3) and Wakayama (64.8). (Ref. Chart A-9)

Dysentery (Int. List No. 27)

There were 7,824 deaths from this disease. The death rate was 9.5 per 100,000 population (Ref. Tables 13 and 14), the second lowest of record. In the preceding year the rate was 6.4. Although the percent increase was sharp, the 1949 figure was only a little greater than one-fourth the rate (34.8) recorded in 1939.

The death rate for all "shi" combined was 9.3 and for all "gun" 9.6.

Public Health and Welfare in Japan - 1949

Prefectural death rates ranged from 2.0 in Shiga to 26.3 in Ibaraki and Saitama prefectures. Other prefectures having rates well above the national average include Gumma (24.3), Tochigi (19.2), Tokyo (15.0), Niigata (14.9), Gifu (14.5) and Shizuoka (14.4). Those having rates well below the average include Hokkaido (2.3), Nara (3.1), Miyagi (3.2), Aomori (3.9) and Akita (4.1).

The peak death rate was in August (34.8) which is usually the month of highest incidence of mortality. (Ref. Table 16)

Cases of dysentery totalled 24,001 and the case rate was 29.2 per 100,000 population. (Ref. Charts A-23, A-26 and Table 81)

Prefectural case rates ranged from 6.0 in Nara to 90.9 in Gumma. Other prefectures having rates well above the national average include Saitama (82.1), Niigata (66.4), Tokyo (60.7), Ibaraki (50.9), and Iwate (47.6). Those having rates well below the average include Shiga (8.5), Kagoshima (9.2), Hyogo (10.2), Okayama (10.2) and Aomori (10.9).

As usual the peak monthly rate was in August (113.7). Although it was nearly twice the rate for the same month last year (64.0), it is still much lower than rates recorded in previous years. The explosive characteristic of this disease during the summer months is shown by the fact that about three out of 10 cases were reported in August and more than 7 out of 10 during July-September.

Congenital Malformations (Int. List No. 157)

There were 6,556 deaths from congenital malformations with a death rate of 8.0 per 100,000 population, the highest of record.

For all "shi" combined the death rate was the same as that recorded for all "gun" which was 8.0.

Prefectural death rates ranged from 6.0 in Shiga to 10.5 in Miyagi and Yamanashi. Other prefectures having rates well above the national average include Gumma (10.4), Tottori (10.2), Fukushima (10.1), Aomori (10.0) and Toyama (10.0). Those having rates well below the average include Nara (6.3) Shimane (6.4), Kochi (6.5), Hyogo (6.5) Kanagawa (6.7), and Nagano (6.7).

The seasonal distribution of mortality rates was uneventful.

Pleurisy Not Specified as Tuberculous (Int. List No. 110)

There were 6,030 deaths from this cause resulting in a death rate of 7.3 per 100,000 population. This extended the sharp reduction in the rate which was 24.1 in 1943.

The death rate for all "shi" combined was 6.8 and for all "gun" 7.6.

Prefectural rates ranged from 5.2 in Gumma to 10.8 in Kumamoto. Other prefectures having rates well above the national average include Oita (10.2), Nagasaki (9.8), Yamaguchi (9.7), Miyazaki (8.5) and

Public Health and Welfare in Japan - 1949

Kagoshima (9.2). Those having rates well below the average include Shizuoka (5.3), Osaka (5.7), Nagano (5.7), Nara (5.8), Kanagawa (5.9) and Miyagi (5.9).

The monthly distribution of death rates was uneventful. (Ref. Table 16)

Beriberi (Int. List No. 68)

There were 5,591 deaths attributed to this disease, the death rate of 6.8 per 100,000 population being the lowest of record.

The death rate for all "shi" combined was 6.4 and for all "gun" 7.0.

Prefectural rates ranged from 1.7 in Yamanashi to 16.6 in Aomori. Other prefectures having rates well above the national average include Iwate (14.1), Ishikawa (11.7), Saga (11.5), Toyama (11.0) and Osaka (10.7). Those having rates well below the average include Kagawa (1.9), Ehime (3.0), Tokushima (3.4), Wakayama (3.5) and Nagano (3.6).

The monthly distribution of death rates was uneventful.

Syphilis (Int. List No. 30)

Deaths from this disease totalled 5,492 in 1949. The death rate was 6.7 per 100,000 population. Although the rate increased above that (5.4) for the preceding year it was one-half to almost one-third the rates recorded between 1923 and 1937.

The death rate for all "shi" combined was 9.0 and for all "gun" 5.4.

Prefectural death rates ranged from 3.6 in Yamanashi to 9.7 in Fukuoka. Other prefectures having rates well above the national average included Tokyo (9.2), Nagasaki (9.0), Kyoto (9.0), Saga (8.8), Osaka (8.3), Kanagawa (8.3). All of these prefectures with the exception of Saga contain large cities. Prefectures having rates well below the average include Niigata (4.2), Fukui (4.3), Ehime (4.4), Hiroshima (4.4) and Toyama (4.4).

There were 188,204 cases of syphilis reported and the case rate was 229.0 per 100,000 population (Ref. Chart A-28).

The monthly distribution of death rates was uneventful. (Ref. Table 16)

Prefectural case rates ranged from 102.6 in Shimane to 472.8 in Kanagawa. Other prefectures having case rates well above the national average include Fukuoka (449.3), Nagasaki (444.6), Osaka (411.7), Kyoto (382.7) and Hyogo (356.4). Those having rates well below average include Niigata (111.4), Ibaraki (112.3), Gifu (116.0), Kagoshima (121.8) and Saitama (122.8).

Public Health and Welfare in Japan - 1949

Diseases of Pregnancy, Childbirth and Puerperium (Int. List No. 140-150)

Deaths from all maternal causes totalled 4,495 and the death rate of 5.5 per 100,000 population equalled the all-time low record of the preceding year. The rate per 1,000 live births was 1.7.

For all "shi" combined the death rate was 5.7 and for all "gun" 5.4.

Prefectures having death rates well above the national average include Akita (9.2), Kagoshima (8.2), Miyazaki (8.1), Iwate (8.0) and Aomori (7.4). Those having rates well below the average include Shiga (3.1), Wakayama (3.7), Nara (3.8), Tokushima (4.3), Shizuoka (4.4) and Aichi (4.4).

Expressed in terms of live births the rates would be as follows: Akita (2.5), Kagoshima (2.3), Miyazaki (2.1), Iwate (2.1), Aomori (1.8), Shiga (1.1), Wakayama (1.3), Nara (1.4), Tokushima (1.3), Shizuoka (1.3) and Aichi (1.4). (Ref. Tables 26, 73, 74 and 75)

The seasonal distribution of the death rates was uneventful.

Appendicitis (Int. List No. 121)

There were 3,267 deaths from appendicitis. The death rate was 4.0. Although this marks the second consecutive year of reduction, the rate has not yet decreased to the secular trend of previous years which reached 3.1 in 1943.

The death rate for all "shi" was 5.5 and for all "gun" 3.2.

Prefectural rates ranged from 2.3 in Kochi to 5.8 in Niigata. Other prefectures having rates well above the national average include Osaka (5.0), Kyoto (4.9), Oita (4.8), Shiga (4.8), and Iwate (4.8). Those having rates well below the average include Tokushima (2.5), Kagoshima (2.5), Nagasaki (2.6), Shizuoka (2.8) and Mie (3.0).

There was a slight tendency for the monthly death rates to rise to a peak (5.0) in August and then decline. (Ref. Table 16)

Tetanus (Int. List No. 12)

There were 1,694 deaths from tetanus with a death rate of 2.4 per 100,000 population. Although 1949 marks the second consecutive year of decrease in the rate, the reduction has been slight and the rate was equal to the low points established in 1939 and 1941.

The death rates for all "shi" and all "gun" were equal at 2.4.

Prefectural death rates ranged from 0.8 in Shiga to 5.6 in Miyazaki. Other prefectures having rates well above the national average include Kagoshima (5.5), Chiba (5.4), Ibaraki (4.9), Saga (4.3) and Kagawa (4.2). Those having rates well below the average

Public Health and Welfare in Japan - 1949

include Kyoto (1.0), Toyama (1.1), Iwate (1.1), Hokkaido (1.2), Hyogo (1.2), Osaka (1.2) and Tottori (1.2).

Death rates rose slightly to 3.2 in September and then decreased to 1.9 in December. Rates in July, August and September are generally a little higher than rates for other months. (Ref. Table 16)

Cases of tetanus totalled 2,199 and the case rate was 2.7 per 100,000 population, a slight increase above the 1948 figure (2.4).

Prefectural case rates ranged from 1.0 in Hokkaido to 7.4 in Ibaraki. Other prefectures having case rates well above the national average include Kagoshima (5.5), Miyazaki (5.4), Chiba (4.8), Ehime (4.8), Kagawa (4.6). Those having rates well below the average include Shiga (1.1), Osaka (1.2), Toyama (1.3), Hyogo (1.4) and Yamagata (1.4).

Diabetes Mellitus (Int. List No. 61)

There were 1,885 deaths attributed to this cause and the death rate, 2.3 per 100,000 population, was slightly higher than the all-time low record (2.2) established in the previous year.

The death rate for all "shi" combined was 1.8 and for all "gun" 2.5.

Prefectural rates ranged from 1.2 in Aomori to 6.0 in Kochi. Other prefectures having rates well above the national average include Ishikawa (4.7), Tottori (4.3), Shimane (4.3) and Kagawa (4.0). Those having rates well below the average include Akita, Fukushima, Osaka, Tochigi and Tokyo, all of which had a rate of 1.4.

The distribution of monthly death rates was uneventful.

Homicide (Int. List No. 165-168)

Homicide was responsible for 1,669 deaths. The death rate was 2.0 per 100,000 population, equalling the all-time high record of the preceding year.

For all "shi" combined the death rate was 2.5 and for all "gun" 1.8.

Prefectural rates ranged from 0.9 in Toyama to 4.0 in Chiba. Other prefectures having rates well above the national average include Fukuoka (3.9) and Kumamoto (3.8). Those having rates well below the average include Kagoshima (1.0) and Niigata (1.1).

The death rate in August and September was 2.4, slightly higher than for other months. (Ref. Table 16)

Diphtheria (Int. List No. 10)

Diphtheria caused 1,659 deaths with a death rate of 2.0 per 100,000 population the lowest of record. (Ref. Tables 13 and 14)

Public Health and Welfare in Japan - 1949

The death rate for all "shi" combined was 2.5 and for all "gun" 1.8.

Prefectural death rates ranged from 0.5 in Wakayama to 5.1 in Miyazaki. (Ref. Table 70) Other prefectures having rates well above the national average include Saga (4.1), Aomori (3.9), Nagasaki (3.8), Kagoshima (3.5), Hokkaido (3.4) and Oita (3.3). Those having rates well below the average include Kyoto (0.7), Aichi (0.8), Saitama (0.8), Nagano (1.0) and Shiga (1.0).

Monthly death rates decreased from 3.5 in January to 0.7 in August. In general a peak is reached each year in December following which the rate decreases to a low point in August in the following year, completing the cycle. (Ref. Table 16)

There were 14,825 cases of diphtheria and the case rate was 18.0 per 100,000 population, the lowest of record. (Ref. Charts A-24, A-26 and Tables 27 and 28)

As usual, the first and last quarters of the year had the highest case rates and midsummer the lowest. The highest monthly case rate was for February (25.7) and the lowest for August (8.7).

Prefectural case rates ranged from 8.0 in Yamanashi to 48.8 in Miyazaki. Other prefectures having rates well above the national average include Saga (44.8), Oita (30.0), Shimane (29.8), Akita (29.1) and Fukuoka (27.8). Those having rates well below the average include Osaka (10.0), Wakayama (10.1), Chiba (10.2), Gifu (10.7) and Kyoto (11.0).

Japanese "B" Encephalitis (Int. List No. 37b)

This disease caused 1,182 deaths resulting in a death rate of 1.4 per 100,000 population; a considerable reduction below the rate (3.3) in 1948 but much higher than the 1947 figure (0.3).

The death rate for all "shi" combined was 2.2 and for all "gun" 1.0.

Prefectural death rates ranged from 0.0 (less than 0.1) in Fukushima and Hokkaido to 4.4 in Kumamoto. Rates were also noticeably high in Toyama (4.3), Fukui (3.5) and Shiga (3.4). They were comparatively low in Kochi (0.1), Yamagata (0.1) and Iwate (0.2) in addition to the two prefectures referred to above.

A review of the seasonal distribution of case rates shows practically no cases during the first half of 1949. A gradual increase in July and August lead to a sudden rise in September to 11.6. (Ref. Table 78)

There were 1,284 cases of this disease reported and the case rate was 1.6 per 100,000 population, almost one-sixth of the rate (9.0) recorded in the previous year. (Ref. Tables 29 and 30)

Case rates were comparatively high in Yamaguchi (4.9), Kanagawa (4.8), Kumamoto (4.7), Hiroshima (4.0) and Tokyo (3.9). Correspondingly

Public Health and Welfare in Japan - 1949

no cases were reported from Aomori, Iwate, Akita, Fukushima, Fukui, Wakayama and Shimane prefectures. However, the highest rates were low compared to the rates recorded in many prefectures last year. (Ref. Chart A-10)

Injury at Birth (Int. List No. 160)

Injuries at birth were responsible for 1,082 deaths, the death rate of 1.3 per 100,000 population, equaling the all-time high rate recorded in the preceding year.

For all "shi" combined the death rate was 1.5 and for all "gun" 1.2.

Prefectural death rates were comparatively high in Toyama (2.5), Kochi (2.3), Miyagi (2.1), Yamagata (2.1) and Kyoto (2.0). On the other hand they were comparatively low in Yamanashi (0.7), Yamaguchi (0.8) and Nagano (0.8) prefectures.

The monthly distribution of death rates was uneventful.

Typhoid Fever (Int. List No. 1)

Typhoid fever caused 970 deaths in 1949. The death rate of 1.2 per 100,000 population was the lowest ever recorded.

The death rate for all "shi" combined was 1.9 and for all "gun" 0.8.

Prefectural rates ranged from 0.2 in Kumamoto and Kagoshima to 2.7 in Okayama. Other prefectures having rates well above the national average include Nara (2.6), Kochi (2.4), Mie (2.3) and Hiroshima (2.1). Those having rates well below the average include Oita (0.3), Miyazaki (0.3), Yamanashi (0.4) and Ehime (0.5).

Monthly death rates increased from a low point (0.8) in January to 2.1 in August and then decreased to 0.9 in December.

Cases totalled 6,489 and the annual case rate was 7.9 per 100,000 population. The seasonal distribution of case rates shows a peak of 13.9 in August, with rates decreasing on either side of it to 4.0 in March and 6.1 in November.

Prefectural case rates ranged from 0.7 in Kagoshima to 19.0 in Mie. Other prefectures having rates well above the national average include Fukui (18.1), Tokyo (16.1), Hiroshima (13.7), Gifu (13.4) and Miyagi (12.8). Those having rates well below the average include Yamanashi (1.3), Kumamoto (1.4), Oita (2.6), Miyazaki (3.2) and Yamaguchi (3.3). (Ref. Chart A-11 and Table 31)

Poliomyelitis (Int. List No. 36)

There were 1,024 deaths from this disease in 1949 and the death rate of 1.2 per 100,000 population was the same as in the preceding year.

Public Health and Welfare in Japan - 1949

The rate for all "shi" combined was 1.1 and for all "gun" 1.3.

Prefectural death rates ranged from 0.3 in Tottori to 3.1 in Oita. Other prefectures having rates above the national average include Kagoshima (1.9), Aomori (1.8) and Aichi (1.7). Those having rates below the average include Yamanashi (0.7), Fukui (0.8), Saitama (0.8), Tokyo (0.8) and Yamagata (0.8).

There was a very slight tendency for the death rate to rise in September and October. (Ref. Table 16)

There were 3,140 cases of poliomyelitis and the case rate was 3.8 per 100,000 population. Comparatively high rates were recorded for Aomori (15.0), Hokkaido (12.2) and Miyagi (10.2) prefectures. Correspondingly, low rates were recorded for Hiroshima (0.6), Yamanashi (0.7) and Shiga (0.8).

Cerebrospinal Meningitis (Int. List No. 6)

There were 497 deaths from this disease. The death rate was 0.6 per 100,000 population.

The death rate for all "shi" combined was 1.0 and for all "gun" 0.4.

Prefectural death rates ranged from zero in Tokushima and Wakayama where no deaths were recorded to 1.6 in Tottori. Among other prefectures with comparatively high rates were Osaka (1.4), Tokyo (1.3), Kyoto (1.2) and Hokkaido (1.2). On the other hand others with low rates include Fukui (0.1), Nara (0.1) and Tochigi (0.1).

The monthly distribution of death rates was uneventful.

The usual rise in the case rates for the spring months of March (2.6), April (2.4) and May (2.0) was observed in 1949. The September rate was 3.0.

There were 1,467 cases and the case rate was 1.8 per 100,000 population. Case rates in Tokyo (4.6), Tottori (4.6), Miyagi (3.8) and Hokkaido (3.7) were comparatively high. On the other hand they were low in Tokushima (0.1), Ishikawa (0.4), Wakayama (0.4), Kumamoto (0.4) and Kagoshima (0.4).

Influenza (Int. List No. 33)

There were 502 deaths from influenza. The death rate was 0.6 per 100,000 population, the same as the all-time low rate recorded last year.

For all "shi" combined the death rate was 0.5 and for all "gun" 0.7.

Compared to the average annual death rate, the rates in Aomori (1.9), Ehime (1.5), Saga (1.5), Toyama (1.2), Kagawa (1.1) and Shimane

Public Health and Welfare in Japan - 1949

(1.1) were high. Correspondingly, rates in Ibaraki (0.0), Saitama (0.1), Miyagi (0.1) and Tottori (0.2) were low (Ref. Tables 71 and 72).

The monthly distribution of death rates was rather uneventful with a very slight increase to 1.2 in February. (Ref. Table 68)

There were 2,957 cases of influenza reported in 1949 and the case rate was 3.6 per 100,000 population. (Ref. Table 81) Prefectures having high rates included Kagawa (16.6), Fukui (16.6), Oita (16.0), Okayama (12.8) and Yamaguchi (10.8). No cases were reported in Akita or Iwate, while rates in Yamagata (0.1), Chiba (0.2) and Ibaraki (0.2) were low. (Ref. Chart A-8)

Leprosy (Int. List No. 23)

There were 162 deaths from leprosy. The death rate was 0.2 per 100,000 population, the same as that recorded since 1941. Data are not available for 1944-1946.

The death rate for all "shi" combined was 0.0 and for all "gun" 0.3.

No deaths were recorded from this disease in 10 prefectures. Especially high was the rate in Kumamoto (3.1), the location of a large leprosarium.

The monthly distribution of death rates was uneventful. (Ref. Table 16)

Leprosy cases reported in 1949 totalled 782. Of that number, 153 were reported in Gumma prefecture and in the following prefectures: 50 in Tokyo, 48 in Fukuoka, 46 in Kyoto and 35 in Kumamoto. (Ref. Table 82) Every prefecture reported at least one case.

Paratyphoid Fever (Int. List No. 2)

There were 116 deaths from this disease and the death rate was 0.1 per 100,000 population, the lowest of record.

The death rate for all "shi" combined was 0.2 and for all "gun" 0.1.

In 10 prefectures there were no deaths from paratyphoid fever. The highest death rate for any prefecture was 0.5 which was recorded in both Mie and Miyagi.

The monthly distribution of death rates was uneventful.

The seasonal distribution of case rates was somewhat similar to that for typhoid fever excepting that the monthly rates were generally one-third to one-half as great. The peak monthly rate was in August (5.1). Rates for June and July were the same (4.1) while that for September was 3.9. (Ref. Tables 77 and 78)

Cases totalled 2,226 with a case rate of 2.7 per 100,000 population. Every prefecture reported at least 5 cases. Tokyo reported

Public Health and Welfare in Japan - 1949

488 cases, Fukushima 136, Miyagi 116 and Shizuoka 103.

Scarlet Fever (Int. List No. 8)

Fifty-one deaths were recorded for scarlet fever resulting in a death rate of 0.1 per 100,000 population. (Ref. Tables 14 and 68)

The death rate for all "shi" combined was 0.1 and for all "gun" was less than 0.1.

In 18 prefectures no deaths were recorded from this disease. In 5 prefectures the rate was less than one-tenth; in 21 prefectures it was 0.1 and in 2 prefectures it was 0.2. (Ref. Tables 70 and 72)

The monthly distribution of death rates was uneventful.

There were 4,667 cases of this disease and the case rate was 5.7 per 100,000 population. Case rates higher than the national average were recorded for Tokyo (19.5), Nagano (17.9), Kyoto (14.3), Shiga (12.3), Kanagawa (10.5) and Hokkaido (10.0).

The seasonal distribution of the case rates has been quite irregular in previous years. In 1949 the peak monthly rate (7.9) was in June. The last time such a distribution was observed was in 1919.

Malaria (Int. List No. 28)

There were 122 deaths from malaria with a death rate of 0.1 per 100,000 population.

For all "shi" combined the death rate was 0.1 and for all "gun" 0.2.

All but five prefectures reported deaths from this disease. The highest rate (0.7) was recorded for both Shiga and Tokushima prefectures. Shimane and Yamaguchi prefectures each had a rate of 0.4.

The monthly distribution of death rates was uneventful (Ref. Table 16).

Cases of malaria totalled 3,732 and the case rate was 4.5 per 100,000 population. The rate for Shiga prefecture was 246.0 (Ref. Table 81).

The peak monthly case rate was 11.6 for July. The June and August rates were 10.4 and 8.2 respectively.

Rabies (Int. List No. 38b)

Seventy-six persons died from rabies in 1949 resulting in a death rate of 0.1 per 100,000 population.

The rate was the same (0.1) for all "shi" combined and all "gun".

Of the 76 deaths, 22 died in Tokyo, 19 in Chiba, 14 in Saitama, 8 in Kanagawa, 7 in Gumma, 3 in Ibaraki and 1 in Tochigi.

Public Health and Welfare in Japan - 1949

The largest number of the deaths occurred in August (12) and there were ten in both September and October.

Of the 76 cases, 23 occurred in Chiba, 18 in Tokyo-to, 14 in Saitama, 9 in Kanagawa, 6 in Gumma, 3 in Ibaraki, 1 in Tochigi and 1 in Niigata. All but 2 of the cases occurred in an area close to Tokyo (Ref. Table 82).

Typhus Fever and Typhus-like Diseases (Int. List No. 39)

Sixty-seven persons died from typhus fever and typhus-like diseases in 1949, the death rate being 0.1 per 100,000 population.

The rate for both all "shi" combined and also all "gun" was 0.1.

Of the 67 deaths the largest number (11) occurred in Okayama. The rest were widely scattered among 21 prefectures. There were no deaths recorded from this disease in 24 prefectures. The highest death rate was 0.7 in Okayama prefecture and the second highest 0.4 in Wakayama and Ehime. The monthly distribution of death rates was uneventful.

There were 121 cases reported. Of that number, 27 were reported by Osaka, 23 by Tokyo-to and 13 by Miyagi. (Ref. Table 81)

Glanders (Int. List No. 26a)

There was one death from glanders in Miyagi prefecture. This was the first death recorded from this disease since 1935. Data are not available for 1944-1946. The national death rate was less than 0.1.

Smallpox (Int. List No. 34)

These were 21 deaths from smallpox, the death rate being less than 0.1 per 100,000 population. The deaths were widely scattered among 11 prefectures (Ref. Table 81), with the most (7) occurring in Osaka.

For all "shi" combined there were 16 deaths and for all "gun" 5 deaths.

Of the 21 deaths 18 occurred during the first half of the year. The majority of deaths from this disease have usually occurred in the first half of the year (Ref. Tables 79 and 80).

There were 124 cases with a case rate of 0.2 per 100,000 population. Half of them were reported from Osaka. There were 21 cases in Fukuoka and 12 in Yamaguchi. Most of the cases were reported during the period March-June with 56 for May alone.

Yellow Fever (Int. List No. 38a)

No cases or deaths were recorded for yellow fever.

Plague (Int. List No. 3)

There were no cases or deaths recorded for plague.

Public Health and Welfare in Japan - 1949

Cholera (Int. List No. 4)

There were no cases or deaths recorded for cholera.

Anthrax (Int. List No. 7)

There were no deaths from anthrax in 1949. Of the eleven cases reported, 4 were from Toyama, 2 from Kanagawa and one each from Tokyo-to, Ishikawa, Osaka, Hyogo and Saga.

Chancroid (Int. List No. 44a)

There were no deaths attributed to this disease in 1949. There were 22,051 cases and the case rate was 26.8 per 100,000 population.

The seasonal distribution of case rates showed rates during the first half of the year were higher than those in the last half, particularly in the first quarter with a peak of 37.9 in February. (Ref. Table 78)

Prefectural case rates ranged from 3.7 in Yamagata to 87.8 in Kanagawa. Other prefectures having rates well above the national average include Kyoto (80.2), Osaka (57.1), Nara (56.1), Fukuoka (55.7), and Wakayama (54.5).

Gonorrhea (Int. List No. 25)

There were no deaths attributed to this disease in 1949. There were 181,187 cases and the case rate was 220.4 per 100,000 population.

The seasonal distribution of case rates shows the highest monthly rate in June (253.3). The rates generally decrease from June toward the end of the year with a low point (186.7) in November. Similarly, the rates decrease toward the beginning of the year with a low point (193.3) in January.

Prefectural case rates ranged from 58.3 in Niigata to 741.1 in Kanagawa. Other prefectures having rates well above the national average include Fukuoka (500.1), Yamaguchi (399.8), Hiroshima (377.2), Wakayama (345.3), Nagasaki (308.5), Tokyo (296.8) and Kyoto (296.2). Those having rates well below the average include Iwate (69.1), Ibaraki (77.3), Shimane (81.8), Saitama (85.8), Akita (87.0), Tokushima (92.7) and Yamagata (93.7)

Lymphogranuloma Venereum (Int. List No. 44a)

There were no deaths from this disease in 1949.

There were 644 cases reported and the case rate was 0.8 per 100,000 population. Miyagi and Nagano prefectures reported no cases. The highest rates were for the following prefectures: Kyoto (3.8), Osaka (2.7), Kanagawa (2.1), Hiroshima (1.9), Nara (1.8) and Hyogo (1.7). Four of the largest cities are located in these prefectures.

Public Health and Welfare in Japan - 1949

Trachoma (Int. List No. 88)

There were 176,279 cases of trachoma reported with a case rate of 214.5 per 100,000 population.

The seasonal distribution of case rates shows a high point (546.7) in June. In May it was 328.5 and in July 325.7. The January and December rates were 105.0 and 136.1, respectively.

Prefectural rates ranged from 66.2 in Yamaguchi to 720.7 in Aomori. Other prefectures having rates well above the national average include Gumma (640.4), Aichi (386.9), Miyagi (382.6), Miyazaki (364.3) and Kagawa (361.8). Those having rates well below the average include Saga (75.2), Niigata (93.2), Nagano (93.8), Shiga (95.0) and Ishikawa (95.6).

Puerperal Infection (Int. List No. 140, 142a and 147)

Deaths from puerperal infection totalled 543, the death rate being 0.7 per 100,000 population or 0.2 per 1,000 live births.

The seasonal distribution of case rates was uneventful. The highest rate was 1.4 per 100,000 population which was recorded in January, April, and October. (Ref. Table 83)

There were 986 cases of puerperal infection reported and the case rate was 1.2 per 100,000 population.

Prefectural rates ranged from zero in Nara and Osaka to 4.0 in Toyama. Other prefectures having rates above the national average include Shimane (2.9), Miyazaki (2.9), Hokkaido (2.6), Aomori (2.5) and Fukui (2.5). Those having rates below the average include Yamaguchi (0.2), Osaka (0.2), Tokyo-to (0.3), Chiba (0.4) and Mie (0.5).

Dengue Fever (Int. List No. 38f)

There were no reported deaths from this disease. There were five cases and the rate was less than 0.1 per 100,000 population. Of the five cases, two were in Fukuoka prefecture, two in Kumamoto and one in Osaka.

INFANT MORTALITY

There were 168,989 deaths of infants under one year of age in 1949. Approximately 18 out of each 100 deaths recorded were in this age group. The provisional infant death rate 62.0 per 1,000 live births was a little higher than the rate (61.0) recorded in the previous year but still the second lowest of record. (Ref. Chart A-12 and Table 3)

For all "shi" combined the rate was 55.1 and for all "gun" 65.6.

Public Health and Welfare in Japan - 1949

The monthly distribution of rates shows the lowest rate was 40.8 in September. In the two preceding years the low point was also in September. The rates decreased regularly from 76.3 in March to the low point referred to above and then increased to 82.1 in December (Ref. Tables 84 and 85). Because of the customary way in which births have been registered for December occurrences, it is likely that the January rate (60.4) is too low and the December rate too high. This is due to the use of the number of live births as the basis for computing the infant mortality rate. For further information reference is made to the paragraph titled "Births". (Ref. Chart A-13 and Tables 34 and 35) Prefectural rates ranged from 43.6 in Kanagawa to 93.1 in Toyama. Other prefectures having rates well above the national average include Aomori (89.8), Iwate (89.3), Akita (85.0), Ishikawa (82.3) and Yamagata (81.5). Those having rates well below the average include Tokyo (46.5), Kagoshima (52.0), Nagano (52.2) and Kumamoto (52.4). (Ref. Chart A-14 and Tables 36 and 37)

INFANT DEATHS FROM SELECTED CAUSES

The ten leading causes of death totalled 142,330 which is 84.2% of infant deaths from all causes. In order of their importance they were as follows: congenital debility, pneumonia, diarrhea and enteritis and ulceration of intestines, premature birth, bronchitis, other diseases peculiar to the first year of life, congenital malformations, whooping cough, measles and beriberi. (Ref. Chart A-15 and Tables 40 and 41)

Infant death rates were the lowest of record for the following causes: congenital debility, diarrhea and enteritis and ulceration of intestines, beriberi, ill-defined and unknown causes, meningitis not due to meningococcus, non-puerperal septicemia and purulent infection, tetanus, erysipelas and typhoid fever. The death rate for congenital malformations was the highest of record.

Congenital Debility (Int. List No. 158)

There were 37,285 deaths from congenital debility, the principal cause of infant deaths, and the death rate was 13.7 per 1,000 live births, the lowest of record. In 1940, the rate (26.0) was nearly double this figure. (Ref. Tables 38 and 39).

The rate for all "shi" combined was 10.7 and for all "gun" 15.2.

Monthly rates decreased from a high point (18.7) in March to a low point in September (9.0), following which they rose to 18.1 at the end of the year. The rate recorded for January was 15.0. As previously explained, infant death rates for January are considered to be higher than the recorded rate and rates of December, lower. There is no way of making corrections. However, with the change in Japanese age practices effective at the beginning of 1950, it is expected that these errors will be materially reduced. (Ref. Table 87)

Public Health and Welfare in Japan - 1949

Prefectural infant death rates ranged from 7.2 in Kanagawa to 23.9 in Akita. Other prefectures having rates well above the national average include Aomori (21.5), Yamagata (20.4), Fukui (19.9), Saga (19.5) and Shiga (19.0). Those having rates well below the average include Tokyo (8.0), Hokkaido (9.7), Yamanashi (10.4), Nagano (10.8) and Kagoshima (10.9). (Ref. Tables 89, 90 and 91)

Pneumonia, All Forms (Int. List No. 107-109)

Pneumonia, the second leading cause of infant deaths, claimed the lives of 27,724 infants. The death rate was 10.2 per 1,000 live births, the second lowest of record. In the preceding year the rate was 8.4.

For all "shi" combined the death rate was 10.7 and for all "gun" 9.9.

The monthly rate decreased from 15.4 in February to a low point of 3.3 in August after which it increased to 17.8 in December. (Ref. Tables 86 and 87)

Prefectural infant death rates ranged from 7.2 in Miyazaki and Saga to 15.6 in Iwate. Other prefectures having rates well above the national average include Osaka (12.9), Fukushima (12.8), Gumma (12.2), Aomori (12.0), Aichi (11.3), and Toyama (11.3). Those having rates well below the average include Kochi (7.5), Okayama (7.9), Kagoshima (7.9), Kumamoto (8.0) and Tottori (8.2). (Ref. Table 89)

Diarrhea, Enteritis and Ulceration of Intestines (Int. List No. 119)

The third leading cause of infant deaths was responsible for 26,815 deaths and a death rate of 9.8 per 1,000 live births, the lowest of record. (Ref. Tables 38 and 39)

For all "shi" combined the death rate was 8.0 and for all "gun" 10.8. (Ref. Tables 88 and 89)

The seasonal distribution of death rates shows an increase from 6.9 in January to a summer peak of 14.0 in July followed by a decrease to a low point of 7.5 in October. (Ref. Table 87)

Prefectural infant death rates ranged from 5.3 in Kanagawa to 18.2 in Toyama. Other prefectures having rates well above the national average include Aomori (17.5), Akita (17.0), Ishikawa (15.8), Yamagata (15.5) and Iwate (15.4). Those having rates well below the average include Tokyo (5.8), Gumma (6.3) and Kochi (6.9).

Premature Birth (Int. List No. 159)

The fourth cause of death of infants in importance was premature birth, which was responsible for 13,667 deaths. The death rate was 5.0 per 1,000 live births, the third highest of record notwithstanding the fact that a slight reduction was recorded for the second consecutive year. In 1947 it was 5.4 and in 1948, 5.1.

Public Health and Welfare in Japan - 1949

The death rate for all "shi" was 4.6 and for all "gun" 5.2.

A review of the seasonal distribution of the death rates shows a rate of 5.5 in March which decreased to 4.4 in October and then rose to 5.3 in November.

Prefectural rates ranged from 2.9 in Kanagawa and Yamaguchi to 7.8 in Toyama. Other prefectures having rates above the national average include Chiba (7.7), Ibaraki (7.7), Iwate (7.5), Kochi (7.2) and Yamagata (7.1). Those having rates below average include Hokkaido (3.1), Kagoshima (3.8), Fukuoka (3.9), Nagasaki (3.9) and Yamanashi (3.9). (Ref. Table 89)

Bronchitis (Int. List No. 106)

Bronchitis was the fifth cause in importance of infant deaths. There were 9,574 deaths with a death rate of 3.5 per 1,000 live births which was a small increase above the rate (3.2) in the preceding year. (Ref. Tables 38 and 39)

For all "shi" combined the death rate was 2.4 and for all "gun" 4.1. (Ref. Table 89)

The mortality rate from this disease decreased from 5.6 in March to 1.2 in September after which it rose to 5.6 in December. (Ref. Table 87)

Prefectural infant death rates ranged from 1.7 in Kanagawa to 7.0 in Toyama. Other prefectures having rates above the national average include Iwate (6.3), Yamagata (5.8), Ishikawa (5.4), Aomori (5.3) and Mie (5.1). Those having rates below the average include Tokyo (1.8), Kyoto (2.0), Osaka (2.4), Kumamoto (2.4) and Gumma (2.5)

Other Diseases Peculiar to the First Year of Life (Int. List No. 161)

This was the sixth leading cause of death among infants. There were 8,967 deaths and the death rate was 3.3 per 1,000 live births the same as that recorded in the preceding year.

The death rate for all "shi" combined was 2.7 and for all "gun" 3.6.

The monthly distribution of death rates of infants was uneventful.

Prefectural death rates ranged from 2.0 in Niigata to 4.8 in Shizuoka. Other prefectures having rates above the national average include Oita (4.7), Saga (4.6), Mie (4.6), Kagawa (4.5) and Gifu (4.5). Those having rates below the average include Tokyo (2.1), Kanagawa (2.5), Aomori (2.6), Fukushima (2.6), Gumma (2.6) and Yamanashi (2.6).

Congenital Malformations (Int. List No. 157)

The seventh leading cause of infant deaths was congenital malformations which was responsible for 5,255 deaths and a death rate of 1.9

Public Health and Welfare in Japan - 1949

per 1,000 live births, the highest of record.

For all "shi" combined the death rate was 1.9 and also the same figure for all "gun".

The seasonal distribution of death rates was rather uneventful, but with a slight tendency for the rates to decrease from the latter part of the first quarter of the year to a low point (1.4) in August following which they increased to the end of the year.

Prefectural death rates ranged from 1.4 in Akita to 2.7 in Gumma. Other prefectures having rates above the national average include Miyagi (2.5), Tochigi (2.5), Yamanashi (2.5), and Toyama (2.4). Those having rates below the average include Kagoshima (1.5), Shimane (1.5), Kanagawa (1.6), Kumamoto (1.6), Oita (1.6) and Tokyo (1.6). (Ref. Table 89)

Whooping Cough (Int. List No. 9)

The eighth cause of infant deaths in order of importance was whooping cough. It caused 5,021 deaths resulting in a death rate of 1.8 per 1,000 live births which was the second lowest of record since 1922. In 1948 the death rate was 0.9. (Ref. Tables 38 and 39)

For all "shi" combined the death rate was 2.1 and for all "gun" 1.7. (Ref. Table 89)

The death rate rose from 0.9 in January to 2.8 in June at which level it remained until August, after which it decreased to 1.5 in November. (Ref. Table 87)

Prefectural rates ranged from 0.6 in Kochi, Miyazaki and Wakayama to 3.4 in Yamagata. Other prefectures having rates above the national average include Gumma (2.9), Shiga (2.6) and Miyagi (2.5). Those having rates below the average include Kagoshima (0.7), Tokushima (0.7), Ibaraki (0.9), Kumamoto (0.9), Nagasaki (1.0) and Shimane (1.0).

Measles (Int. List No. 35)

Measles was the ninth leading cause of death among infants with 4,508 deaths. The death rate was 1.7 per 1,000 live births. The year 1949 was on the upper end of the regular 2-year cycle for measles and it was to be expected that there would be a temporary rise in the rate. (Ref. Tables 38 and 39)

The death rate for all "shi" combined was 2.0 and for all "gun" 1.5. (Ref. Table 89)

The monthly death rate increased from 0.6 in January to 5.7 in May after which it declined to 0.2 in October. (Ref. Table 87)

Prefectural rates ranged from 0.3 in Kochi and Okayama to 3.9 in Nara. Other prefectures having rates above the national average

Public Health and Welfare in Japan - 1949

include Osaka (3.3), Hokkaido (2.9), Toyama (2.9), Fukuoka (2.7) and Niigata (2.7). Those having rates below the average include Kagawa (0.4), Tokushima (0.4), Tochigi (0.5), Miyazaki (0.6) and Tottori (0.7).

Beriberi (Int. List No. 68)

Beriberi was tenth in importance as a cause of infant deaths, being responsible for 3,514 deaths. The death rate was 1.3 per 1,000 live births, the lowest of record.

The death rate for all "shi" was 1.1 and for all "gun" 1.4.

The monthly death rate increased from 1.3 in January to 1.8 in March, following which it decreased to 0.7 in September and then increased to 1.7 in December.

Prefectural death rates ranged from 0.2 in Kagawa and Yamanashi to 3.1 in Aomori and Iwate. Other prefectures having rates above the national average include Ishikawa (2.8), Yamagata (2.4), Saga (2.3), Tottori (2.0) and Toyama (2.0). Those having rates below the average include Ehime (0.5), Tokushima (0.6), Gumma (0.7), Kanagawa (0.7) and Nagano (0.7).

Ill-Defined and Unknown Causes (Int. List No. 199-200)

There were 2,587 deaths from this cause with a death rate of 0.9 per 1,000 live births, the lowest of record. The trend in the rate has been gradually downward for many years. (Ref. Tables 38 and 39)

The death rate for all "shi" combined was 0.8 and for all "gun" 1.0. (Ref. Table 89)

Death rates were somewhat higher during the first four months and last two months of the year. (Ref. Table 87)

Prefectural death rates ranged from 0.4 in Kumamoto and Tottori to 2.0 in Aomori. Prefectures having rates above the national average include Fukui (1.8), Nara (1.7), Hokkaido (1.6), Toyama (1.5) and Iwate (1.4). Those having rates below the average include Hiroshima (0.5), Nagasaki (0.5), Fukuoka (0.6), Kagoshima (0.6), Kyoto (0.6) and Miyazaki (0.6).

Meningitis, Not Due to Meningococcus (Int. List No. 81)

There were 2,543 deaths of infants from this cause and the death rate, 0.9 per 1,000 live births was the lowest of record.

The death rates for all "shi" combined and all "gun" were both 0.9.

Death rates for April and May were 1.2. The average rate for the second quarter of the year was a little higher than that for other quarters, whereas that for the third quarter was a little lower.

Public Health and Welfare in Japan - 1949

Prefectural death rates ranged from 0.5 in Nagano, Tokyo and Yamaguchi to 1.6 in Akita. Among the other prefectures having rates above the national average were Ishikawa (1.4), Aichi (1.3) and Nara (1.3). Those having rates below the average include Gumma (0.6), Ibaraki (0.6) and Tochigi (0.6). (Ref. Table 89)

Accidental Deaths (Int. List No. 169-195)

There were 1,931 deaths from accidental causes. The death rate was 0.7 per 1,000 live births.

The death rate was the same for all "shi" and all "gun", 0.7. Monthly distribution of deaths was uneventful.

Prefectural death rates ranged from 0.3 in Chiba and Saitama to 1.5 in Shimane. Other prefectures having rates above average include Fukui (1.2), Kagoshima (1.1), Fukushima (1.0), Iwate (1.0) and Nagasaki (1.0). Those having rates below average include Ibaraki (0.4), Kagawa (0.4) and Nara (0.4).

Convulsions (Int. List No. 86)

Deaths of infants from convulsions totalled 1,409 and the death rate, 0.5 per 1,000 live births, equalled the lowest of record. During the 3-year period 1947-1949, the rate has remained the same. (Ref. Tables 38 and 39).

The death rate for all "shi" combined was 0.2 and for all "gun" 0.7. (Ref. Table 89)

The monthly distribution of death rates was uneventful. The lowest rate was 0.3 recorded in both August and September.

Prefectural death rates ranged from less than 0.1 in Miyazaki and Nara to 6.0 in Toyama. Other prefectures having rates well above the national average include Ishikawa (5.5), Aomori (4.0), Akita (3.1) and Hokkaido (1.2). Those having rates well below average include Aichi, Ehime, Fukuoka, Fukushima, Gifu, Gumma, Hyogo, Kagoshima, Kanagawa, Kochi, Kumamoto, Okayama, Osaka, Saga, Saitama, Shizuoka and Tokyo, all of which had a rate of 0.1.

Tuberculosis, All Forms (Int. List No. 13-22)

Tuberculosis caused the deaths of 1,332 infants resulting in a death rate of 0.5 per 1,000 live births equaling the second lowest of record. Ever since 1929 the rate has been either 0.4 or 0.5 and most frequently the latter. (Ref. Tables 38 and 39)

The death rate for all "shi" combined was 0.8 and for all "gun" 0.3. (Ref. Table 89)

The distribution of the monthly death rates was uneventful although, as in the preceding year, there was a slight tendency for the

infant mortality rates to rise from 0.3 in January to 0.8 in May. In June and July the rate was 0.7 following which it decreased to 0.4 in August at which level it remained for four months. (Ref. Table 87)

Prefectural death rates ranged from less than 0.1 in Yamanashi to 1.0 in Osaka. Other prefectures having rates above the national average include Hokkaido (0.9), Kyoto (0.8), Tokyo (0.8), Fukuoka (0.7), Hyogo (0.7) and Shimane (0.7). Those having rates below the average include Gumma (0.1), Ibaraki (0.1), Tochigi (0.1), Chiba (0.2), Fukui (0.2), Fukushima (0.2), Nagano (0.2), Saitama (0.2) and Shiga (0.2).

Syphilis (Int. List No. 30)

Syphilis caused the death of 1,146 infants. The death rate was 0.4 per 1,000 live births equalling the all-time low point reached in 1947 and also the rate for 1948.

For all "shi" combined the death rate was 0.6 and for all "gun" 0.3.

Except for a very slight rise in the death rate to 0.5 in March and 0.6 in April the monthly distribution of rates was uneventful.

Prefectural death rates ranged from 0.1 in Nagano to 1.0 in Fukuoka. Other prefectures having rates above the national average include Kochi (0.9), Nara (0.8), Kyoto (0.7), Miyazaki (0.7) and Osaka (0.7). Those having rates below average included Ehime, Gifu, Ishikawa, Niigata, Shimane, Tokushima, Wakayama and Yamanashi all of which had a rate of 0.2. (Ref. Table 89)

Injury at Birth (Int. List No. 160)

Deaths of infants caused by injury at birth totalled 1,062. The death rate was 0.4 per 1,000 live births, which equalled the highest point of record established in the preceding year. (Ref. Tables 38 and 39)

The death rate for all "shi" combined was 0.4, the same as that for all "gun".

The seasonal distribution of death rate was uneventful.

Prefectural death rates ranged from 0.2 in Aomori and Yamanashi to 0.8 in Kochi and Toyama. Other prefectures having rates above the national average include Kyoto (0.7), Yamagata (0.7), Miyagi (0.6) and Chiba (0.6). Sixteen prefectures had a rate of 0.3. (Ref. Table 89)

Non-Puerperal Septicemia and Purulent Infection (Int. List No. 24)

There were 898 deaths of infants from this cause and the death rate was 0.3 per 1,000 live births, the lowest of record and one-third the rate recorded in 1940. (Ref. Tables 38 and 39)

Public Health and Welfare in Japan - 1949

For all "shi" combined the death rate was the same as that for all "gun", 0.3.

The monthly distribution of the death rates was uneventful.

Prefectural rates ranged from 0.2 in nine prefectures to 0.7 in Oita. The death rate was 0.6 in Kagoshima and 0.5 in Fukuoka, Ishikawa, Nagasaki and Shimane.

Tetanus (Int. List No. 12)

Deaths of infants from tetanus totalled 868. The death rate was 0.3 per 1,000 live births, the lowest of record.

The death rate for all "shi" combined was 0.2 and for all "gun" 0.4.

As in the preceding year, there was a slight increase in the death rates from this disease during the summer months. (Ref. Table 87)

Prefectural death rates ranged from 0.1 in Fukui, Hyogo, Kyoto, Nara, Osaka, Tokyo and Tottori to 0.9 in Chiba and Ibaraki. In Saga prefecture the rate was 0.8 and in Gumma, Kagawa and Yamanashi, 0.7.

Erysipelas (Int. List No. 11)

There were 770 deaths from this disease and the death rate, 0.3, was the lowest of record.

The death rate for all "shi" combined was 0.2 and for all "gun" 0.3.

The seasonal distribution of monthly death rates was uneventful.

Prefectural death rates ranged from 0.1 to 0.5. Prefectures having rates above average included Kagoshima (0.5), Nara (0.5) and Shiga (0.5). The rate was 0.1 in Kanagawa and Tokyo.

Diphtheria (Int. List No. 10)

Diphtheria was responsible for 212 deaths of infants and the death rate was 0.1 per 1,000 live births, which has been recorded many times in past years and is the lowest point which the rate for the disease has reached. (Ref. Tables 38 and 39)

For all "shi" combined and for all "gun" the death rate was 0.1. (Ref. Table 89)

Monthly death rates in June, August, September and October were less than 0.1, slightly lower than the rate of 0.1 recorded for other months. (Ref. Table 87)

Public Health and Welfare in Japan - 1949

Prefectural death rates ranged from less than 0.1 in thirteen prefectures to 0.2 in Miyazaki, Oita, Okayama and Tottori.

Influenza (Int. List No. 33)

Deaths of infants from influenza totalled 165. The death rate was 0.1 per 1,000 live births, the second lowest of record. It was less than 0.1 in the preceding year. (Ref. Tables 38 and 39)

The death rate for all "shi" combined and that for all "gun" were the same (0.1).

Monthly death rates were slightly lower in the Summer and Fall than at other times during the year. (Ref. Table 87)

Prefectural death rates ranged from zero (no deaths) to 0.2. No deaths were recorded in Ibaraki, Kochi, Miyagi, Shiga and Tottori. The rate was 0.2 in Aomori, Fukui and Shimane. (Ref. Table 89)

Dysentery (Int. List No. 27)

Dysentery caused 113 deaths of infants. The death rate was less than 0.1 equal to the lowest of record. Prior to 1948 the rate remained at a level of 0.1. (Ref. Tables 38 and 39)

For all "shi" combined and for all "gun" the rates were the same, less than 0.1.

The monthly death rates increased to 0.1 during the summer months (June-September). (Ref. Table 87)

Prefectural death rates ranged from zero (no deaths) to 0.1. No deaths were recorded in Kochi, Kumamoto, Kyoto, Miyagi, Shiga and Wakayama. (Ref. Table 89)

Cerebrospinal (Meningococcus) Meningitis (Int. List No. 6)

There were 81 deaths from this cause with a death rate less than 0.1. It has remained at this level from the time data were available in 1920.

The death rate for all "shi" combined was 0.1 and for all "gun" less than 0.1.

The monthly distribution of death rates was uneventful. (Ref. Table 87).

Prefectural death rates ranged from zero (no deaths) in twenty-one prefectures to 0.1 in eight prefectures.

Public Health and Welfare in Japan - 1949

Japanese "B" Encephalitis (Int. List No. 37b)

There were eight deaths of infants attributed to this disease and the death rate was less than 0.1 both in 1949 and 1948, which are the only years for which data are available.

The death rates for all "shi" and all "gun" were equal (less than 0.1).

No deaths from this disease were recorded January-May, July, November and December. Of the eight deaths recorded during 1949, one occurred in June, one in August, four in September and two in October.

Prefectural death rates ranged from zero (no deaths) in 38 prefectures to less than 0.1 in all others.

Scarlet Fever (Int. List No. 8)

There were five deaths recorded from scarlet fever in 1949 and the death rate was less than 0.1 per 1,000 live births the same as it has been for many years. (Ref. Tables 38 and 39)

For both all "shi" combined and all "gun" the death rate was less than 0.1. (Ref. Table 89)

Of the five deaths, one each was recorded in January, February, April, May and July. No deaths were recorded from this disease for the last five months of the year. All monthly death rates were considerably below 0.1 per 1,000 live births. (Ref. Tables 86 and 87)

Four prefectures recorded deaths. They were Aomori, Hokkaido, Okayama and Osaka. All had one death except Hokkaido which had two deaths. All rates were well below 0.1 per 1,000 live births.

Typhus Fever and Typhus-Like Diseases (Int. List No. 39)

Only two deaths were recorded for infants from these diseases and the death rate was considerably less than 0.1 per 1,000 live births. Data are not available for 1944-1946. During the quarter of a century which has elapsed since 1925, deaths have been recorded in only five. They are 1935 (1), 1940 (1), 1941 (1), 1943 (2) and the present year (2). (Ref. Tables 38 and 39).

The death rate was less than 0.1 per 1,000 live births for both all "shi" combined and all "gun".

One death occurred in January and the other in February. (Ref. Table 86)

Typhoid Fever (Int. List No. 1)

No deaths of infants were recorded for typhoid fever. This is the first time this has happened according to data collected beginning

Public Health and Welfare in Japan - 1949

with 1920. In previous years the annual number of deaths recorded has ranged from 5 to 32 but the death rate has remained at the level of less than 0.1 per 1,000 live births.

Paratyphoid Fever (Int. List No. 2)

There were no deaths of infants from paratyphoid fever. This is not unusual as a similar record has been observed 10 times in the last quarter of a century. Over the same period the number of deaths has usually been one and never more than two.

Cholera (Int. List No. 4)

No infants died from cholera in 1949. The last record of death from this disease was in 1940 when two deaths occurred. Prior to that date the last reported deaths were in 1930 when two were recorded.

Anthrax (Int. List No. 7)

There were no deaths of infants from anthrax. The last record of such an occurrence was one death in 1947. During the past quarter of a century not more than one death was recorded in any single year. Other than the death referred to above they occurred in 1940, 1937, 1933, 1932, 1930, 1928 and 1926.

Leprosy (Int. List No. 23)

There were no deaths of infants from leprosy. Very few deaths of infants have been recorded from this disease. According to data collected beginning with 1920, deaths were recorded in only three years. In 1920, 1924 and 1943 there was one death each and in 1940, three deaths were recorded.

Glanders (Int. List No. 26a)

There were no deaths of infants from glanders. In only three years since 1920 has there been a death from this disease. One death was recorded in 1921, 1923 and 1935.

Malaria (Int. List No. 28)

No deaths of infants from malaria were recorded in 1949. In only two other years since 1920 has there been a similar record. The number of deaths has ranged up to 15 deaths in any single year (1921 and 1923). The death rate in years in which deaths have been recorded has always remained at the same level -- less than 0.1 per 1,000 live births.

Public Health and Welfare in Japan - 1949

Yellow Fever (Int. List No. 38a)

There were no deaths of infants from yellow fever. Data are not available for 1944-1946. Beginning with 1923 there have been only two years in which infants have died. They are 1923 and 1933 in which one death each was recorded.

Rabies (Int. List No. 38b)

No deaths of infants were recorded from rabies in 1949. Data are not available for 1944-1946. Since 1920 there have been only four years in which deaths of infants have been recorded from this disease. There were two deaths in 1923, 26 in 1924, 3 in 1925 and 1 in 1935.

STILLBIRTHS

Stillbirths totalled 192,970 and the provisional stillbirth rate (70.8 per 1,000 live births) was the highest of record in more than a quarter of a century (Ref. Chart A-12, A-16). The 1949 rate was more than 70% above the 1943 figure. (Ref. Table 4)

The death rate for all "shi" combined was 95.7 and for all "gun" 58.0.

The death rate was highest in August (82.2) with a secondary peak (79.7) in May. A review of the seasonal distribution of stillbirth rates shows the 1949 distribution to be unusual. Excluding rates for January and December because of the effect which the registration practices (as described in the paragraph "Births") have on their computation, some interesting observations may be made. During the decade 1920-1929 the peak rate was in April and in the following decade (1930-1939) generally in June. Data are not available by month for 1942-1945, but during 1940-1949 the peak was most often in May. (Ref. Chart A-17 and Tables 42 and 43)

Prefectural death rates ranged from 56.0 in Aomori to 94.8 in Tottori. Other prefectures having rates well above the national average include Nagano (89.7), Okayama (88.0), Hyogo (84.4), Osaka (84.3), Shimane (81.7), Kyoto (81.5), and Gumma (80.5). Those having rates well below the average include Hokkaido (58.8), Tochigi (60.0), Chiba (60.9), Toyama (60.9) and Kochi (61.4) (Ref. Tables 44 and 45). It is striking that six of the prefectures having very high stillbirth rates are located close together on the southern end of the island of Honshu. Those having very low rates are widely scattered geographically. (Ref. Tables 62, 92 and 93) Note: All stillbirth data are for stillbirths after the third month of uterogestation. There were 141,663 stillbirths after the fifth month of uterogestation in 1949, or 52.0 per 1,000 live births.

MARRIAGES

There were 846,014 marriages and the marriage rate was 10.3 per 1,000 population. The sharp rise in the marriage rate during 1947 and 1948 has ended with a decrease back to the level of 1943. (Ref. Tables 4, 46, 47, 48 and 62)

The marriage rate for all "shi" combined was 9.8 and for all "gun" 10.5. (Ref. Tables 94 and 95)

The seasonal distribution of marriage rates for the 3-year period 1947-1949 has been quite different from that recorded in earlier years. In 1947 there was a sharp rise in May (16.9) and the December rate (14.1) was also much higher than usual. In 1948 the usual decline in April did not occur and the rate (15.7) remained the same. In 1949 the January rate (11.9) was the highest of record for that month. Although the rate increased to 14.6 in February, it was a little below the rate (14.9) for the same month in the previous year. The usual increase in the rate from February to March did not occur. Instead the rate decreased to 12.9 in March. From then until the end of the year the seasonal pattern of the rates was rather normal. Except for the January rate all other monthly rates showed marked reductions compared to the 1947 and 1948 figures. It is expected that this will be reflected in a substantially reduced birth rate in 1950. (Ref. Charts A-18, A-19 and Table 47).

Prefectural marriage rates ranged from 8.9 in Kyoto and Tokyo to 12.3 in Yamagata. Other prefectures having rates well above the national average include Saga (11.7), Fukushima (11.4), Fukui (11.2), Iwate (11.2) and Nagasaki (11.2). Those having rates well below the average include Kanagawa (9.0), Chiba (9.2), Aichi (9.5) and Yamanashi (9.5). (Ref. Chart A-20 and Tables 49 and 95)

DIVORCES

Divorces totalled 82,872 and the provisional divorce rate was 1.0 per 1,000 population the same as that in each of the two preceding years and equal to the highest of record. (Ref. Chart A-21 and Tables 50 and 51)

For all "shi" combined the rate was 1.1 and for all "gun" 1.0. (Ref. Tables 96 and 97)

The seasonal distribution of rates for 1949 was rather similar to that of the preceding year except that the January rate (0.9) increased and the November rate (0.9) decreased. The monthly distribution of rates was uneventful. (Ref. Chart A-22)

Prefectural rates ranged from 0.7 to 1.4. Those having rates above the national average include Akita (1.4), Kagawa (1.3), Kochi (1.3) and Nagasaki (1.3). Those having rates below the average include Chiba (0.7), Ibaraki (0.7), Nagano (0.7) and Saitama (0.7). (Ref. Tables 52 and 53)

Public Health and Welfare in Japan - 1949

HOSPITAL STRENGTH

Number of Hospitals

In January 1946, the first month for which the number of civilian hospitals in Japan is available, there were 2,940 hospitals of 10 beds or more, not including tuberculosis sanatoria and leprosaria. The number rose generally each month to a peak of 3,483 in April 1948 and decreased slightly the next month to 3,462. In June 1948 the basis for counting hospitals was changed to include tuberculosis sanatoria and leprosaria of 10 beds or more (3,746). The apparent increase of 284 hospitals is believed to consist almost entirely of sanatoria and leprosaria. Under the new reporting system the number of hospitals continued to rise, slightly but continuously, to a peak of 3,828 in October of the same year. In November 1948 there was a change in the legal definition of "hospital" which established 20 beds as the minimum size of places entitled to be called hospitals. For that month 2,859 hospitals of 20 beds or more were reported. The decrease of nearly a thousand in the number of hospitals is believed to be due almost exclusively to elimination of hospitals of 10 or more beds but less than 20. From December 1948 (2,852) through December 1949 (3,136) the number of hospitals steadily increased.

Statistics on the number of hospitals can be subdivided into four categories: tuberculosis sanatoria, mental hospitals, leprosaria and all other hospitals. (Ref. Table 98) There was a small increase in the number of tuberculosis sanatoria during 1949 from 290 in January to 294 in December. Of the 294 sanatoria, half of them were located in nine prefectures as follows: Tokyo (30), Fukuoka (23), Hyogo (21), Hokkaido (15), Chiba (14), Kanagawa (13), Osaka (11), Niigata (10) and Aichi (10). Over 60 were located in the Tokyo area and vicinity.

The number of mental hospitals remained practically the same in December as in January (123). Mental hospitals were widely distributed, there being one or more in each prefecture except in Wakayama and Miyazaki where there were none. The greatest number (12) was in Tokyo. The number of leprosaria (13) did not change during the year. They were located in 10 prefectures. There was one in each of the following: Aomori, Miyagi, Gumma, Tokyo, Yamaneeni, Kagawa and Kagoshima. There were two each in Shizuoka, Okayama and Kumamoto. The number of general hospitals rose from 2,447 in January to 2,706 in December. Of the 2,706, more than one-third were located in seven prefectures as follows: Tokyo (222), Hokkaido (213), Osaka (129), Aichi (119), Fukuoka (110), Kanagawa (102) and Hyogo (101).

Prefectural hospital statistics show that the total number of hospitals increased in 35 prefectures, remained about the same in five and decreased slightly in the remaining five. The greatest increase was in Hokkaido, with 42 more hospitals at the end of the year than at the beginning. The greatest decrease was in Fukushima with four less.

Bed Capacity

The daily average bed capacity in civilian hospitals in Japan (i.e. the rated capacity which may in some cases be less than the

Public Health and Welfare in Japan - 1949

number of in-patients) has shown a less regular increase than the number of hospitals. There were 201,263 beds available daily in January 1946. The number increased through June (208,768) and in July there was a negligible decrease to 208,737. Thereafter the number rose each month to a peak of 222,498 in May 1947, after which it decreased regularly through November (211,659). In December (211,903) there was a slight rise, but subsequent months showed a decline again through March 1948, at which time the number of beds available had decreased to 208,290. April and May showed increases so that the daily average bed capacity in May was 210,452. The sudden rise in June 1948 to 258,020 reflects the addition of sanatoria and leprosaria to the hospital tabulations. The upward movement continued through October (261,374). The decrease in November 1948 (246,273) is believed to be almost entirely due to the elimination of hospitals of less than 20 beds from the tabulations. The number decreased to 243,478 in February 1949, after which the trend turned upward reaching a bed capacity of 253,506 in December.

In January 1949 the bed capacity of tuberculosis sanatoria (Ref. Table 98) was 53,350. Thereafter it increased generally to 56,150 in December. If the beds available in tuberculosis wards of general hospitals are included, there was a rise in the number of tuberculosis beds each month, from 73,200 in January to 82,069 in December. The bed capacity for mental hospitals was 15,377 in January. It decreased slightly to its lowest point in April (15,210), rising generally thereafter to a December total of 15,928. The total beds available for mental patients, including those in general hospitals, followed much the same pattern. In January it was 16,883; in April, 16,726; and in December, 17,665. The seasonal distribution of beds available in leprosaria followed a less regular pattern. The highest daily average capacity was 9,169 in February and lowest in October (9,011). Beds available in all other hospitals (exclusive of mental hospitals, leprosaria and sanatoria) totalled 165,956 in January, decreased slightly to 165,758 in February and then increased generally to 172,390 in December.

The percent of beds in general hospitals reserved for tuberculosis patients rose during the year. In January tuberculosis wards contained 19,850 beds or 12% of the total (165,956). In December there were 25,919 tuberculosis beds, or 16% of the total (172,390). The number of ward beds for mental patients rose by a smaller amount from 1,506 in January to 1,737 in December and the percent of total beds in general hospitals remained about one percent each month.

A review of the data shows that 34 prefectures had greater hospital bed capacities in December than in January, 11 had smaller and one had the same number. The greatest increase was in Osaka Prefecture, from 14,950 beds available in January to 17,867 in December. The greatest decrease was in Tokyo-to from 30,555 to 28,960.

Total Patient Load

The daily average number of all patients in civilian hospitals, including both in-patients and out-patients but excluding those in sanatoria and leprosaria, was 310,809 in January 1946. The number

Public Health and Welfare in Japan - 1949

increased each month until September at which time it had reached 419,983. From October through January 1947 the number decreased to 353,643. Thereafter it rose to 44,062 in April. During the period May-September it increased irregularly to 444,472. As in the previous year, there was a period of decrease between October and January 1948. A low point of 333,767 was reached. Then the number of hospital patients generally increased and the addition of leprosaria and sanatoria to hospital tabulations in June 1948 had no marked effect upon the upward trend. The seasonal pattern of the two previous years was repeated with a peak in September of 465,826 patients, followed by a decrease in October to 444,862. The November decrease (392,927) is believed to be largely due to the elimination of hospitals of less than 20 beds from the tabulations. Part of the decrease may be attributed to the normal seasonal trend. In January 1949 there were 368,628 hospital patients. The high point of the year was reached in August with 521,104 patients, following which the number decreased each month to a December total of 443,466.

The number of patients in tuberculosis sanatoria (Ref. Table 98) increased with some monthly fluctuations from 40,442 in January 1949 to a high point of 51,974 in October. Decreases were recorded in the last two months of the year, ending with 50,622 in December. The number of patients in mental hospitals in January was 11,366. The high point of the year was reached in September with 13,741. By December the total had decreased to 13,237. The number of patients in leprosaria varied from month to month, but the trend was generally upward with 8,072 in January and 8,440 in December. For all other hospitals (exclusive of sanatoria, leprosaria and mental hospitals) total patients in January numbered 308,748. An increase was recorded each month until a high point was reached in August (451,605). After that the number decreased through December (371,167).

The total number of hospital patients in January 1949 was lower than in December in all but two prefectures, Hokkaido and Aomori. The greatest increase was in Fukuoka prefecture.

In-Patient Load

The daily average number of in-patients for hospitals of 10 beds or more, excluding sanatoria and leprosaria, was 91,949 in January 1946. "In-patients" include all patients who spent at least one night in the hospital. The number rose each month until August (116,646) after which it decreased regularly until January (96,873). It rose to 110,494 in July, decreased in August and returned to 110,400 in September. The usual seasonal downward trend was recorded from October 1947 to January 1948 with a low point of 88,499 in the latter month. From February to May the number increased reaching 104,848 in May. The sharp rise to 132,881 in June is believed, to a very large extent, to be due to the addition of sanatoria and leprosaria to the tabulations. The upward trend continued, reaching a high point of 143,397 in September. There was a decrease in October to 141,280. The marked reduction in November to 132,480 largely resulted from the elimination of hospitals of less than 20 beds from the tabulations.

Public Health and Welfare in Japan - 1949

The usual downward seasonal trend followed with a low point being reached in January 1949 of 130,364. After January the number increased regularly, reaching a high point of 177,074 in September. A downward trend was then observed to 165,322 in December.

In-patients in tuberculosis sanatoria (Ref. Table 98) numbered 36,609 in January 1949. The trend was upward through October (47,773). In the last two months it was downward reaching 47,131 in December. The total number of tuberculosis in-patients, including those in general hospitals, followed the same monthly pattern. The lowest number for the year was in January (52,065) and the highest in October (73,007). The number of in-patients in mental hospitals rose from January (10,868) through July, decreased slightly in August, and rose again in September, remaining at approximately 13,000 for the remainder of the year. The total for mental patients, including those in general hospitals, also increased from a low point in January (11,953) through July, declined in August, and rose again in September (14,677). The number in the three months following showed slight successive decreases. The number of patients in leprosaria rose irregularly from 8,059 in January to its highest point (8,487) for the year in November, following which it decreased slightly to 8,432 in December. The number of patients in all other hospitals (exclusive of sanatoria, leprosaria and mental hospitals) increased each month from January (74,828) through September (108,319) while in each of the three following months it decreased.

In January 1949 there were 15,456 tuberculosis patients in general hospitals, 21% of the total number of patients in them. Correspondingly, by December the number had risen to 24,610 or 25% of the total (96,936). Of the 52,065 in-patients with tuberculosis in January, both in sanatoria and general hospitals, approximately 30% were in general hospitals. In December the ratio had risen to almost 35%. The average for the year was about 33%. Similarly, the number of mental patients in general hospitals rose from 1,085 to 1,397, representing a little more than 1% of the total in each month of all patients in general hospitals. Approximately 10% of all mental patients were cared for in general hospitals.

All prefectures except Aomori and Tochigi reported more in-patients in December than in January. In Aomori there were 3,411 in-patients in January and 2,728 in December. The decrease in Tochigi was negligible. Greatest increases were in Tokyo-to (18,064 to 21,356), Hokkaido (6,796 to 9,623) and Osaka Prefecture (7,397 to 10,005).

Out-Patient Load

The daily average number of out-patients in hospitals of 10 beds or more, exclusive of sanatoria and leprosaria, was 228,860 in January 1946. It includes both out-patients and patients visited in their homes by physicians on the hospital staffs. The number varied from month to month in a similar way to that observed for the in-patients, rising generally from February through September and decreasing from October through January of the following year. However, the number of out-patients in June 1948 (291,304), the first month for which sanatoria and leprosaria were included in the tabulations, was less than in the

Public Health and Welfare in Japan - 1949

preceding month (316,070). There was a marked drop in November 1948 (260,447) from October (303,583). This was partly due to the exclusion of hospitals of less than 20 beds from the tabulations. Although the trend in the number of out-patients was upward during the first nine months in 1949, unlike the previous three years the high point was reached one month earlier (August) with 354,077, after which the number decreased regularly to the end of the year.

The daily average number of out-patients of tuberculosis sanatoria in 1949 rose irregularly from January (3,833) to July (4,913) and decreased each month thereafter, reaching 3,491 in December. In January the number of out-patients of mental hospitals was 498, at which level it remained generally through June. It decreased to 399 in July practically equalling the November figure (398). By December the number was 414. The number of out-patients of leprosanatoria is small with a daily average ranging from 7 to 23 during the 12 months of 1949. Out-patients of all other hospitals (exclusive of sanatoria, leprosanatoria and mental hospitals) averaged 233,920 daily in January. The number increased each month through April, decreased slightly in May, but rose again to a peak in August (348,673), with the last four months showing successive decreases.

The daily average number of out-patients was greater in December than in January in 41 prefectures and smaller in five. The greatest increase was in Fukuoka Prefecture, with 24,425 out-patients in January and 30,810 in December. Osaka Prefecture reported an increase in the daily average of more than 4,000 in December compared to January. The greatest decrease occurred in Hokkaido, with 29,028 in January and 25,038 in December.

Bed Occupancy

The bed occupancy ratio, the number of beds occupied per 100 beds of rated capacity, was 40.7 in January 1946. The percent rose through August to 55.4 and then decreased to 44.7 in January 1947. The highest percent in 1947 was reached in September (51.5) after which, following the usual seasonal pattern, it decreased to 41.8 in January 1948. In September it reached 55.0, after which it decreased to 53.5 in January 1949. Repeating the usual seasonal trend, it rose to a high point of 70.3 in September, subsequently decreasing to 65.2 in December.

The bed occupancy for tuberculosis sanatoria in January 1949 was 68.6%. The percent in each succeeding month (except August) showed a higher ratio through October, when it reached a high point of 86.3, decreasing a little in November and December (83.9). Bed occupancy percents for all tuberculosis patients, including those in general hospitals, were higher than those for sanatoria but followed approximately the same seasonal pattern. In January the percent was 71.1 but the high point for the year was reached in September (90.9), practically equalling the October figure (90.8). By December the percent had decreased to 87.4. The bed occupancy percent for mental hospitals was 70.7 in January, following which it rose each month (except August) to a high point of 84.6 in September. Then it decreased each month to 80.5 in December. The occupancy percents of beds for all mental patients, including those in general hospitals, were approximately the same as for patients in mental hospitals alone. The January percent

Public Health and Welfare in Japan - 1949

was 70.8. It rose irregularly to a high point (84.5) in September, decreasing thereafter to 80.5 in December. Bed occupancy percents in leprosaria were irregular during 1949, but during the second half of the year they were higher than during the first half. The percent in January was 88.4. The high point of the year was in November (93.9), and the December figure (93.3) was only a little lower. The percent of beds occupied in general hospitals rose steadily from 45.1 in January to 62.9 in September, and then declined to 56.2 in December.

Over half (24) of the prefectures reported bed occupancy percents for 1949 within a range of 10% plus or minus the annual average for all Japan (63.6). Thirteen prefectures had percents which were below this range and nine above. The percent in Yamanashi Prefecture (48.5) was about one-fourth below the national average, while in Shimane the highest percent (74.1) was about one-sixth above. All but two prefectures reported higher percents in December than in January. Aomori Prefecture reported 92.4 in January and 72.2 in December. The greatest increases occurred in Nagasaki (32.3 to 54.8), Fukui (45.5 to 68.8), and Hyogo (44.7 to 66.8).

About one-third (15) of the prefectures reported bed-occupancy ratios for tuberculosis sanatoria within 10% of the national average (79.1). Twenty were above this range and 11 below. Yamanashi Prefecture had the highest ratio (106.5), over 30% above the national figure. The excess (6.5) over 100% indicates that more patients were admitted than the rated capacity of the hospitals. Five additional prefectures (Shimane, Wakayama, Miyazaki, Miyagi, and Kagawa) had percents over 20% above. Fukushima Prefecture had a rate (56.3) nearly 30% below the national average. Four additional prefectures (Ishikawa, Akita, Osaka and Aichi) had rates more than 20% below. Ratios were higher in December than in January in all but four prefectures. The ratio in Kagawa Prefecture decreased from 100.0 in January to 94.0 in December, while lesser decreases were recorded in Toyama, Nagasaki, and Tokyo. The greatest increase occurred in Nara (45.8 to 89.0), Hyogo (43.8 to 86.1), and Saga (57.2 to 99.0).

About one-third (15) of the prefectures had bed-occupancy ratios for mental hospitals within a range of 10% of the national average (79.4). There were 18 with ratios above this range and 11 below. Two prefectures (Wakayama and Miyazaki) have no mental hospitals. The bed-occupancy ratio in Iwate Prefecture (160.0) was double the national average, and ratios in Shimane and Toyama were more than 50% above. Ratios in Aomori (46.5) and Oita (47.7) were each about 40% below the national figure. Percents were higher in December than in January in 35 prefectures, lower in 8, and the same both months in one. The greatest increase in percents of occupancy occurred in Iwate Prefecture, from 106.7 in January to 178.3 in December, (above rated capacity each month). There was a decrease in Shimane, from 153.1 to 110.5. Bed-occupancy percents in the ten prefectures having leprosaria ranged from 64.6 in Yamanashi, about 30% below the national rate (90.5), to 107.6 in Shizuoka, almost 20% above. Ratios were higher in December than in January in 6 prefectures and lower in 4. The greatest increase (72.6 to 91.6) occurred in Miyagi Prefecture and the greatest decrease (107.0 to 82.0) in Shizuoka.

Public Health and Welfare in Japan - 1949

Nearly half (21) of the prefectures reported bed occupancy percents for general hospitals within a range of 10% plus or minus the national average (55.8). Fifteen were lower, and ten were higher. The ratio in Yamanashi Prefecture (37.0) was one-third below the national figure, and in Fukuoka (67.1) one-fifth above. All but four prefectures reported higher occupancy percents in December than in January. Aomori Prefecture reported a January ratio of 99.6 and a December ratio of 61.1, while lesser decreases were reported by Tochigi, Nara, and Ehime. The greatest increase was in Kagoshima 34.5 to 61.6.

VENEREAL DISEASE CLINICS ACTIVITIES

Monthly reports of activities of venereal disease clinics (Ref. Table 54) have been received since November 1947. During that period there has been a considerable increase in the activities of these clinics. The average monthly case load for all venereal diseases during 1947, for which data were available, was 29,537. The monthly average for 1948 (59,510) was approximately twice as high and that for 1949 (79,681), more than two and a half times the 1947 average. This pattern of increasing case load was followed in approximately the same proportions for each of the venereal diseases as a result of improvement in the case finding program.

Cases of primary syphilis accounted for a slightly greater proportion of all syphilis cases treated in 1948 than in 1947. In 1947 the average monthly number of cases of primary syphilis (3,150) amounted to 17% of all syphilis cases (18,107), while in 1948 primary syphilis (7,816) accounted for 21% of all syphilis (37,794). In 1949 the average monthly number of cases of primary syphilis (10,459) had decreased slightly in proportion to total syphilis cases (52,128) and represented 20% of that total. The proportionate decrease was quite regular throughout the year (Ref. Table 99). In January 1949 cases of primary syphilis (9,865) accounted for 21% of the total (46,602). By May primary syphilis (10,580) accounted for 20% of the total (51,981). The greatest number of cases was reached in August, when there were 56,231 cases of all stages of syphilis, of which primary syphilis (11,133) still accounted for 20%. Thereafter the number of cases declined. In December there were 51,606 cases under treatment of which 9,698 (19%) were primary.

The average number of cases of congenital syphilis per month in 1947 was 891, or 5% of all syphilis. In 1948 the average number had risen to 1,959 but was still 5% of the total. In 1949 there was an average of 3,137 cases of congenital syphilis under treatment each month, or 6% of the total. Through 1949 there was an almost continuous but gradual rise both in numbers and percent in each successive month (Ref. Table 99) from 2,596 (6%) in January to 3,462 (7%) in December.

The average monthly case load for gonorrhea in 1947 was 9,923 of which 6,320 (64%) were acute cases. In 1948 the average gonorrhea monthly case load had risen to 18,333, and acute cases (12,207) accounted for 67% of the total. In 1949 the average monthly case load (23,216) and number of acute cases (15,328) had again risen, but the proportion

Public Health and Welfare in Japan - 1949

of acute to total cases was slightly lower (66%). During 1949 the number of cases of gonorrhea rose somewhat irregularly from 21,697 in January to a high point of 24,832 in September and decreased, again somewhat irregularly, to 23,687 in December. The acute cases in January (14,128) represented 65% of the total and rose generally throughout the year, reaching 68% in December (16,136).

The number of new cases of all venereal diseases, not previously treated by any clinic or physician, admitted to venereal disease clinics during the last two months of 1947 was 25,669. Over 50% of all new cases in 1947 were gonorrhea (13,211) and about 40% were syphilis (10,131). In 1948 there were 212,889 new cases, of which slightly less than 50% were gonorrhea (101,563) and about 45% syphilis (96,819). In 1949 the number of new cases had decreased to 176,642, of which about 45% (80,740) were gonorrhea and nearly 50% (85,716) were syphilis. In 1947 new cases of gonorrhea accounted for two-thirds of the total gonorrhea case load, in 1948 slightly less than half, and in 1949 less than a third. New cases of syphilis accounted for over a fourth of the total case load for syphilis in 1947, about a fifth in 1948, and about one-seventh in 1949.

In January 1949 the total number of cases of all venereal diseases, not previously treated by any clinic or physician, was 14,021. (Ref. Table 100). The number increased with some irregularity to a high point of 16,848 in May and decreased thereafter to a low point of 11,595 in December. The number of new cases of syphilis in January was 6,524. The high point for the year was reached in July (8,177) and the low point in December (5,529). New cases of gonorrhea in January amounted to 6,425. The high point for the year was reached in May (7,856) and the low in December (5,420). New cases of syphilis in January accounted for 14% of the total syphilis case load for the month. The percentage increased to 17 in February and decreased thereafter, with new cases accounting for 11% of the December case load. New cases of gonorrhea accounted for 30% of the gonorrhea case load in January, reached a high point (35%) in May and decreased to 23% by December.

During the last two months of 1947 a total of 10,673 venereal disease cases discontinued treatment without permission. The number of syphilis cases (6,317) discontinuing treatment amounted to 17% of the syphilis case load, and the number of gonorrhea cases (3,779) to 19%. In 1948, 65,101 cases of syphilis discontinued treatment, 14% of the syphilis case load and 27,438 gonorrhea cases, or 12% of the gonorrhea case load. In 1949, 69,431 syphilis cases (11%) and 20,195 gonorrhea cases (7%) discontinued treatment. In 1948, 11,633 delinquent cases of all venereal diseases were visited out of a total of 96,544 or approximately 12 out of each 100 cases. Correspondingly, in 1949 22,648 delinquent cases were visited out of a total of 92,923 or about 24 out of each 100 cases. Although this shows improvement in 1949, 3 out of every 4 delinquent cases remained unvisited.

The trend in percent of cases discontinuing treatment each month in 1949 was irregular. (Ref. Table 101) In January 6,801 syphilis cases or 15% of the syphilis case load in 1949 discontinued treatment without permission. The proportion decreased to 11% in February and

Public Health and Welfare in Japan - 1949

remained at about the same level until the low point for the year was reached in October (9% or 5,064 cases). The following month the number discontinuing (6,849) was the highest in numbers of any month of the year, and the second highest proportionally (13%). In December the number decreased to 5,040 or 10% of the syphilis case load. Gonorrhea cases discontinuing treatment in January (1,706) amounted to 8% of the gonorrhea case load. The trend was fairly level (about 7%) through most of the year, but in September rose to 2,209 cases (9%). In October the number decreased to 1,507 (6%), but increased again in November to 2,358 (10%) a high point for the year. In December the number of cases decreased again to 1,283 (5%), the low point for the year.

During the last two months of 1947 there were 13,205 contacts reported, 8,733 for syphilis, 3,885 for gonorrhea, and the remainder for chancroid and the other venereal diseases. For 1948 there were 99,848 total contacts reported, including 62,584 for syphilis and 26,611 for gonorrhea. In 1949 there was a slight decrease in the total contacts reported to 97,815. However both syphilis (67,473) and gonorrhea (28,105) had slight increases. The decrease was accounted for by chancroid and other venereal disease contacts, which decreased from a total of 10,653 in 1948 to 2,237 in 1949.

During 1949 the number of contacts reported ranged from 7,300 to 8,300 each month except in September when it increased to 10,525 and in October was 9,124. (Ref. Table 102) For syphilis the number was between 5,000 and 6,000 each month except in September when it was 7,707. For gonorrhea the number was between 2,000 and 2,500 each month except in September (2,673) and October (2,660).

There were 10,635 visits made for contact tracing by venereal disease clinics in November and December 1947. For the year 1948 there were 75,013 visits, and in 1949 nearly as many (74,289). During 1949 the number of visits per month was lowest in February (4,623) and highest in September (7,999). (Ref. Table 103) Cases found by contact tracing in 1949 totalled 15,378 compared to 13,503 in 1948. The average number of visits per case found was 4.8 in 1949 and 5.6 in 1948.

NUTRITION SURVEYS

For the discussion of results of nutrition surveys refer to Volume 1. These data are shown in Tables 55-58, 104-111.

SCHOOL LUNCH PROGRAM

For the discussion of the school lunch program refer to Volume 1. A tabulation of the data is shown in Table 59.

Public Health and Welfare in Japan - 1949

PUBLIC ASSISTANCE

For the discussion of the public assistance program refer to Volume 1. A tabulation of the data is indicated in Table 6C.

NON-NATIONALS

Vital events of non-nationals (exclusive of those connected with the Occupation of Japan) are shown in Table 112. There were 21,869 births in 1949 compared to 20,549 in 1948 and 13,301 in 1947. Deaths also increased in 1949, there being 5,222 compared to 4,432 in 1948 and 4,184 in 1947. Infant deaths under one year totalled 1,475 compared to 1,196 and 1,142 recorded in the two preceding years. Correspondingly, there were 1,261 stillbirths compared to 1,037 and 789.

Marriages increased to 896 compared to 748 in 1948. The 1947 record (2,838) of marriages is not comparable because it includes many marriages between persons one of whom was not a Japanese national. It is known to contain 700 marriages between members of the Occupation Forces and Japanese Nationals. Divorces decreased to 24 from 34 in 1948. However, the 138 divorces recorded in 1947 is not comparable, for the reason already explained concerning marriages.

BIRTHS, DEATHS AND INFANT DEATHS OF JAPANESE NATIONALS OUTSIDE OF JAPAN

The number of live births of Japanese Nationals which occurred outside Japan Proper in 1949 and which were recorded in Japan totalled 5,958 compared to 8,896 for 1948 and 24,633 for 1947. There were 27,113 deaths compared to 142,062 in 1948 and 470,268 in 1947. Infant deaths totalled 567 compared to 966 in 1948 and 3,435 in 1947. Data for the above by months may be found in Table 113.

TABLE 1. - POPULATION BY PREFECTURE, JAPAN, 1949

Prefecture	1949	Prefecture	1949
All Japan	82,200,000		
Aichi	3,305,900	Nagano	2,131,100
Akita	1,315,400	Nagasaki	1,604,300
Amori	1,248,400	Nara	797,900
Chiba	2,193,400	Niigata	2,495,700
Ehime	1,517,700	Osaka	1,276,500
Fukui	751,500	Okayama	1,691,100
Fukushima	3,394,500	Osaka	3,602,100
Gifu	2,076,600	Saga	954,400
Gunma	1,562,500	Saitama	2,184,900
	1,648,700	Shiga	894,400
Hiroshima	2,096,500	Shimane	925,900
Hokkaido	4,120,500	Shizuoka	2,466,600
Hyogo	3,234,900	Tochigi	1,596,400
Ibaraki	2,095,100	Tokushima	890,800
Ishikawa	965,100	Tokyo	5,551,800
Iwate	1,326,200	Tottori	607,500
Kagawa	957,200	Toyama	1,023,000
Kagoshima	1,810,200	Wakayama	1,004,200
Kanagawa	2,374,800	Yamagata	1,379,800
Kochi	887,800	Yamaguchi	1,542,700
		Yamanashi	835,600
Kumamoto	1,830,200		
Kyoto	1,828,900		
Mie	1,487,000		
Miyagi	1,635,800		
Miyazaki	1,078,500		

Footnotes: Estimate is of 1 July 1949 based on the census of 1 October 1947, net natural increase and migration. Distributions by prefecture are based on the ration count of 1 August 1948.

This table corresponds to Table 1, Statistical Annex, Public Health and Welfare in Japan, 1949.

TABLE 2. - 1/POPULATION BY AGE: JAPAN, 1948-1949

Year	All Ages	0-4	5-9	10-14	15-19	20-24	25-30	30-34
1949	82,603,000	10,577,000	9,842,000	8,785,000	8,604,000	7,723,000	6,037,000	5,115,000
1948	80,697,000	10,154,000	9,443,000	8,884,000	8,516,000	7,447,000	5,740,000	5,191,000

Year	35-39	40-44	45-49	50-54	55-59	60 & over
1949	5,043,000	4,455,000	4,050,000	3,374,000	2,709,000	6,291,000
1948	4,985,000	4,309,000	3,988,000	3,288,000	2,687,000	6,066,000

Footnotes: 1/ All data are as of 1 October each year. Estimates by age are provisional, and were made by Institute of Population Problems, Ministry of Welfare.

This Table corresponds to Table 2, Statistical Annex, Public Health and Welfare in Japan, 1948.

TABLE 3. 1/POPULATION, LIVE BIRTHS, DEATHS, INFANT DEATHS, STILLBIRTHS,
MARRIAGES AND DIVORCES: JAPAN, 1947-1949

Year	2/ Population	LiveBirths	Deaths	3/ Infant Deaths	4/ Stillbirths	Marriages	Divorces
*1949	82,200,000	2,723,873	950,719	168,989	192,970	846,014	82,872
*1948	80,200,000	2,709,871	950,610	165,406	144,095	960,275	79,415
1947	78,000,000	2,678,792	1,138,238	205,360	123,837	934,170	79,551

Footnotes: * All 1949 data are provisional. All 1948 data are provisional except deaths and infant deaths. Death data for 1948 and all data for 1947 are from Final Annual Report, Ministry of Welfare. Other data from Monthly Schedule Report, Ministry of Welfare.

1/ Only events occurring to Japanese nationals in Japan.

2/ Estimates are as of 1 July each year.

3/ Deaths under one year of age.

4/ Stillbirths after third month.

This Table corresponds to Table 3, Statistical Annex, Public Health and Welfare in Japan, 1948.

TABLE 4. - 1/LIVE BIRTH, DEATH, INFANT DEATH, STILLBIRTH, MARRIAGE AND DIVORCE RATES: JAPAN, 1947-1949

Year	Live Births	Deaths	2/ Infant Deaths	Stillbirths	Marriages	Divorces
*1949	33.1	11.6	62.0	70.8	10.3	1.0
*1948	33.8	11.9	61.0	53.2	12.0	1.0
1947	34.3	14.6	76.7	46.2	12.0	1.0

Footnotes: * All 1949 data are provisional. All 1948 data are provisional except deaths. Death data for 1948 and all data for 1947 are from Final Annual Report, Ministry of Welfare, other data from Monthly Schedule Report, Ministry of Welfare.

1/ Birth, death, marriage and divorce rates are the number of events per 1,000 population, infant death and stillbirth rates are the number of events per 1,000 live births in the corresponding period. Tabulations include only events occurring to Japanese nationals in Japan.

2/ Deaths under one year of age.

This Table corresponds to Table 4, Statistical Annex, Public Health and Welfare in Japan, 1948.

TABLE 5. - LIVE BIRTHS BY MONTH: JAPAN, 1947-1949

Year	Total Births	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
*1949	2,723,873	326,166	244,506	249,484	220,191	203,738	189,937	211,921	219,500	221,808	219,877	211,220	204,825
*1948	2,709,871	314,700	260,481	257,373	224,293	201,575	187,732	208,502	215,083	215,088	219,017	219,476	190,451
1947	2,678,792 ¹	295,465	226,018	235,891	209,159	195,574	194,633	226,560	236,331	231,874	229,058	210,764	186,961

Footnotes: * Data for 1948 and 1949 are provisional from Monthly Schedule Report, Ministry of Welfare. Data for 1947 are final from Final Annual Report, Ministry of Welfare. Data refer to births of Japanese nationals in Japan.
 1/ Total includes 4 births of unknown month which are not distributed.

This Table corresponds to Table 5, Statistical Annex, Public Health and Welfare in Japan, 1948.

TABLE 6. - LIVE BIRTH RATES BY MONTH: JAPAN, 1947-1949
(Rates per 1,000 population)

Year	Annual	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
*1949	33.1	46.7	38.8	35.7	32.7	29.2	28.1	30.4	31.4	32.8	31.5	31.3	29.3
*1948	33.8	46.3	41.0	37.0	34.1	29.7	28.6	30.3	31.7	32.7	32.1	33.4	28.0
1947	34.3	44.6	37.8	35.6	32.6	29.5	30.4	34.2	35.8	36.2	34.6	32.9	28.2

Footnotes: * Data for 1948 and 1949 are provisional from Monthly Schedule Report, Ministry of Welfare. Data for 1947 are final from Final Annual Report, Ministry of Welfare. All rates were computed by Public Health and Welfare Section, GHQ, SCAP.

This Table corresponds to Table 6, Statistical Annex, Public Health and Welfare in Japan, 1948.

TABLE 7. - LIVE BIRTHS BY PREFECTURE: JAPAN, 1947-1949

Year	All Japan	Aichi	Akita	Aomori	Chiba	Ehime	Fukui	Fukuoka	Fukushima	Gifu	Gumma	
*1949	2,723,873	107,117	47,736	51,703	68,116	52,797	24,925	125,514	73,867	49,989	52,417	
*1948	2,709,871	110,905	44,005	47,064	67,326	53,501	26,106	119,550	72,032	53,419	52,414	
1947	2,678,792	104,377	47,838	48,997	73,955	52,775	25,445	108,237	71,232	51,432	54,605	
Year	Hiroshima	Hokkaido	Hyogo	Ibaraki	Ishikawa	Iwate	Karawa	Kagoshima	Kanagawa	Kochi	Fumamoto	Kyoto
*1949	63,039	166,215	100,354	67,326	32,250	49,740	31,107	64,875	75,555	26,587	63,551	52,930
*1948	62,385	154,763	103,478	67,624	34,501	47,301	33,968	63,746	73,661	27,460	61,264	54,783
1947	67,757	141,577	97,371	69,164	37,289	46,083	34,284	58,769	74,699	28,656	61,213	53,828
Year	Mie	Miyagi	Miyazaki	Nagano	Nagasaki	Nara	Niigata	Oita	Okayama	Osaka	Saza	Saitama
*1949	43,610	57,430	40,508	59,316	61,711	21,946	84,471	42,511	50,235	113,241	34,391	71,855
*1948	47,459	56,699	41,261	60,594	57,903	23,534	82,482	43,923	52,899	113,163	33,027	70,838
1947	46,234	55,512	38,641	61,920	53,021	23,989	86,204	42,330	53,528	103,782	31,958	77,234
Year	Shiga	Shimane	Shizuoka	Tochigi	Tokushima	Tokyo	Tottori	Toyama	Wakayama	Yamagata	Yamaguchi	Yamanashi
*1949	25,794	29,678	81,390	54,054	29,838	171,309	18,801	34,392	29,139	44,340	50,760	25,443
*1948	27,474	30,638	83,600	54,397	32,233	164,867	20,011	35,962	30,101	42,286	48,224	25,140
1947	25,869	32,534	81,560	54,870	31,646	157,306	19,348	41,428	30,803	43,941	49,246	26,305

Footnotes: * Data for 1948 and 1949 are provisional. Data for 1948 and 1949 are from Monthly Schedule Reports, Ministry of Welfare. Data for 1947 are from Final Annual Report, Ministry of Welfare. All data refer only to births of Japanese nationals in Japan.

This Table corresponds to Table 7, Statistical Annex, Public Health and Welfare in Japan, 1949.

TABLE 8.- LIVE BIRTH RATES BY PREFECTURE: JAPAN, 1947-1949
(Rates per 1,000 population)

Year	All Japan	Aichi	Akita	Aomori	Chiba	Ehime	Fukui	Fukuoka	Fukushima	Gifu	Gumma
1949	33.1	32.4	36.3	41.4	31.1	34.8	33.2	37.0	35.6	32.0	31.8
1948	33.8	34.4	34.3	38.6	31.5	36.1	35.6	36.1	36.6	35.0	32.6
1947	34.3	33.5	38.1	41.6	35.0	36.3	35.1	34.1	36.8	34.5	34.8

Year	Hiroshima	Hokkaido	Hyogo	Ibaraki	Ishikawa	Iwate	Kagawa	Kagoshima	Kanazawa	Kochi	Kumamoto	Kyoto
1949	30.1	40.3	31.0	32.1	33.4	37.5	32.5	35.8	31.8	29.9	34.7	28.9
1948	30.5	38.5	32.8	33.1	36.6	36.6	36.4	36.1	31.8	31.7	34.3	30.7
1947	33.7	36.8	31.9	34.4	40.2	36.5	37.4	33.7	33.7	33.8	34.7	31.0

Year	Mie	Miyagi	Miyazaki	Nagano	Nagasaki	Nara	Niigata	Oita	Okayama	Osaka	Saga	Saitama
1949	29.3	35.1	37.6	27.8	38.5	27.5	33.8	33.3	29.7	31.4	36.0	32.9
1948	32.7	35.5	39.2	29.1	37.0	30.2	33.9	35.3	32.1	32.2	35.5	33.2
1947	32.7	35.5	37.7	30.1	34.7	30.8	35.7	34.4	33.1	31.2	34.9	36.8

Year	Shiga	Shimane	Shizuoka	Tochigi	Tokyo	Tottori	Tokushima	Toyama	Wakayama	Yamagata	Yamaguchi	Yamanashi
1949	28.8	32.1	33.0	33.9	30.9	30.9	33.5	33.6	29.0	32.1	32.9	30.4
1948	31.5	33.9	34.7	34.9	30.4	33.7	37.1	36.0	30.7	31.4	32.0	30.8
1947	30.2	36.4	34.7	35.8	31.5	33.0	37.1	42.4	32.1	32.9	33.3	32.6

Footnotes: * Data for 1948 and 1949 are provisional. Rates are the number of births per 1,000 population. Data for 1948 and 1949 are from Monthly Schedule Report, Ministry of Welfare. Data for 1947 are from Final Annual Report, Ministry of Welfare. All data refer only to births of Japanese nationals in Japan. Rates were computed by Public Health and Welfare Section, GHQ, SCAP.

This Table corresponds to Table 8, Statistical Annex, Public Health and Welfare in Japan, 1948.

TABLE 9. - DEATHS BY MONTH: JAPAN, 1948-1949

Year	Total Deaths	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Unknown
*1949	950,719	86,814	77,763	90,267	84,155	78,768	74,297	79,443	77,981	73,630	72,120	73,347	82,134	
1948	950,610	93,717	88,435	90,876	78,832	74,316	75,385	78,640	79,279	72,523	69,977	70,948	77,605	77

Footnotes: *Data for 1949 for provisional from Monthly Schedule Reports, Ministry of Welfare. Data for 1948 are final from Annual Report, Ministry of Welfare. All data refer to deaths of Japanese nationals in Japan.

This Table corresponds to Table 9, Statistical Annex, Public Health and Welfare in Japan, 1948.

TABLE 10. - DEATH RATES BY MONTH: JAPAN, 1948-1949
(Rates per 1,000 population)

Year	Annual	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
*1949	11.6	12.4	12.3	12.9	12.5	11.3	11.0	11.4	11.2	10.9	10.3	10.9	11.8
*1948	11.9	13.8	13.9	13.4	12.0	10.9	11.5	11.6	11.7	11.0	10.3	10.8	11.4

Footnotes: *Data for 1949 are provisional, for 1948 are final. Rates are per 1,000 population and are computed on an annual basis. Deaths refer to deaths of Japanese nationals in Japan. Data for 1949 are from Monthly Schedule Reports, Ministry of Welfare, and for 1948 are from Final Annual Report, Ministry of Welfare. Rates were computed by Public Health and Welfare Section, GHQ, SCAP.

This Table corresponds to Table 10, Statistical Annex, Public Health and Welfare in Japan, 1948.

TABLE 11. - DEATHS BY PREFECTURE: JAPAN, 1948-1949

Year	All Japan	Aichi	Akita	Aomori	Chiba	Ehime	Fukui	Fukuoka	Fukushima	Gifu	Gunma	
* 1949	950,719	36,463	17,233	16,842	26,037	17,645	9,954	40,041	23,841	18,536	18,452	
1948	950,610	35,370	18,501	18,024	26,211	17,325	13,662	37,107	23,823	18,545	18,388	
Year	Hiroshima	Hokkaido	Hyogo	Ibaraki	Ishikawa	Iwate	Kagawa	Kagoshima	Kanagawa	Kochi	Kumamoto	Kyoto
* 1949	23,512	48,361	74,695	24,854	13,015	18,427	11,247	21,455	22,985	10,577	21,401	19,707
1948	23,909	47,227	75,264	24,274	13,475	19,194	11,497	21,622	22,683	10,672	21,165	19,892
Year	Mie	Miyagi	Miyazaki	Nagano	Nagasaki	Nara	Niigata	Oita	Okayama	Osaka	Saga	Saitama
* 1949	17,526	18,184	12,913	22,818	19,709	9,584	31,550	17,001	19,557	38,883	12,306	25,698
1948	17,588	18,575	12,180	23,692	18,899	9,700	32,737	16,697	21,165	37,142	12,630	24,789
Year	Shiga	Shimane	Shizuoka	Tochigi	Tokushima	Tokyo	Tottori	Toyama	Wakayama	Yamagata	Yamaguchi	Yamanashi
* 1949	11,217	12,033	25,380	17,855	11,081	54,240	7,356	14,287	11,183	17,656	18,508	8,914
1948	11,294	12,119	24,762	17,448	11,333	51,624	7,673	14,032	11,037	18,232	18,433	8,995

FOOTNOTES: * Data for 1949 are provisional.

Data refer only to deaths of Japanese nationals in Japan.

Data for 1948 are from Final Annual Report, Ministry of Welfare, and for 1949 from Monthly Schedule Reports, Ministry of Welfare.

This table corresponds to table 11, Statistical Annex, Public Health and Welfare in Japan, 1948.

TABLE 12. DEATH RATES BY PREFECTURE: JAPAN, 1948-1949
(Rates per 1,000 population)

Year	All Japan	Aichi	Akita	Aomori	Chiba	Ehime	Fukui	Fukuoka	Fukushima	Gifu	Gumma	
*1949	11.6	11.0	13.1	13.5	11.9	11.6	13.2	11.8	11.5	11.9	11.2	
1948	11.9	11.0	14.4	14.8	12.2	11.7	18.6	11.2	11.8	12.2	11.4	
Year	Hiroshima	Hokkaido	Hyogo	Ibaraki	Ishikawa	Iwate	Karawa	Karoshima	Kanagawa	Kochi	Kumamoto	Kyoto
*1949	11.2	11.7	10.7	11.9	13.5	13.9	11.7	11.9	9.7	11.9	11.7	10.8
1948	11.7	11.7	11.2	11.9	14.3	14.8	12.3	12.2	9.8	12.3	11.9	11.2
Year	Mie	Miyagi	Miyazaki	Nagano	Nagasaki	Nara	Niigata	Oita	Okayama	Osaka	Saga	Saitama
*1949	11.8	11.1	12.0	10.7	12.3	12.0	12.6	13.3	11.6	10.8	12.9	11.8
1948	12.1	11.6	11.6	11.4	12.1	12.5	13.4	13.4	12.8	10.6	13.6	11.6
Year	Shiga	Shimane	Shizuoka	Tochigi	Tokushima	Tokyo	Tottori	Toyama	Wakayama	Yamagata	Yamaguchi	Yamanashi
*1949	12.5	13.0	10.3	11.2	12.4	9.8	12.1	14.0	11.1	12.8	12.0	10.7
1948	12.9	13.4	10.3	11.2	13.0	9.5	12.9	14.1	11.3	13.5	12.2	11.0

FOOTNOTES: *Data for 1949 are provisional.

Data refer only to deaths of Japanese nationals in Japan.

Data for 1948 are from Final Annual Report, Ministry of Welfare, and for 1949 from Monthly Schedule

Reports, Ministry of Welfare.

Rates were computed by Public Health and Welfare Section, GHQ, SCAP.

This Table corresponds to Table 12, Statistical Annex, Public Health and Welfare in Japan, 1948.

TABLE 13. - 1/DEATHS FROM SELECTED CAUSES: JAPAN, 1948-1949

Year	All Causes	2/Typhoid fever (1)	Paratyphoid fever (2)	Plague (3)	Cholera (4)	Cerebrospinal meningitis (6)	Anthrax (7)	3/Scarlet fever (8)	Whooping cough (9)	4/Diphtheria (10)
*1949	950,719	970	116	-	-	497	-	51	9,147	1,659
1948	950,610	1,433	170	-	-	650	-	42	4,746	1,903

Year	Tetanus (12)	5/Tuberculosis (all forms) (13-22)	Leprosy (23)	Glanders (26a)	Dysentery (27)	Malaria (28)	6/Syphilis (30)	Influenza (33)	Smallpox (34)	Measles (35)
*1949	1,964	138,764	162	1	7,824	122	5,492	502	21	12,440
1948	2,138	143,909	150	-	5,157	224	4,356	515	3	5,598

Year	Japanese "B" encephalitis (37b)	Yellow fever (38a)	Rabies (38b)	Typhus fever and typhus-like diseases (39)	Cancer and other malignant tumors (45-55)	Diabetes mellitus (61)	Beriberi (68)	8/Simple meningitis (81a)
*1949	1,182	-	76	67	58,890	1,885	5,591	9,108
1948	2,620	-	40	93	55,677	1,789	6,281	10,414

Year	9/Intracranial lesions of vascular origin (83)	10/Diseases of the heart (90-95)	Bronchitis (106)	12/Pneumonia (all forms) (107-109)	Pleurisy not specified as tuberculous (110)	13/Ulcer of stomach or duodenum (117)
*1949	100,650	52,625	25,695	56,636	6,030	19,521
1948	94,329	49,046	25,932	52,979	7,700	20,705

TABLE 13. - 1/DEATHS FROM SELECTED CAUSES: JAPAN, 1948-1949, Cont'd

Year	10/Diarrhea, enteritis and ulceration of intestines (all ages) (119-120)	15/Diarrhea, enteritis and ulceration of intestines (under 2 years) (119)	16/Diarrhea, enteritis and ulceration of intestines (2 years and over) (120)	17/ Appendicitis (130-132)	Nephritis (130-132)
• 1949	72,186	41,822	30,364	3,267	33,859
1948	83,264	48,444	34,820	3,808	36,561

Year	18/Diseases of pregnancy, childbirth, puerperium (140-150)	Conenital malformations (157)	Conenital debility (158)	Premature birth (159)	Injury at birth (160)	19/ Senility (162)	Suicide (163-164)
• 1949	4,495	6,556	37,285	13,667	1,082	66,699	14,211
1948	4,437	5,751	38,204	13,720	1,030	64,074	12,753

Year	Homicide (165-168)	Accidental deaths (169-195)	20/All-defined and unknown (200 a & c)	All other causes
• 1949	1,669	34,498	16,900	126,457
1948	1,641	38,960	17,188	130,620

Footnotes:

• Data for 1949 are provisional.

Data for 1948-1949 refer to deaths of Japanese nationals only in Japan.

1/All data for 1948-1949 are based on death certificates. Data for 1948 are from Final Annual Report, Ministry of Welfare, and for 1949 from Monthly Schedule Report, Ministry of Welfare.

2/Typhoid fever excludes typhoid pneumonia.

3/Scarlet fever excludes puerperal scarlet fever.

4/Diphtheria includes diphtheric endocarditis and excludes croupous bronchitis and laryngeal croup.

5/Tuberculosis includes spondylitis and excludes chronic pleuropneumonia and nontubercular adenopathy.

6/Includes general paralysis of insane and aneurysm of the aorta.

TABLE 13. - 1/DEATHS FROM SELECTED CAUSES: JAPAN, 1945-1949, Cont'd

Footnotes: Cont'd

- 7/ includes tubercle-like diseases.
- 8/ excludes acute cerebrospinal meningitis not due to meningococcus.
- 9/ excludes alcoholic erythema congestio.
- 10/ excludes hypertrophic endocarditis
- 11/ includes croupous bronchitis.
- 12/ includes typhoid pneumonia.
- 13/ excludes ulcer of duodenum.
- 14/ excludes malnutrition, 1 year of age and over.
- 15/ excludes malnutrition, 1 to 2 years of age.
- 16/ excludes malnutrition, 2 years of age and over.
- 17/ excludes alcoholic nephritis.
- 18/ includes puerperal scarlet fever.
- 19/ includes senility with mention of senile dementia.
- 20/ includes malnutrition 1 year of age and over, excludes sudden death and found dead.

"NA" indicates data not available.
A dash (-) indicate no deaths were reported.

This Table corresponds to Table 13, Statistical Annex, Public Health and Welfare in Japan, 1949.

TABLE 14. - 1/DEATH RATES FOR SELECTED CAUSES: JAPAN, 1948-1949
(per 100,000 population)

Year	All Causes	2/Typhoid fever (1)	Paratyphoid fever (2)	Plague (3)	Cholera (4)	Cerebrospinal (meningococcus) meningitis (6)	Anthrax (7)	3/Scarlet fever (8)	Whooping cough (9)	4/Diphtheria (10)
1949	1156.6	1.2	0.1	-	-	0.6	-	0.1	11.1	2.0
1948	1185.3	1.8	0.2	-	-	0.8	-	0.1	5.9	2.4

Year	Tetanus (12)	5/Tuberculosis (all forms) (13-22)	Leprosy (23)	Glanders (26a)	Dysentery (27)	Malaria (28)	Syphilis (30)	Influenza (33)	Smallpox (34)	Measles (35)
1949	2.4	168.8	0.2	0.0	9.5	0.1	8.7	0.6	0.0	15.1
1948	2.7	179.4	0.2	-	6.4	0.3	5.4	0.6	0.0	7.0

Year	Japanese "B" encephalitis (37b)	Yellow fever (38a)	7/Typhus fever and typhus-like diseases (39)	Cancer and other malignant tumors (45-55)	Diabetes mellitus (61)	Beriberi (66)	8/Simple meningitis (81a)
1949	1.4	-	0.1	71.6	2.3	6.8	11.1
1948	3.3	-	0.0	69.4	2.2	7.8	13.0

Year	9/Intracranial lesions of vascular origin (83)	10/Diseases of the heart (90-95)	11/E Bronchitis (106)	12/Pneumonia (all forms) (107-109)	Pleurisy not specified as tuberculous (110)	13/Ulcer of the stomach or duodenum (117)
1949	122.4	64.3	31.3	68.9	7.3	23.7
1948	117.6	61.2	32.3	66.1	9.6	25.8

TABLE 14. - 1/DEATH RATES FOR SELECTED CAUSES: JAPAN, 1948-1949, Cont'd
(per 100,000 population)

Year	14/Diarrhea, enteritis and ulceration of intestines (all ages) (119-120)	15/Diarrhea, enteritis and ulceration of intestines (under 2 years) (119)	16/Diarrhea, enteritis and ulceration of intestines (2 years and over)(120)	17/ Nephritis (130-132)			
1949	87.8	50.9	36.9	4.0			
1948	103.8	60.4	43.4	4.7			
				45.6			
Year	18/Diseases of pregnancy, childbirth, puerperium (140-150)	Conenital malformations (157)	Conenital debility (158)	Premature birth (159)	Injury at birth (160)	19/Senility (162)	Suicide (163-164)
1949	5.5	8.0	45.4	16.6	1.3	81.1	17.3
1948	5.5	7.2	47.6	17.1	1.3	79.9	15.9
Year	Homicide (165-168)	Accidental deaths (169-195)	20/ill-defined and unknown (200 a & c)	All other causes			
1949	2.0	42.0	20.6	153.8			
1948	2.0	48.6	21.4	162.9			

Footnotes: • Data for 1949 are provisional.

1/Rates are the number of deaths per 100,000 population estimated as of 1 July of the year concerned, and were computed by Public Health and Welfare Section, GHQ, SCAP. 1948 data are from Final Annual Report, Ministry of Welfare, and for 1949 from Monthly Schedule Reports, Ministry of Welfare.

2/Excludes typhoid pneumonia.

3/Excludes puerperal scarlet fever.

4/Includes diphtheric endocarditis and excludes chronic pleuro pneumonia and non-tubercular adenopathy.

5/Includes spondylitis and excludes chronic pleuro pneumonia and non-tubercular adenopathy.

6/Includes general paralysis of insane and aneurysm of aorta.

TABLE 14. - 1/DEATH RATES FOR SELECTED CAUSES: JAPAN, 1948-1949, Cont'd

Footnotes: Cont'd

- 7/Includes typhus-like diseases.
- 8/Excludes acute cerebrospinal meningitis not due to meningococcus.
- 9/Excludes alcoholic epilepsy and congestion.
- 10/Excludes diphtheric endocarditis.
- 11/Includes croupous bronchitis.
- 12/Includes typhoid pneumonia.
- 13/Includes ulcer of duodenum.
- 14/Excludes malnutrition, 1 year of age and over.
- 15/Excludes malnutrition, 1 to 2 years of age.
- 16/Excludes malnutrition, 2 years of age and over.
- 17/Excludes alcoholic nephritis.
- 18/Includes puerperal scarlet fever.
- 19/Includes senility with mention of senile dementia.
- 20/Includes malnutrition 1 year of age and over, excludes sudden death and found dead.

A dash (-) indicates that no deaths were reported.

A rate of 0.0 indicates that there were some deaths but that the rate was less than 0.1.

This Table corresponds to Table 14, Statistical Annex, Public Health and Welfare in Japan, 1948.

TABLE 15. - 1/ DEATHS FROM SELECTED CAUSES BY MONTH: JAPAN, 1920 - 1949
2/ TYPHOID FEVER (1)

Year	Total Deaths	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Un- known
*1949	970	57	57	72	54	68	82	96	147	123	90	59	65	-
1948	1,433	84	86	65	72	127	123	123	223	163	114	96	83	-
1947	2,926	238	153	144	164	180	218	297	543	415	289	181	104	-
1946	5,388	454	430	573	384	403	445	411	624	523	397	452	292	-
1945	7,999	223	224	266	279	313	402	526	921	1,319	1,451	1,107	968	-
1944	7,844	299	357	384	375	524	621	844	1,140	1,070	837	715	678	-
1943	8,858	221	260	299	360	487	484	722	926	1,081	866	596	623	-
1942	6,432	218	303	300	312	347	377	576	715	681	625	457	600	-
1941	6,913	516	408	430	418	513	614	756	854	772	656	485	491	-
1940	7,117	450	371	348	307	423	498	693	891	954	865	718	599	-
1939	6,957	435	388	324	356	471	547	649	806	943	803	695	540	-
1938	7,819	439	359	350	371	495	549	765	1,038	1,164	942	767	580	-
1937	7,124	431	433	431	407	494	559	641	867	869	796	635	561	-
1936	6,871	355	301	289	305	382	510	666	895	1,015	928	666	559	-
1935	7,240	559	369	388	300	397	514	688	883	1,020	914	701	507	-
1934	8,149	540	413	381	360	408	592	785	1,083	1,052	1,097	780	658	-
1933	7,652	506	396	320	286	378	580	694	970	1,083	959	753	727	-
1932	6,954	524	398	362	291	390	484	606	796	909	928	684	602	-
1931	8,176	624	414	421	377	449	680	799	953	1,058	1,012	785	604	-
1930	8,350	536	450	428	384	462	588	850	1,138	1,223	1,002	663	626	-
1929	8,032	526	431	426	380	507	626	735	1,040	1,129	912	730	590	-
1928	8,780	513	499	446	381	502	615	808	1,148	1,210	1,130	865	663	-
1927	7,843	510	372	336	335	393	562	753	956	1,124	1,068	825	609	-
1926	9,783	1,083	966	605	452	497	571	741	1,049	1,259	1,095	841	624	-
1925	10,033	926	641	554	437	457	620	842	1,184	1,298	1,359	986	729	-
1924	14,063	1,231	1,092	883	677	718	809	1,207	1,916	1,779	1,534	1,139	1,078	-
1923	12,939	682	515	434	394	496	687	1,015	1,569	1,798	2,142	1,759	1,443	5
1922	12,439	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1921	11,953	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1920	12,106	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA

See footnotes at end of table.

TABLE 15. - 1/ DEATHS FROM SELECTED CAUSES BY MONTH: JAPAN, 1920 - 1949 - Cont'd
PARATYPHOID FEVER (2)

Year	Total Deaths	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Un-known
* 1949	116	7	3	13	4	4	11	16	15	20	10	9	4	-
1948	170	7	11	10	15	14	20	13	28	14	19	10	9	-
1947	316	18	18	15	21	22	22	35	49	53	33	17	13	-
1946	474	15	17	36	26	42	49	46	60	68	38	46	31	-
1945	526	9	20	15	21	23	32	44	51	80	77	86	68	-
1944	564	11	26	25	24	44	39	64	93	71	65	56	46	-
1943	535	14	13	26	16	22	38	51	69	85	75	51	48	-
1942	282	5	6	12	14	17	24	33	42	33	28	28	31	-
1941	309	12	15	14	33	21	31	24	42	38	34	23	22	-
1940	315	7	13	19	14	14	33	31	53	49	35	22	25	-
1939	295	14	13	16	11	17	19	36	50	35	51	17	16	-
1938	297	13	10	12	9	17	18	40	40	44	37	34	23	-
1937	264	7	10	9	10	23	22	37	46	34	38	15	13	-
1936	277	12	10	3	10	13	26	26	36	52	42	26	21	-
1935	238	11	12	10	14	16	16	32	33	31	25	20	18	-
1934	322	11	10	13	16	25	39	37	61	28	29	37	16	-
1933	334	18	11	8	11	6	28	44	40	75	42	33	18	-
1932	372	26	12	17	16	13	27	40	46	68	52	31	24	-
1931	312	19	23	9	9	10	26	26	41	39	50	34	26	-
1930	373	20	19	14	12	18	35	51	52	65	42	26	19	-
1929	353	17	10	12	14	27	28	38	36	62	48	36	25	-
1928	453	21	21	10	16	20	32	38	74	79	72	41	29	-
1927	490	23	17	15	14	19	31	60	73	86	89	39	24	-
1926	433	30	21	29	16	17	27	41	48	72	66	36	30	-
1925	443	29	16	12	18	20	27	39	48	87	72	37	38	-
1924	597	34	29	29	21	33	51	75	85	85	68	52	35	-
1923	677	35	25	17	15	26	34	75	109	115	99	74	53	-
1922	767	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1921	703	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1920	784	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-

See footnotes at end of table.

TABLE 15. - 1/ DEATHS FROM SELECTED CAUSES BY MONTH: JAPAN, 1920 - 1945 - Cont'd
PLAGUE (3)

Year	Total Deaths	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Un- known
* 1949	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1948	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1947	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1946	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1945	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1944	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1943	1	-	-	-	-	-	-	-	-	-	-	-	-	-
1942	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1941	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1940	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1939	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1938	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1937	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1936	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1935	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1934	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1933	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1932	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1931	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1930	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1929	2	-	-	-	-	1	1	-	-	-	-	-	-	-
1928	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1927	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1926	6	-	-	-	-	-	-	3	1	-	-	-	-	2
1925	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1924	3	1	-	-	-	-	1	1	-	-	-	-	-	-
1923	1	-	-	-	-	-	-	-	-	1	-	-	-	-
1922	67	-	-	-	-	-	-	10	1	1	27	22	6	-
1921	2	-	-	1	-	-	-	-	1	-	-	-	-	-
1920	14	-	-	-	-	-	-	-	-	5	4	4	1	-

See footnotes at end of table.

TABLE 15. - 1/ DEATHS FROM SELECTED CAUSES BY MONTH: JAPAN, 1920 - 1949 - Cont'd
CHOLERA (4)

Year	Total Deaths	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Un-known
(*) 1949	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1948	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1947	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1946	528	-	-	-	-	-	55	90	265	99	2	3	14	-
1945	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1944	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1943	1	-	-	-	-	-	-	-	-	-	-	-	-	-
1942	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1941	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1940	5	-	-	-	-	-	-	3	-	1	-	1	-	-
1939	1	-	-	-	-	-	1	-	-	-	-	-	-	-
1938	10	-	-	-	-	-	-	5	2	2	1	-	-	-
1937	11	-	-	-	-	-	-	-	-	6	3	3	1	-
1936	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1935	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1934	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1933	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1932	1	-	-	-	-	-	1	-	-	-	-	-	-	-
1931	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1930	2	-	-	-	-	1	-	-	1	-	-	-	-	-
1929	114	-	-	-	-	-	-	-	19	81	8	5	1	-
1928	1	-	-	-	-	-	-	-	-	-	1	-	-	-
1927	3	-	-	-	-	-	-	-	1	2	-	-	-	-
1926	13	10	-	-	-	-	-	-	-	-	1	2	-	-
1925	363	-	-	-	-	-	1	3	4	160	134	47	14	-
1924	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1923	31	-	-	3	2	3	2	5	6	5	1	4	-	-
1922	542	-	-	-	-	-	-	5	3	149	378	7	-	-
1921	-	-	1	2	2	1	6	6	7	5	8	1	-	-
1920	3,426	3	3	4	-	1	317	1,284	1,016	621	151	23	3	-

See footnotes at end of table.

TABLE 15. - 1/ DEATHS FROM SELECTED CAUSES BY MONTH: JAPAN, 1920 - 1949 - Cont'd
CEREBROSPINAL (MENINGOCOCCUS) MENINGITIS (6)

Year	Total Deaths	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Un- known
*1949	497	37	35	59	44	33	48	30	46	80	38	23	24	-
1948	660	45	57	85	71	52	48	48	91	53	33	28	39	-
1947	1,187	53	102	203	187	150	94	74	96	87	63	37	41	-
1946	424	12	32	45	52	47	52	27	35	28	31	30	33	-
1945	1,072	19	110	200	233	194	108	74	22	42	28	25	17	-
1944	396	20	40	44	58	47	36	31	27	29	16	23	25	-
1943	572	19	28	55	40	70	39	18	37	25	25	19	23	-
1942	443	21	30	38	32	42	30	25	13	23	28	25	25	-
1941	534	35	49	72	67	62	40	40	43	35	34	34	23	-
1940	725	51	59	98	113	91	70	45	45	36	35	47	35	-
1939	854	34	65	111	136	127	83	75	90	78	34	24	37	-
1938	580	31	53	60	75	82	53	42	43	61	36	23	21	-
1937	479	29	53	43	51	32	49	34	71	39	32	16	30	-
1936	604	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1935	701	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1934	603	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1933	244	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1932	165	8	8	11	15	16	10	11	24	18	21	10	13	-
1931	165	8	11	12	26	12	10	11	9	9	19	17	13	-
1930	169	8	12	12	8	19	11	11	36	14	15	10	13	-
1929	209	7	12	25	23	25	16	16	38	23	3	9	12	-
1928	171	10	10	12	16	20	17	14	15	23	13	7	14	-
1927	214	18	10	21	21	27	20	15	25	25	9	14	9	-
1926	236	12	16	17	17	24	19	20	24	42	22	8	15	-
1925	275	16	23	30	37	41	27	19	19	25	15	12	11	-
1924	1,334	23	28	35	28	21	21	24	672	384	54	21	23	-
1923	396	20	29	53	44	39	42	35	22	40	41	14	17	-
1922	592	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1921	466	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1920	533	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-

See footnotes at end of table.

TABLE 15. - 1/ DEATHS FROM SELECTED CAUSES BY MONTH: JAPAN, 1920 - 1949 - Cont'd
ANTHRAX (7)

Year	Total Deaths	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Un- known
*1949	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1948	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1947	2	-	-	-	1	1	-	-	-	-	-	-	-	-
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1943	9	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1942	15	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1941	9	-	-	-	1	-	-	4	2	1	-	-	1	-
1940	15	1	2	4	2	2	-	2	1	-	-	1	-	-
1939	19	2	1	1	1	6	1	4	-	1	1	1	-	-
1938	10	1	1	1	-	1	2	2	1	1	-	-	-	-
1937	17	3	4	1	3	-	-	-	3	-	2	-	1	-
1936	18	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1935	14	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1934	16	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1933	18	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1932	9	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1931	8	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1930	12	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1929	12	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1928	17	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1927	18	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1926	11	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1925	8	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1924	15	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1923	30	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1922	13	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1921	1	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1920	6	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-

See footnotes at end of table.

TABLE 15. - 1/ DEATHS FROM
3/ SCARLET FEVER (8)
SELECTED CAUSES BY MONTH: JAPAN, 1920 - 1949 - Cont'd

Year	Total Deaths	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Un- known
*1949	51	6	4	1	7	5	5	3	5	2	3	7	3	-
1948	42	1	2	4	6	4	3	3	4	1	2	5	7	-
1947	71	9	9	7	5	10	4	5	7	2	4	4	5	-
1946	100	3	14	23	19	9	5	3	3	6	2	6	7	-
1945	82	8	12	4	3	8	7	7	3	7	4	11	8	-
1944	114	10	10	11	22	9	7	8	9	9	8	5	6	-
1943	197	17	19	18	12	14	15	13	12	10	13	7	15	-
1942	217	18	20	14	19	17	18	21	25	9	14	18	32	-
1941	268	28	35	22	28	20	19	15	17	13	17	26	28	-
1940	389	37	42	47	42	45	31	21	14	18	21	26	45	-
1939	476	44	52	47	44	36	34	34	26	31	19	50	59	-
1938	398	47	42	39	24	31	33	15	24	16	37	48	42	-
1937	454	32	46	44	51	49	38	35	19	22	25	43	50	-
1936	467	42	56	50	39	55	30	14	19	25	40	52	45	-
1935	489	56	55	50	40	33	39	28	29	32	38	45	44	-
1934	505	55	49	61	44	44	41	31	24	24	34	44	54	-
1933	401	35	31	47	34	35	22	29	26	24	30	49	39	-
1932	336	37	33	33	36	27	25	17	15	17	16	33	47	-
1931	327	38	30	37	33	23	25	29	17	27	23	27	28	-
1930	303	24	23	20	25	27	28	12	13	12	22	54	43	-
1929	248	21	27	28	26	27	18	20	8	12	15	25	21	-
1928	220	20	30	20	23	17	14	17	15	14	20	9	21	-
1927	219	22	20	18	20	20	19	16	4	13	19	19	25	-
1926	300	34	30	26	25	27	38	40	19	8	17	15	21	-
1925	284	18	15	23	42	31	29	23	17	8	12	36	30	-
1924	130	14	5	15	14	15	12	6	4	3	10	14	18	-
1923	104	13	11	12	10	11	9	5	4	3	6	8	12	-
1922	107	10	11	12	12	13	11	7	5	3	14	1	8	-
1921	93	13	10	10	6	8	8	6	-	5	11	9	7	-
1920	102	9	13	4	10	9	8	4	5	4	13	13	10	-

See footnotes at end of table.

TABLE 15. - 1/ DEATHS FROM SELECTED CAUSES BY MONTH: JAPAN, 1920 - 1949 - Cont'd
WHOOPING COUGH (9)

Year	Total Deaths	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Un- known
*1949	9,147	566	556	530	580	728	928	1,070	1,186	922	591	588	902	-
1948	4,746	468	375	283	308	355	417	561	552	423	256	296	450	2
1947	17,001	2,123	2,763	2,187	1,621	1,614	1,607	1,594	1,351	852	436	355	457	1
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1943	9,407	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1942	9,510	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1941	10,740	816	826	777	711	809	965	1,345	1,238	1,018	691	659	885	-
1940	8,792	894	778	631	614	689	803	1,014	1,073	746	458	450	642	-
1939	10,103	771	748	861	821	842	949	1,174	1,323	852	577	484	701	-
1938	8,871	887	810	678	563	587	757	968	1,086	851	575	460	649	-
1937	10,985	890	792	875	836	993	1,160	1,337	1,380	968	588	439	727	-
1936	9,751	867	870	719	574	660	869	1,237	1,306	978	590	465	616	-
1935	12,216	957	965	958	988	1,097	1,329	1,648	1,448	986	632	495	709	-
1934	8,520	483	478	481	411	508	772	1,136	1,315	948	627	558	803	-
1933	6,157	725	625	560	468	525	590	642	680	503	302	219	318	-
1932	14,657	1,242	1,280	1,210	1,198	1,361	1,781	1,865	1,782	1,153	627	497	661	-
1931	8,260	467	454	428	449	504	695	946	1,059	1,047	709	578	924	-
1930	7,437	711	583	622	575	607	768	942	955	669	339	287	379	-
1929	9,822	744	689	758	789	816	973	1,277	1,452	930	481	419	494	-
1928	11,440	949	1,015	957	868	903	1,133	1,335	1,441	1,022	638	495	684	-
1927	9,147	627	528	531	556	698	871	1,182	1,322	1,007	558	516	751	-
1926	7,312	488	495	608	568	637	673	893	922	752	480	327	469	-
1925	8,456	889	848	720	562	635	841	1,046	994	719	445	336	421	-
1924	10,256	852	712	783	757	933	1,070	1,250	1,410	906	521	408	653	1
1923	8,484	350	379	430	478	658	932	1,222	1,389	1,010	589	457	590	-
1922	6,273	528	567	458	362	407	585	838	923	618	358	274	325	-
1921	6,952	493	502	467	464	524	659	783	1,041	783	460	335	441	-
1920	7,922	655	663	638	551	630	804	1,094	1,076	679	357	327	448	-

See footnotes at end of table.

TABLE 15. - 1/ DEATHS FROM SELECTED CAUSES BY MONTH: JAPAN, 1920 - 1949 - Cont'd
4/ DIPHTHERIA (10)

Year	Total Deaths	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Un-known
* 1949	1,699	245	196	184	165	115	75	82	49	60	100	178	210	-
1948	1,903	278	233	202	132	93	79	50	59	93	146	245	293	-
1947	3,900	541	428	432	323	269	182	126	94	102	218	292	383	-
1946	3,891	684	550	492	293	207	231	138	137	196	266	376	361	-
1945	7,840	845	923	884	730	653	492	352	261	338	565	873	924	-
1944	6,192	704	648	644	473	334	291	242	276	275	510	700	1,095	-
1943	7,122	544	593	650	557	477	366	253	143	176	364	520	776	-
1942	5,153	461	503	481	370	232	184	161	126	198	344	494	878	-
1941	5,004	679	621	483	414	304	243	168	195	252	419	547	679	-
1940	4,744	626	601	536	386	293	210	164	137	230	377	506	678	-
1939	5,272	648	647	711	552	388	330	227	157	178	353	468	613	-
1938	4,135	572	511	478	378	269	206	162	125	166	254	406	608	-
1937	4,253	537	498	507	390	309	274	199	126	171	298	434	590	-
1936	4,332	612	579	552	391	294	176	142	112	159	290	486	539	-
1935	4,462	593	561	512	433	344	244	154	177	170	301	422	551	-
1934	5,235	805	657	694	500	366	299	199	151	194	369	482	599	-
1933	5,437	615	658	704	546	422	296	189	151	216	398	537	705	-
1932	4,822	542	507	560	410	307	252	174	144	158	329	519	620	-
1931	4,886	606	591	577	450	357	275	191	142	163	283	412	539	-
1930	4,079	515	482	463	323	263	181	144	129	198	387	455	539	-
1929	4,712	662	673	702	491	326	227	130	102	141	303	456	499	-
1928	4,410	565	543	568	376	304	241	176	151	167	319	450	550	-
1927	3,922	469	467	471	376	317	226	150	89	162	269	402	524	-
1926	3,631	418	400	430	338	287	198	152	106	100	289	421	492	-
1925	3,608	505	463	517	334	299	193	119	98	136	228	299	417	-
1924	3,561	462	384	459	322	249	202	139	106	149	232	385	472	-
1923	3,388	539	395	390	300	219	183	140	91	129	223	326	433	-
1922	3,599	491	471	422	243	207	178	114	92	128	239	407	547	-
1921	3,866	540	453	419	308	213	211	159	128	168	314	466	487	-
1920	3,827	510	404	358	271	241	213	134	126	165	322	478	605	-

See footnotes at end of table.

TABLE 15. - 1/ DEATHS FROM SELECTED CAUSES BY MONTH: JAPAN, 1920 - 1949 - Cont'd
TETANUS (12)

Year	Total Deaths	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Un-known
* 1949	1,964	136	142	136	138	156	170	188	212	217	169	145	135	-
1948	2,138	152	117	138	205	176	181	237	234	196	152	183	167	-
1947	2,221	192	135	137	175	166	173	247	232	260	192	185	127	-
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1943	1,984	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1942	1,843	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1941	1,778	119	107	107	115	145	131	227	185	189	163	129	161	-
1940	1,841	136	112	121	149	152	154	191	194	174	183	131	144	-
1939	1,702	122	97	127	140	124	138	173	177	181	169	121	133	-
1938	1,775	133	122	126	132	157	150	165	183	190	145	134	137	-
1937	1,926	134	129	145	149	162	161	208	209	190	162	151	126	-
1936	1,989	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1935	2,127	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1934	2,192	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1933	2,296	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1932	2,445	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1931	2,430	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1930	2,501	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1929	2,668	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1928	2,774	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1927	2,756	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1926	2,864	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1925	2,901	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1924	2,986	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1923	3,183	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1922	2,894	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1921	3,059	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1920	3,096	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-

See footnotes at end of table.

TABLE 15. - $\frac{1}{2}$ DEATHS FROM SELECTED CAUSES BY MONTH: JAPAN, 1920 - 1949 - Cont'd
 $\frac{5}{2}$ TUBERCULOSIS (ALL FORMS) (13-22)

Year	Total Deaths	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Un-known
*1949	138,764	11,135	10,106	12,279	12,493	13,005	12,287	12,095	11,833	11,526	11,240	10,525	10,240	-
1948	143,909	10,946	10,741	12,424	12,862	13,147	12,677	12,776	12,488	11,914	11,810	11,106	11,008	10
1947	146,241	12,107	11,312	12,608	12,861	13,789	12,647	13,141	12,596	11,776	11,770	10,761	10,868	5
1946	198,900	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1945	203,000	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1944	176,800	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1943	171,473	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1942	161,484	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1941	154,344	12,251	11,275	12,290	12,477	12,959	12,811	13,797	14,159	13,497	13,460	12,560	12,808	
1940	153,154	12,605	11,934	12,882	12,151	12,821	12,437	13,505	13,742	13,195	13,193	12,327	12,362	
1939	154,371	12,110	11,053	12,785	12,058	12,513	12,632	13,961	14,118	13,588	13,747	12,728	13,078	
1938	148,827	11,847	10,877	12,201	11,961	12,454	12,445	13,552	13,566	13,167	12,749	12,043	11,964	1
1937	144,620	11,658	10,624	12,216	12,062	12,590	12,468	12,699	13,020	12,418	12,101	11,328	11,474	2
1936	145,160	11,932	11,152	11,836	11,694	12,531	12,266	12,984	12,788	12,442	12,436	11,506	11,593	
1935	132,151	10,530	9,479	10,713	10,462	11,290	11,153	11,749	12,019	11,519	11,667	10,671	10,899	
1934	131,525	10,568	9,790	11,163	10,856	11,700	11,242	11,531	11,729	11,047	11,303	10,416	10,179	1
1933	126,703	9,850	9,451	10,735	10,481	11,066	10,800	11,192	11,504	11,161	10,537	9,867	10,059	
1932	119,196	9,295	9,052	10,052	9,791	10,253	10,028	10,625	10,725	10,356	10,141	9,257	9,280	
1931	121,875	9,585	9,127	10,256	9,899	10,526	10,328	10,769	11,060	10,617	10,550	9,637	9,521	
1930	119,635	9,853	8,940	10,371	10,312	10,842	10,143	10,546	10,534	10,093	9,747	9,070	9,144	
1929	123,490	9,528	8,953	10,284	10,197	10,645	10,553	11,162	11,132	10,783	10,624	9,880	9,749	
1928	119,632	9,400	9,333	10,208	10,035	10,460	9,703	10,598	10,616	10,474	10,130	9,450	9,284	1
1927	119,439	8,704	8,299	9,511	9,858	10,824	10,283	10,982	11,280	10,675	10,520	9,339	9,162	2
1926	113,045	9,152	8,398	9,512	9,106	9,634	9,775	9,950	10,319	9,871	10,179	8,769	8,380	
1925	115,956	9,093	8,929	9,536	9,587	10,088	9,781	10,563	10,552	9,967	10,187	8,819	8,851	3
1924	114,229	8,934	8,552	9,556	9,020	10,108	9,585	10,367	10,478	10,141	9,942	8,724	8,621	1
1923	118,216	9,177	8,545	9,984	9,807	10,124	10,109	10,838	10,954	10,602	10,286	9,044	8,722	4
1922	125,506	9,337	9,394	11,185	10,685	11,374	10,744	11,084	11,819	11,010	10,484	9,330	9,060	
1921	120,719	8,998	8,773	9,863	9,685	10,369	9,968	10,866	11,524	11,055	10,943	9,558	9,117	
1920	125,165	10,530	9,723	10,604	10,304	10,691	10,189	11,121	11,544	11,224	10,636	9,447	9,152	

See footnotes at end of table.

TABLE 15. - 1/ DEATHS FROM SELECTED CAUSES BY MONTH: JAPAN, 1920 - 1949 - Cont'd
LEPROSY (23)

Year	Total Deaths	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Un-known
* 1949	162	17	15	12	18	14	12	12	6	9	15	19	13	-
1948	150	16	16	20	7	19	9	11	11	9	8	12	13	-
1947	189	18	17	20	23	17	7	14	13	17	18	12	13	-
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1943	131	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1942	169	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1941	208	15	12	20	25	20	20	NA	NA	16	15	12	12	-
1940	264	30	28	32	31	12	23	17	18	23	15	16	19	-
1939	312	34	35	37	32	19	22	27	16	25	22	21	22	-
1938	337	31	43	27	24	27	28	28	26	28	14	29	32	-
1937	400	44	21	51	33	33	31	25	32	34	39	25	32	-
1936	451	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1935	482	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1934	482	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1933	514	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1932	541	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1931	669	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1930	640	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1929	637	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1928	717	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1927	759	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1926	779	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1925	789	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1924	823	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1923	920	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1922	976	99	80	112	72	82	85	67	90	72	78	72	67	-
1921	1,094	90	106	97	87	72	82	93	89	111	98	84	85	-
1920	1,118	103	123	111	95	81	77	79	86	103	83	78	99	-

See footnotes at end of table.

TABLE 15. - 2/ DEATHS FROM SELECTED CAUSES BY MONTH: JAPAN, 1920 - 1949 - Cont'd
GLANDERS (26a)

Year	Total Deaths	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Un- known
1949	1	-	-	-	-	1	-	-	-	-	-	-	-	-
1948	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1947	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1943	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1942	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1941	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1940	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1939	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1938	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1937	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1936	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1935	1	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1934	2	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1933	-	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1932	-	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1931	1	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1930	1	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1929	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1928	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1927	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1926	1	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1925	5	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1924	16	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1923	21	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1922	1	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1921	2	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1920	-	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-

See footnotes at end of table.

TABLE 15. - $\frac{1}{2}$ DEATHS FROM SELECTED CAUSES BY MONTH: JAPAN, 1920 - 1949 - Cont'd
DYSENTERY (27)

Year	Total Deaths	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Un-known
*1949	7,824	47	54	60	111	225	534	1,850	2,430	1,638	569	191	115	-
1948	5,157	38	55	67	125	194	453	1,392	1,487	831	344	105	66	-
1947	9,573	113	91	102	152	285	583	2,626	3,233	1,586	563	173	66	-
1946	13,198	133	45	88	77	148	396	1,505	4,077	3,038	2,175	1,123	393	-
1945	20,107	49	46	59	96	186	418	2,117	4,789	6,068	3,528	1,927	824	-
1944	11,208	38	74	80	114	229	547	1,861	3,152	2,380	1,482	791	460	-
1943	12,641	112	116	149	211	364	779	2,092	2,693	1,863	1,130	415	284	-
1942	14,283	172	222	292	384	731	1,003	2,290	2,314	2,212	1,372	825	520	-
1941	16,325	316	284	379	551	929	1,740	3,743	3,875	2,385	1,282	516	325	-
1940	22,032	503	512	619	759	1,273	2,380	5,064	5,116	2,985	1,646	752	423	-
1939	24,907	436	365	533	637	1,110	2,012	4,594	6,003	4,240	2,915	1,404	658	-
1938	21,966	326	304	465	666	1,007	2,062	4,663	5,391	3,621	2,009	891	561	-
1937	19,726	363	355	468	663	1,019	1,634	3,557	4,263	3,476	2,404	1,060	464	-
1936	16,721	259	252	414	538	785	1,420	3,345	4,242	2,770	1,540	732	424	-
1935	15,921	394	298	423	543	897	1,522	3,459	3,590	2,367	1,403	677	348	-
1934	15,490	218	234	373	527	815	1,396	3,325	3,619	2,419	1,545	658	361	-
1933	14,880	262	265	306	403	722	1,415	2,938	3,526	2,504	1,541	637	361	-
1932	13,550	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1931	13,005	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1930	13,019	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1929	13,427	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1928	12,044	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1927	10,217	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1926	8,530	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1925	7,515	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1924	8,658	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1923	9,400	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1922	9,116	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1921	7,879	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1920	8,155	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-

See footnotes at end of table.

TABLE 15. - 1/ DEATHS FROM SELECTED CAUSES BY MONTH: JAPAN, 1920 - 1949 - Cont'd
MALARIA (28)

Year	Total Deaths	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Un-known
* 1949	122	13	11	19	7	9	9	9	8	17	6	9	5	-
1948	224	23	26	23	21	27	11	15	20	26	19	10	3	-
1947	456	43	35	38	44	42	47	36	35	38	31	33	34	-
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1943	245	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1942	220	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1941	153	17	12	11	10	5	12	6	12	22	15	14	15	-
1940	224	22	28	16	18	26	9	19	17	17	19	18	15	-
1939	222	36	24	11	9	17	11	12	19	22	24	22	15	-
1938	207	6	8	9	6	8	3	13	30	25	36	26	37	-
1937	65	2	5	2	7	5	6	5	10	4	7	6	6	-
1936	76	-	6	5	-	4	9	14	13	8	3	6	8	-
1935	51	4	1	2	1	5	3	4	8	6	7	7	3	-
1934	53	4	9	4	5	4	3	2	6	3	4	4	5	-
1933	74	7	8	4	6	3	7	7	10	3	5	7	7	-
1932	68	5	3	1	1	5	6	7	6	9	13	4	8	-
1931	63	4	6	6	5	8	6	1	7	5	5	2	8	-
1930	60	3	4	3	5	5	11	7	5	2	5	3	7	-
1929	97	9	5	10	4	7	20	6	10	11	9	3	3	-
1928	164	7	6	7	5	5	7	15	33	34	21	15	9	-
1927	172	13	6	7	11	4	17	15	37	19	21	18	4	-
1926	190	9	9	4	9	11	11	9	18	25	39	23	25	-
1925	122	9	5	5	9	5	13	12	11	23	17	6	7	-
1924	150	15	9	7	9	6	24	10	17	23	8	14	8	-
1923	169	10	5	4	5	18	17	18	27	20	16	19	10	-
1922	191	8	10	6	10	4	16	24	39	30	19	9	16	-
1921	219	20	4	10	11	19	20	31	39	20	22	11	12	-
1920	243	22	7	8	8	10	15	37	48	30	22	17	19	-

See footnotes at end of table.

TABLE 15. - 1/ DEATHS FROM SELECTED CAUSES BY MONTH: JAPAN, 1920 - 1949 - Cont'd
2/ SYPHILIS (30)

Year	Total Deaths	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Un-known
* 1949	5,492	518	467	518	467	397	390	404	393	399	477	515	547	-
** 1948	4,356	358	401	417	337	321	309	318	322	334	358	388	493	-
** 1947	4,444	524	416	409	344	354	370	364	300	290	336	361	376	-
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1943	7,854	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1942	7,930	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1941	7,772	739	699	687	624	586	556	584	550	614	682	652	799	-
1940	7,706	873	682	747	610	552	531	535	568	625	644	632	707	-
1939	8,067	832	695	697	615	571	579	606	628	584	762	698	800	-
1938	8,749	795	795	795	651	673	560	675	676	696	767	763	740	-
1937	9,440	898	771	898	778	717	654	713	735	746	764	816	950	-
1936	9,402	1,026	908	900	752	646	666	684	687	680	800	763	889	-
1935	9,323	891	841	857	706	692	635	664	705	754	777	835	966	-
1934	9,367	975	782	894	800	670	632	685	652	662	850	890	875	-
1933	9,553	951	922	879	717	705	718	697	724	731	777	837	895	-
1932	9,340	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1931	9,868	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1930	9,797	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1929	10,313	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1928	10,956	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1927	10,884	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1926	11,110	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1925	10,722	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1924	10,920	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1923	11,410	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1922	9,282	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1921	10,416	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1920	10,170	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-

See footnotes at end of table.

TABLE 15. - 1/ DEATHS FROM SELECTED CAUSES BY MONTH: JAPAN, 1920 - 1949 - Cont'd
INFLUENZA (33)

Year	Total Deaths	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Un- known
* 1949	502	57	78	62	53	51	25	13	16	20	19	32	76	-
1948	515	86	64	91	55	27	33	23	16	19	29	25	47	-
1947	1,903	429	684	344	127	79	48	34	14	17	32	28	67	-
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1943	3,412	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1942	3,981	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1941	4,351	916	1,330	831	333	177	101	66	62	59	81	136	259	-
1940	3,164	636	713	564	300	163	120	59	57	38	69	107	338	-
1939	5,848	839	1,178	1,845	911	292	117	57	61	41	85	112	310	-
1938	7,646	1,490	2,508	1,872	641	289	108	55	57	84	84	113	345	-
1937	2,991	438	534	446	312	190	121	70	56	82	101	148	493	-
1936	8,670	2,612	3,251	1,447	470	218	88	67	38	53	80	109	237	-
1935	3,070	512	523	561	340	217	98	80	89	62	96	122	370	-
1934	10,142	2,624	3,038	2,364	842	365	143	72	54	50	100	162	328	-
1933	4,765	607	764	1,086	748	374	149	73	61	78	109	166	550	-
1932	5,370	582	699	1,135	1,209	668	211	102	79	94	113	174	344	-
1931	15,673	3,030	4,510	3,564	1,828	1,124	495	197	96	75	130	192	432	-
1930	5,207	613	721	989	868	637	288	124	115	110	119	179	444	-
1929	8,480	1,034	1,411	2,299	1,465	776	308	169	110	135	153	194	426	-
1928	9,669	883	1,292	3,298	2,063	828	258	110	98	94	151	217	577	-
1927	8,151	1,368	1,927	1,505	926	891	458	177	130	116	144	164	345	-
1926	4,610	602	654	769	590	502	250	118	91	110	165	242	516	1
1925	10,806	1,274	4,142	3,000	907	344	181	100	73	106	137	183	358	1
1924	6,512	751	1,165	1,449	943	640	368	150	163	148	160	235	340	-
1923	6,520	720	1,218	1,438	980	596	332	188	112	138	188	210	400	-
1922	12,688	1,167	4,072	4,157	1,399	543	245	154	116	140	142	205	348	-
1921	10,304	1,215	1,496	2,140	1,820	956	432	287	231	281	329	472	645	-
1920	108,428	39,562	31,006	17,726	10,780	4,987	1,260	450	302	355	458	581	961	-

See footnotes at end of table.

TABLE 15. - 1/ DEATHS FROM SELECTED CAUSES BY MONTH: JAPAN, 1920 - 1949 - Cont'd
SMALLPOX (34)

Year	Total Deaths	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Un- Known
*1949	21	-	2	2	5	6	3	1	1	-	1	-	-	-
1948	3	-	-	1	-	-	1	1	-	-	-	-	-	-
1947	85	14	6	12	14	8	11	8	4	2	1	2	3	-
1946	2,823	109	374	747	696	485	235	43	8	17	5	17	96	-
1945	253	6	5	3	20	53	42	17	26	18	7	28	28	-
1944	44	1	3	13	3	14	6	1	1	-	1	-	1	-
1943	74	2	11	7	24	15	8	-	-	1	1	2	2	-
1942	61	1	20	20	6	3	6	-	-	-	-	-	-	-
1941	92	11	11	14	9	15	20	5	1	2	2	1	1	-
1940	60	3	2	18	22	4	8	1	-	1	-	-	1	-
1939	33	3	7	7	3	8	2	3	-	-	-	-	-	-
1938	6	-	-	-	-	-	-	-	-	-	-	1	5	-
1937	8	-	3	2	1	2	-	-	-	-	-	-	-	-
1936	20	1	-	-	3	5	3	1	1	3	1	-	2	-
1935	16	4	1	2	1	4	3	-	-	-	-	1	-	-
1934	35	3	2	4	6	5	1	-	-	-	-	2	12	-
1933	55	2	11	8	11	12	5	6	-	-	-	-	-	-
1932	44	2	1	5	11	20	4	-	-	-	1	-	-	-
1931	8	1	-	2	2	1	-	-	-	-	-	-	2	-
1930	4	-	-	1	1	1	-	-	-	1	-	-	-	-
1929	11	-	-	7	2	1	1	-	-	-	-	-	-	-
1928	100	13	29	17	26	8	5	-	-	-	-	1	-	-
1927	90	8	10	6	4	9	8	8	7	4	4	2	19	-
1926	158	3	11	32	31	49	15	5	1	2	-	-	9	-
1925	69	1	2	5	12	21	18	6	2	-	1	-	1	-
1924	294	89	69	42	49	19	10	10	4	2	-	-	-	-
1923	999	23	78	59	96	49	58	23	5	5	2	8	53	-
1922	120	10	17	12	23	21	23	7	1	1	-	1	4	-
1921	203	5	19	58	38	37	21	17	2	1	-	2	3	-
1920	729	58	82	124	162	160	91	37	8	3	2	-	2	-

See footnotes at end of table.

TABLE 15. - 1/ DEATHS FROM SELECTED CAUSES BY MONTH: JAPAN, 1920 - 1949 - Cont'd
MEASLES (35)

Year	Total Deaths	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Un-known
* 1949	12,440	575	681	1,403	1,787	2,758	2,209	1,442	679	233	169	229	275	-
1948	5,598	410	411	706	556	731	742	610	377	183	164	277	431	-
1947	20,939	1,248	2,133	3,035	3,136	3,355	3,086	2,205	1,144	418	299	360	520	-
1946	MA	MA	MA	MA	MA	MA	MA	MA	MA	MA	MA	MA	MA	-
1945	MA	MA	MA	MA	MA	MA	MA	MA	MA	MA	MA	MA	MA	-
1944	MA	MA	MA	MA	MA	MA	MA	MA	MA	MA	MA	MA	MA	-
1943	21,881	MA	MA	MA	MA	MA	MA	MA	MA	MA	MA	MA	MA	-
1942	6,794	MA	MA	MA	MA	MA	MA	MA	MA	MA	MA	MA	MA	-
1941	14,616	449	692	1,514	2,176	3,264	2,691	1,597	705	322	329	373	504	-
1940	5,101	376	324	490	529	783	765	710	353	155	155	187	274	-
1939	14,680	492	922	1,880	2,412	3,103	2,598	1,649	634	244	169	246	331	-
1938	4,997	321	309	445	512	923	851	639	287	125	118	187	280	-
1937	10,889	489	690	1,190	1,787	2,209	1,750	1,363	554	182	212	212	281	-
1936	5,264	318	322	428	547	858	956	628	327	126	138	243	373	-
1935	9,816	663	825	1,400	1,595	1,752	1,440	867	405	146	188	222	313	-
1934	9,787	632	716	1,258	1,491	1,893	1,379	935	424	180	164	289	426	-
1933	8,442	383	426	975	1,276	1,733	1,352	919	393	139	133	287	426	-
1932	6,220	573	593	861	859	1,061	753	632	288	113	86	136	265	-
1931	13,691	721	916	1,403	1,856	2,553	2,579	1,462	810	292	221	343	535	-
1930	5,965	520	449	613	688	878	889	674	351	110	164	237	392	-
1929	16,070	1,390	1,768	2,698	2,364	2,609	2,133	1,326	578	218	223	353	410	-
1928	11,122	909	965	1,255	1,158	1,595	1,569	1,219	585	278	237	491	861	-
1927	13,472	749	857	1,398	1,615	2,317	2,145	1,638	789	314	346	532	772	-
1926	10,510	618	635	809	1,166	1,831	1,786	1,329	691	279	267	462	637	-
1925	15,438	1,206	1,340	1,953	2,289	2,762	2,312	1,399	697	310	253	405	512	-
1924	11,986	657	773	1,253	1,495	1,984	1,761	1,421	682	300	305	552	803	-
1923	13,277	833	790	1,157	1,563	2,261	2,256	1,675	1,020	384	368	425	545	-
1922	9,698	1,028	795	925	950	1,384	1,400	1,029	603	211	216	391	726	-
1921	13,894	754	867	1,233	1,586	2,160	2,049	1,806	1,024	454	422	683	856	-
1920	7,502	970	820	720	747	840	921	730	406	206	205	345	592	-

See footnotes at end of table.

TABLE 15. - 1/ DEATHS FROM SELECTED CAUSES BY MONTH: JAPAN, 1920 - 1949 - Cont'd
ACUTE POLIOENCEPHALITIS (36)

Year	Total Deaths	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Un- known
*1949	1,024	76	60	75	67	81	89	109	81	110	113	79	84	-
1948	775	64	52	59	52	47	47	80	96	88	74	51	65	-
1947	1,009	66	92	93	73	92	85	128	126	78	57	52	67	-
1920-1946	Not Available.													

JAPANESE "B" ENCEPHALITIS (37b)

Year	Total Deaths	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Un- known
* 1949	1,182	3	6	8	2	5	14	24	153	774	169	21	3	-
1948	2,620	2	2	2	2	-	7	24	1,657	786	99	26	13	-
1947	228	3	4	1	6	9	9	13	56	86	38	3	-	-
1920-1946	Not Available.													

See footnotes at end of table.

TABLE 15. - 1/ DEATHS FROM SELECTED CAUSES BY MONTH: JAPAN, 1920 - 1949 - Cont'd
YELLOW FEVER (38a)

Year	Total Deaths	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Un- known
*1949	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1948	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1947	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1943	7	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1942	1	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1941	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1940	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1939	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1938	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1937	1	-	-	-	-	-	-	-	-	-	-	1	-	-
1936	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1935	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1934	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1933	1	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1932	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1931	1	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1930	2	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1929	1	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1928	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1927	2	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1926	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1925	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1924	2	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1923	5	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1922	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1921	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1920	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-

See footnotes at end of table.

TABLE 15. - 1/ DEATHS FROM SELECTED CAUSES BY MONTH: JAN., 1920 - 1949 - Cont'd
RABIES (38b)

Year	Total Deaths	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Un- known
*1949	76	2	4	3	6	6	5	6	12	10	10	6	6	-
1948	40	4	5	-	1	1	1	3	5	6	3	4	7	-
1947	16	1	2	-	2	1	-	2	-	-	3	3	2	-
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1943	6	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1942	3	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1941	4	-	-	-	-	1	1	-	1	1	-	-	-	-
1940	6	-	-	1	-	-	-	1	2	-	1	1	-	-
1939	2	-	-	1	-	-	-	-	-	-	-	-	1	-
1938	3	-	-	-	1	2	-	-	-	-	-	-	-	-
1937	4	-	-	-	1	1	-	1	1	-	-	-	-	-
1936	3	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1935	4	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1934	2	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1933	3	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1932	4	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1931	1	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1930	4	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1929	12	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1928	22	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1927	30	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1926	80	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1925	143	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1924	235	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1923	174	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1922	79	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1921	54	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1920	48	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-

See footnotes at end of table.

TABLE 15. - 1/ DEATHS FROM CORRECTED CAUSES BY MONTH: JAPAN, 1920 - 1949 - Cont'd
2/ TYPHUS FEVER AND TYPHUS LIKE DISEASES (39)

Year	Total Deaths	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Un- known
*1949	67	4	8	7	5	6	10	3	9	8	3	2	2	-
1948	93	14	12	6	6	10	6	8	9	5	5	6	6	-
1947	173	26	13	20	27	15	15	15	11	7	5	2	17	-
1946	2,909	32	158	740	722	587	392	117	36	10	79	18	18	-
1945	259	5	11	10	33	64	56	17	26	6	8	3	20	-
1944	622	52	58	74	68	120	125	49	50	16	6	2	2	-
1943	183	-	1	1	5	15	34	9	2	3	15	17	27	-
1942	66	1	2	-	8	2	4	1	-	1	-	-	2	-
1941	37	2	-	2	1	2	7	5	10	5	-	2	1	-
1940	41	-	-	-	1	1	2	14	17	3	1	1	1	-
1939	38	-	-	-	-	-	-	6	14	10	5	2	1	-
1938	39	-	-	-	-	-	-	11	17	6	5	-	-	-
1937	32	-	-	-	-	-	1	7	14	3	5	2	-	-
1936	25	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1935	38	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1934	47	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1933	47	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1932	33	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1931	22	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1930	16	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1929	18	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1928	14	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1927	31	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1926	20	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1925	26	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1924	14	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1923	39	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1922	43	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1921	77	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1920	90	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-

See footnotes at end of table.

TABLE 15. - 1/ DEATHS FROM SELECTED CAUSES BY MONTH: JAPAN, 1920 - 1949 - Cont'd
CANCER AND OTHER MALIGNANT TUMORS (45-55)

Year	Total Deaths	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Un-known
1949	58,890	4,741	4,239	4,737	4,647	4,910	4,837	5,137	5,318	5,155	5,126	4,916	5,127	-
1948	55,677	4,292	3,997	4,373	4,370	4,586	4,642	4,916	4,885	4,942	5,075	4,799	4,806	3
1947	53,047	4,336	3,772	4,265	4,189	4,251	4,438	4,696	4,708	4,645	4,823	4,465	4,459	-
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1943	52,678	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1942	52,936	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1941	51,972	4,118	3,720	4,142	4,026	4,257	4,376	4,707	4,715	4,700	4,602	4,190	4,419	-
1940	50,940	4,123	3,689	3,997	3,959	4,069	4,174	4,737	4,839	4,529	4,366	4,219	4,239	-
1939	51,129	4,116	3,588	4,170	3,927	4,262	4,302	4,715	4,595	4,522	4,491	4,182	4,259	-
1938	50,447	3,974	3,638	3,874	3,892	4,076	4,238	4,593	4,755	4,583	4,565	4,195	4,064	-
1937	50,648	4,001	3,649	3,956	3,939	4,284	4,270	4,509	4,595	4,519	4,582	4,187	4,157	-
1936	49,212	4,108	3,574	3,753	3,703	4,053	4,033	4,391	4,498	4,390	4,543	4,183	3,983	-
1935	49,111	3,826	3,520	3,696	3,813	4,096	4,154	4,505	4,508	4,320	4,525	4,099	4,049	-
1934	47,972	3,853	3,354	3,624	3,662	3,903	4,088	4,387	4,407	4,408	4,431	3,975	3,880	-
1933	46,879	3,616	3,377	3,635	3,610	4,004	3,964	4,237	4,365	4,295	4,375	3,717	3,684	-
1932	45,288	3,424	3,184	3,602	3,486	3,670	3,786	4,188	4,064	4,177	4,111	3,814	3,752	-
1931	44,541	3,538	3,263	3,597	3,519	3,730	3,687	3,956	4,061	3,948	4,060	3,673	3,609	-
1930	44,902	3,554	3,179	3,560	3,497	3,763	3,774	4,126	4,162	4,091	3,962	3,637	3,597	-
1929	43,734	3,449	3,043	3,408	3,521	3,578	3,755	3,956	4,093	3,896	3,921	3,544	3,570	-
1928	44,534	3,430	3,243	3,496	3,437	3,659	3,690	3,944	4,149	4,025	4,159	3,723	3,479	-
1927	42,835	3,284	3,094	3,400	3,325	3,617	3,649	3,906	3,682	3,877	3,798	3,514	3,389	-
1926	42,588	3,355	2,925	3,271	3,318	3,571	3,562	3,871	4,035	3,937	3,871	3,428	3,458	-
1925	41,744	3,163	2,947	3,169	3,221	3,553	3,542	3,822	3,915	3,746	3,912	3,396	3,358	-
1924	41,312	3,111	2,992	3,242	3,154	3,425	3,483	3,904	3,904	3,816	3,715	3,323	3,243	-
1923	41,810	3,181	2,804	3,203	3,565	3,671	3,671	3,859	3,947	3,938	3,908	3,276	3,188	-
1922	40,813	3,095	2,798	3,223	3,084	3,457	3,412	3,710	3,895	3,782	3,841	3,353	3,163	-
1921	40,578	3,051	2,739	3,107	3,068	3,164	3,266	3,704	3,961	3,821	3,897	3,364	3,136	-
1920	40,328	3,131	2,855	3,055	3,025	3,357	3,348	3,581	4,013	3,930	3,668	3,208	3,162	-

See footnotes at end of table.

TABLE 15. - 1/ DEATHS FROM SELECTED CAUSES BY MONTH: JAPAN, 1920 - 1949 - Cont'd
DIABETES MELLITUS (61)

Year	Total Deaths	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Un- known
* 1949	1,885	201	179	202	175	151	130	139	127	129	152	141	159	-
1948	1,789	217	196	140	169	139	117	137	135	119	132	131	157	-
1947	1,827	183	203	196	149	143	135	143	109	134	125	120	186	1
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1943	2,477	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1942	2,619	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1941	2,657	296	310	237	208	212	151	204	202	197	177	220	243	-
1940	2,762	268	292	269	246	233	207	180	202	207	203	201	254	-
1939	2,795	299	259	281	237	217	183	200	216	209	239	212	242	1
1938	3,043	335	260	312	249	244	218	221	228	209	244	238	285	-
1937	2,812	266	245	252	206	207	204	218	223	223	240	238	290	-
1936	2,629	309	307	278	222	193	185	183	184	196	209	182	181	-
1935	2,527	252	245	222	217	198	182	196	215	180	175	196	249	-
1934	2,718	296	280	287	243	216	199	192	185	177	215	211	217	-
1933	2,989	274	246	267	219	200	172	177	193	184	209	219	229	-
1932	2,331	215	220	209	221	165	181	177	175	197	189	173	209	-
1931	2,209	249	213	238	182	221	160	158	147	125	162	150	204	-
1930	2,247	255	208	207	182	199	167	169	140	162	175	187	196	-
1929	2,300	222	210	251	206	174	170	158	172	188	169	189	191	-
1928	2,171	210	206	209	204	168	162	161	142	152	172	155	230	-
1927	2,173	216	238	219	220	173	147	167	139	156	173	143	172	-
1926	2,046	188	180	189	168	159	164	172	164	172	164	172	178	1
1925	1,979	222	198	204	166	162	153	137	137	133	176	137	153	1
1924	1,963	187	160	186	170	152	148	159	153	145	169	159	175	-
1923	1,873	191	186	152	164	157	136	162	133	136	170	127	159	-
1922	1,904	200	197	174	158	143	156	119	132	146	146	156	177	-
1921	1,890	184	164	163	156	145	145	159	145	170	163	135	161	-
1920	1,725	170	180	164	141	120	134	122	129	138	121	155	161	-

See footnotes at end of table.

TABLE 15. - 1/ DEATHS FROM SELECTED CAUSES BY MONTH: JAPAN, 1920 - 1949 - Cont'd
HERIBERI (68)

Year	Total Deaths	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Un- known
*1949	5,591	597	539	603	553	474	411	373	412	361	381	394	493	-
1948	6,281	710	717	658	566	470	471	458	486	424	407	427	487	-
1947	8,596	805	767	895	820	750	633	679	635	563	604	614	831	-
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1943	6,480	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1942	6,684	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1941	6,677	687	603	671	557	494	441	493	453	529	509	551	689	-
1940	7,186	822	687	701	573	508	522	527	607	740	526	553	620	-
1939	9,567	977	785	745	665	614	580	727	916	909	938	823	888	-
1938	12,712	948	747	756	683	748	789	1,033	1,472	1,703	1,585	1,220	1,028	-
1937	10,627	725	653	661	610	593	595	740	1,050	1,217	1,449	1,246	1,088	-
1936	11,097	746	689	788	641	613	708	952	1,211	1,446	1,506	971	826	-
1935	10,062	656	572	639	539	488	529	717	1,019	1,311	1,635	1,076	881	-
1934	11,494	716	597	663	620	637	682	1,003	1,178	1,471	1,847	1,272	808	-
1933	13,828	805	636	685	547	770	840	1,321	1,581	2,279	2,160	1,273	931	-
1932	11,805	729	653	656	571	602	633	1,058	1,396	1,628	1,815	1,210	854	-
1931	17,767	652	625	802	870	1,153	1,006	1,674	2,480	2,850	3,049	1,604	1,002	-
1930	15,419	808	649	763	856	1,144	1,099	1,917	2,182	2,430	1,851	1,010	710	-
1929	15,456	838	706	702	762	957	1,193	1,832	2,249	1,957	1,908	1,403	949	-
1928	19,036	595	520	518	590	708	873	1,661	2,899	3,950	3,455	2,115	1,172	-
1927	12,058	567	446	488	502	646	778	1,283	1,664	1,833	1,921	1,182	748	-
1926	12,109	674	492	632	531	673	888	1,168	1,690	1,865	1,862	986	648	-
1925	13,909	687	558	562	579	788	1,019	1,588	1,796	2,111	2,179	1,228	814	-
1924	18,333	788	680	763	663	998	1,195	2,284	2,959	2,987	2,576	1,516	924	-
1923	26,796	633	561	680	931	1,348	1,542	2,819	5,101	5,670	4,296	2,094	1,120	1
1922	19,162	654	474	628	658	1,063	1,213	2,095	3,290	3,644	3,078	1,521	844	-
1921	21,675	392	348	356	344	639	849	2,186	4,567	5,224	4,584	2,208	978	-
1920	14,239	331	266	258	367	489	558	1,429	2,713	3,707	2,473	1,085	563	-

See footnotes at end of table.

TABLE 15. - 1/ DEATHS FROM SELECTED CAUSES BY MONTH: JAPAN, 1920 - 1949 - Cont'd
g/ SIMPLE MENINGITIS (81a)

Year	Total Deaths	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Un- known
* 1949	9,108	767	706	870	919	889	799	760	771	753	645	615	654	-
** 1948	10,414	984	889	1,004	1,065	1,037	927	987	860	649	662	652	698	-
1947	14,143	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1943	30,088	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1942	29,723	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1941	30,014	2,120	2,109	2,494	2,505	2,761	2,761	3,342	3,122	2,411	2,266	2,096	2,087	-
1940	31,609	2,366	2,365	2,753	2,670	2,926	2,909	3,631	3,261	2,584	2,219	1,962	1,963	-
1939	35,631	2,416	2,350	2,996	2,877	3,024	3,271	3,737	4,538	3,367	2,663	2,238	2,154	-
1938	36,748	2,664	2,582	2,998	2,998	3,155	3,188	4,061	4,207	3,649	2,652	2,308	2,286	-
1937	36,188	2,410	2,295	2,786	2,917	3,085	3,077	3,959	4,882	3,368	2,731	2,320	2,358	-
1936	36,905	2,599	2,655	3,094	3,174	3,122	3,351	3,913	4,128	3,542	2,736	2,196	2,395	-
1935	37,615	2,682	2,501	3,059	3,085	3,215	3,308	4,090	4,963	3,213	2,769	2,320	2,410	-
1934	39,924	2,938	2,707	3,356	3,284	3,620	3,686	4,512	4,219	3,458	2,925	2,596	2,623	-
1933	42,546	3,032	2,861	3,388	3,398	3,818	3,790	4,891	5,179	3,735	3,163	2,661	2,630	-
1932	43,586	3,082	3,084	3,552	3,346	3,716	3,837	5,011	5,320	3,944	3,140	2,696	2,858	-
1931	46,296	3,139	3,166	3,645	3,710	4,071	4,071	4,850	5,428	4,549	3,703	3,058	2,905	1
1930	47,532	3,417	3,212	3,722	3,725	3,903	4,301	5,451	6,275	4,257	3,431	2,899	2,939	-
1929	54,097	3,615	3,470	4,333	4,184	4,306	4,750	6,407	7,751	4,869	3,926	3,331	3,155	-
1928	53,388	3,706	3,596	4,159	4,071	4,430	4,538	6,129	6,157	5,564	4,203	3,385	3,450	-
1927	55,420	3,580	3,571	4,161	4,179	4,499	4,640	6,586	7,705	5,620	4,172	3,273	3,433	1
1926	56,759	3,767	3,471	4,192	4,372	4,787	4,859	7,995	7,682	6,362	4,281	3,558	3,432	1
1925	58,540	4,109	4,063	4,704	4,683	5,027	5,233	6,421	6,775	5,893	4,490	3,568	3,574	-
1924	68,466	4,581	4,398	5,083	5,032	5,227	5,659	8,189	12,151	6,153	4,437	3,708	3,847	1
1923	73,489	4,598	4,239	5,186	5,150	5,388	5,997	8,105	10,530	9,423	6,181	4,382	4,310	-
1922	69,585	4,568	4,731	5,140	4,805	5,136	5,452	7,428	11,140	8,136	5,210	3,956	3,883	-
1921	68,293	4,189	4,271	4,960	4,984	5,058	5,290	7,504	9,609	8,124	5,591	4,370	4,343	-
1920	69,207	5,016	5,019	5,617	5,014	5,205	5,304	7,852	9,876	7,365	4,738	4,015	4,186	-

See footnotes at end of table.

TABLE 15. - 1/ DEATHS FROM SELECTED CAUSES BY MONTH: JAPAN, 1920 - 1949 - Cont'd
9/ INTRACRANIAL LESIONS OF VASCULAR ORIGIN (83)

Year	Total Deaths	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Un- known
* 1949	100,650	9,581	8,446	9,811	9,006	7,947	7,358	7,133	6,724	7,713	8,517	8,753	9,661	-
1948	94,329	10,063	9,418	9,454	7,858	7,462	6,686	6,148	6,500	7,009	7,257	7,884	8,581	9
1947	101,095	10,805	10,856	10,713	8,740	8,031	7,050	6,548	6,014	6,504	7,747	8,215	9,870	2
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1943	120,985	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1942	125,349	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1941	125,124	12,180	11,405	11,568	10,670	9,989	9,249	9,300	9,043	9,834	10,326	10,164	11,396	-
1940	127,847	13,393	12,837	12,674	10,872	10,424	9,150	8,756	9,359	9,623	9,951	9,722	11,086	-
1939	130,826	13,211	11,390	12,352	10,623	10,244	9,388	9,177	10,608	10,453	10,885	10,368	12,127	-
1938	126,861	12,987	11,807	11,369	10,102	9,515	8,977	9,237	9,474	11,106	10,411	10,500	11,376	-
1937	118,761	11,031	9,596	10,751	9,293	8,623	8,623	8,154	10,064	9,717	10,289	9,737	11,951	-
1936	118,152	12,636	11,393	11,536	9,907	9,147	8,317	8,192	8,154	9,268	9,933	9,418	10,251	-
1935	114,554	10,937	9,807	10,376	9,196	8,967	8,309	8,119	10,126	9,283	9,308	9,292	10,834	-
1934	114,447	12,017	10,689	11,219	9,805	8,813	8,158	7,768	8,038	8,412	9,839	9,587	10,101	1
1933	110,719	11,195	10,214	10,837	9,024	8,605	8,069	7,476	8,629	8,517	9,087	8,916	10,150	-
1932	107,378	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1931	107,352	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1930	104,942	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1929	108,439	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1928	102,985	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1927	101,705	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1926	98,688	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1925	96,293	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1924	102,810	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1923	94,615	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1922	91,433	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1921	90,523	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1920	88,186	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	

See footnotes at end of table.

TABLE 15. - 1/ DEATHS FROM SELECTED CAUSES BY MONTH: JAPAN, 1920 - 1949 - Cont'd
10/ ALL HEART DISEASES (90-95)

Year	Total Deaths	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Un- known
1949	52,825	5,111	4,572	5,229	4,878	4,186	3,953	3,864	3,677	3,655	4,221	4,408	5,071	-
1948	49,046	4,788	4,741	4,914	4,132	3,930	3,744	3,670	3,445	3,466	3,818	4,077	4,320	1
1947	48,575	4,606	4,700	4,885	4,060	4,083	3,684	3,534	3,373	3,201	3,778	3,874	4,796	1
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1943	45,428	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1942	43,487	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1941	42,543	4,655	4,330	4,083	3,600	3,358	3,044	3,062	2,840	3,032	3,265	3,345	3,099	-
1940	45,542	5,127	4,893	4,593	3,999	3,640	3,373	3,235	3,024	3,086	3,305	3,274	4,033	-
1939	47,442	5,402	4,581	5,213	3,947	3,599	3,337	3,374	3,242	3,140	3,538	3,575	4,494	-
1938	47,461	5,226	4,954	4,560	3,768	3,541	3,226	3,463	3,351	3,409	3,731	3,879	4,353	-
1937	42,822	4,312	3,739	3,845	3,596	3,409	3,140	3,138	3,033	3,141	3,546	3,461	4,462	-
1936	42,910	5,179	4,706	4,193	3,391	3,194	3,122	3,067	2,853	2,897	3,365	3,279	3,659	-
1935	39,902	3,872	3,686	3,691	3,206	3,116	3,064	2,855	2,826	2,968	3,331	3,270	3,957	-
1934	42,519	4,732	4,245	4,318	3,563	3,259	2,960	2,929	2,881	2,928	3,555	3,446	3,703	-
1933	40,111	4,032	3,791	4,133	3,350	3,117	2,936	2,862	2,704	2,983	3,240	3,286	3,677	-
1932	38,973	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1931	41,867	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1930	41,138	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1929	41,532	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1928	39,907	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1927	38,971	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1926	37,949	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1925	39,895	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1924	40,676	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1923	42,184	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1922	37,312	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1921	37,420	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1920	35,540	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	

See footnotes at end of table.

TABLE 15. - $\frac{1}{2}$ DEATHS FROM COLLECTED CAUSES BY MONTH: JAPAN, 1920 - 1949 - Cont'd
 $\frac{1}{2}$ BRONCHITIS (106)

Year	Total Deaths	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Un-known
*1949	25,695	3,109	3,060	3,370	2,844	2,153	1,688	1,389	1,136	1,142	1,471	1,749	2,584	-
1948	25,932	3,730	3,488	3,266	2,465	1,964	1,582	1,263	1,136	1,324	1,603	1,744	2,347	-
1947	34,923	3,229	4,057	5,370	4,058	3,075	2,448	2,029	1,701	1,439	1,848	2,355	3,314	-
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1943	23,268	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1942	22,416	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1941	22,047	2,601	3,068	2,779	2,090	1,701	1,427	1,299	1,077	1,181	1,380	1,515	1,929	-
1940	22,572	3,032	3,108	2,804	2,103	1,711	1,399	1,276	1,184	1,215	1,340	1,391	2,009	-
1939	25,115	3,148	3,012	3,402	2,636	1,806	1,579	1,372	1,299	1,284	1,536	1,679	2,362	-
1938	26,178	3,439	3,519	3,458	2,442	1,984	1,457	1,509	1,346	1,411	1,537	1,780	2,295	-
1937	24,030	2,640	2,581	2,721	2,204	1,924	1,550	1,411	1,341	1,389	1,684	1,803	2,782	-
1936	26,120	3,780	3,933	3,438	2,391	1,881	1,543	1,490	1,175	1,244	1,541	1,645	2,099	-
1935	24,240	2,946	2,854	2,807	2,279	1,900	1,555	1,379	1,325	1,387	1,613	1,803	2,392	-
1934	27,024	3,545	3,462	3,814	2,673	2,056	1,581	1,411	1,240	1,352	1,685	1,832	2,373	-
1933	25,397	2,935	3,291	3,306	2,531	1,949	1,675	1,360	1,234	1,330	1,652	1,817	2,314	-
1932	27,628	3,187	3,187	3,569	2,860	2,375	1,905	1,666	1,464	1,552	1,757	1,903	2,407	-
1931	30,364	3,746	3,919	3,813	2,959	2,470	2,047	1,830	1,577	1,568	1,853	2,078	2,504	-
1930	27,930	3,423	3,240	3,301	2,612	2,232	1,903	1,645	1,507	1,570	1,892	2,065	2,540	-
1929	31,875	4,076	3,810	4,207	3,249	2,615	2,012	1,881	1,566	1,771	1,910	2,191	2,587	-
1928	32,672	3,650	4,118	4,252	3,160	2,406	1,963	1,850	1,481	1,652	2,180	2,385	3,335	-
1927	33,855	4,269	4,432	4,432	3,218	2,753	2,218	1,923	1,734	1,858	2,071	2,218	2,836	-
1926	31,341	3,607	3,415	3,819	3,110	2,695	2,188	1,792	1,578	1,550	2,097	2,315	3,163	-
1925	35,520	4,397	5,554	5,088	3,388	2,639	2,141	1,924	1,559	1,588	2,148	2,182	2,911	-
1924	35,797	4,376	4,451	4,878	3,301	2,818	2,282	1,871	1,753	1,887	2,289	2,652	3,239	-
1923	36,188	4,484	4,334	4,340	3,339	2,611	2,281	2,108	1,856	2,110	2,643	2,580	3,512	-
1922	51,759	7,332	7,427	7,633	4,792	3,619	2,855	2,398	2,208	2,412	2,989	3,330	4,764	-
1921	52,049	5,898	5,841	6,352	4,860	3,919	3,253	2,792	2,603	2,892	3,533	4,492	5,573	-
1920	52,656	7,276	7,480	6,942	5,025	3,906	2,999	2,696	2,378	2,607	3,047	3,331	4,969	-

See footnotes at end of table.

TABLE 15. - 1/ DEATHS FROM SELECTED CAUSES BY MONTH: JAPAN, 1920 - 1949 - Cont'd
12/ PNEUMONIA, ALL FORMS (107-109)

Year	Total Deaths	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Un-known
*1949	56,636	7,529	6,816	7,277	6,378	4,918	3,452	2,828	2,025	2,251	2,692	4,043	6,427	-
1948	52,979	8,057	7,053	7,447	5,320	3,853	3,055	2,302	2,056	2,550	2,550	3,407	5,171	-
1947	101,601	15,412	18,996	16,152	11,633	8,574	6,060	4,141	2,864	2,606	3,147	4,490	7,521	5
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1943	116,494	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1942	106,022	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1941	104,073	13,506	16,194	15,724	11,435	9,321	6,686	4,964	3,929	3,632	4,363	5,624	8,695	-
1940	111,077	16,657	17,137	16,812	13,215	10,101	7,239	4,830	3,751	3,562	4,336	5,031	8,406	-
1939	131,542	16,152	17,466	22,217	16,196	11,495	9,111	5,943	4,586	4,025	5,047	6,584	12,120	-
1938	118,153	17,671	18,953	18,031	12,325	9,457	6,470	5,055	3,919	4,010	4,900	6,481	10,881	-
1937	108,256	13,108	13,476	14,500	12,718	10,104	7,829	5,764	4,328	4,278	5,194	6,072	10,884	1
1936	112,204	18,136	19,375	15,797	11,279	8,713	6,640	5,026	3,856	3,651	4,794	5,974	8,963	-
1935	105,078	13,482	13,489	14,244	11,636	9,596	7,418	5,224	4,306	4,293	4,757	6,273	10,360	-
1934	124,117	18,067	18,334	20,048	14,071	11,068	7,487	4,996	4,046	3,845	5,261	6,540	10,354	-
1933	106,247	12,924	14,029	16,222	11,994	9,473	7,107	4,759	3,915	3,867	5,229	6,421	10,106	1
1932	112,681	12,918	14,053	17,339	15,321	11,414	7,599	5,541	4,189	3,979	4,952	6,039	9,527	-
1931	129,380	16,488	19,043	20,340	14,853	12,117	9,286	6,095	4,742	4,141	5,233	6,851	10,190	1
1930	101,046	12,840	12,853	13,604	11,545	9,452	6,930	4,979	4,114	4,117	4,995	6,109	9,508	-
1929	121,181	14,701	14,920	19,064	15,611	11,893	8,233	5,961	4,653	4,530	5,227	6,740	9,648	-
1928	125,787	14,644	16,868	20,792	15,594	10,500	7,206	5,754	4,948	4,446	5,907	7,587	11,541	-
1927	119,960	14,754	15,775	17,066	12,862	11,988	8,765	6,176	4,995	4,827	5,510	6,872	10,399	1
1926	108,106	12,465	12,337	14,134	12,743	10,843	7,954	5,841	4,369	4,097	5,596	6,907	10,879	1
1925	129,129	16,428	22,559	21,195	13,604	10,787	8,350	5,927	4,343	4,220	5,471	6,637	9,608	-
1924	123,403	14,867	15,267	18,157	14,116	11,626	9,003	6,544	5,533	4,855	5,570	7,144	10,717	4
1923	119,263	14,658	14,556	15,988	12,363	10,501	8,419	6,992	5,630	5,630	6,552	7,305	10,669	-
1922	112,642	14,261	18,872	18,707	10,985	8,332	6,533	4,949	4,679	4,422	5,312	6,263	9,312	1
1921	113,709	11,824	12,441	15,398	13,044	10,218	8,117	6,397	5,449	5,523	6,476	8,109	10,713	-
1920	175,674	44,594	30,676	25,216	17,858	11,980	7,346	5,844	5,268	4,927	5,630	6,628	9,707	-

See footnotes at end of table.

TABLE 15. - 1/ DEATHS FROM SELECTED CAUSES BY MONTH: JAPAN, 1920 - 1949 - Cont'd
PLEURISY (110)

Year	Total Deaths	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Un-known
* 1949	6,030	515	503	612	548	537	517	540	506	461	469	422	400	-
**1948	7,700	746	690	751	720	713	662	608	627	601	524	520	538	-
1947	10,822	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1943	17,929	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1942	18,049	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1941	18,081	1,465	1,348	1,542	1,628	1,654	1,619	1,543	1,526	1,522	1,493	1,345	1,396	-
1940	20,099	1,787	1,732	1,887	1,860	1,813	1,668	1,785	1,615	1,562	1,504	1,387	1,459	-
1939	21,681	1,708	1,541	1,977	1,963	1,955	1,947	1,933	1,837	1,744	1,738	1,639	1,699	-
1938	20,980	1,693	1,595	1,805	1,931	1,856	1,828	1,819	1,832	1,654	1,720	1,620	1,623	-
1937	18,842	1,355	1,383	1,574	1,536	1,549	1,641	1,634	1,730	1,710	1,631	1,517	1,582	-
1936	18,036	1,518	1,392	1,586	1,648	1,656	1,557	1,572	1,562	1,407	1,415	1,321	1,402	-
1935	17,025	1,406	1,315	1,522	1,474	1,528	1,515	1,437	1,413	1,373	1,440	1,314	1,288	-
1934	17,811	1,467	1,383	1,677	1,632	1,675	1,573	1,532	1,378	1,415	1,407	1,321	1,351	-
1933	17,577	1,369	1,276	1,582	1,567	1,656	1,574	1,543	1,530	1,367	1,406	1,357	1,350	-
1932	17,560	1,458	1,417	1,590	1,606	1,477	1,700	1,503	1,439	1,396	1,327	1,277	1,277	-
1931	17,677	1,359	1,314	1,597	1,637	1,652	1,601	1,538	1,466	1,410	1,475	1,236	1,392	-
1930	16,581	1,436	1,263	1,561	1,490	1,481	1,386	1,452	1,388	1,396	1,337	1,122	1,269	-
1929	17,189	1,364	1,315	1,526	1,503	1,570	1,504	1,556	1,591	1,345	1,358	1,292	1,265	-
1928	16,144	1,239	1,212	1,439	1,425	1,376	1,350	1,421	1,398	1,364	1,380	1,249	1,291	-
1927	16,003	1,226	1,160	1,379	1,401	1,418	1,403	1,489	1,442	1,348	1,308	1,188	1,235	-
1926	15,179	1,151	1,161	1,390	1,347	1,417	1,295	1,295	1,345	1,239	1,302	1,128	1,115	-
1925	16,278	1,295	1,321	1,541	1,504	1,521	1,362	1,442	1,355	1,315	1,311	1,156	1,155	-
1924	16,755	1,370	1,341	1,501	1,482	1,462	1,449	1,470	1,484	1,418	1,290	1,259	1,229	-
1923	16,115	1,215	1,229	1,351	1,317	1,354	1,409	1,403	1,431	1,473	1,441	1,243	1,249	-
1922	16,111	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1921	14,903	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1920	14,008	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-

See footnote at end of table.

TABLE 15. - 1/ DEATHS FROM SELECTED CAUSES BY MONTH: JAPAN, 1920 - 1949 - Cont'd
13/ ULCER OF STOMACH OR DUODENUM (117)

Year	Total Deaths	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Un-known
* 1949	19,521	1,835	1,592	1,836	1,692	1,626	1,475	1,469	1,357	1,343	1,612	1,758	1,926	-
1948	20,705	2,295	1,980	1,984	1,759	1,606	1,554	1,434	1,348	1,445	1,664	1,774	1,862	-
1947	25,908	2,387	2,294	2,442	2,168	2,095	1,947	1,959	1,764	1,824	2,168	2,297	2,523	-
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1943	17,106	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1942	14,911	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1941	12,900	1,189	1,033	1,078	995	1,048	1,007	944	944	1,002	1,104	1,217	1,339	-
1940	12,923	1,342	1,186	1,161	1,052	993	994	908	900	927	1,100	1,134	1,232	-
1939	13,222	1,301	1,166	1,227	1,043	1,032	930	1,001	879	982	1,112	1,257	1,292	-
1938	13,279	1,284	1,101	1,091	1,027	1,024	970	968	922	1,062	1,207	1,299	1,254	-
1937	12,322	1,183	907	1,041	959	989	970	887	856	898	1,147	1,110	1,375	-
1936	11,977	1,207	1,067	1,080	987	918	804	899	840	844	1,094	1,146	1,091	-
1935	11,276	1,062	898	969	869	886	832	878	826	883	1,032	1,064	1,117	-
1934	11,329	1,060	959	1,065	898	824	814	846	776	849	1,037	1,115	1,086	-
1933	10,524	1,005	893	920	782	828	765	780	763	827	931	984	1,046	-
1932	10,056	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1931	9,701	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1930	9,190	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1929	8,689	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1928	8,610	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1927	8,030	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1926	7,837	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1925	7,769	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1924	7,391	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1923	7,142	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1922	6,386	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1921	6,704	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1920	6,245	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-

See footnotes at end of table.

TABLE 15. - 1/ DEATHS FROM SELECTED CAUSES BY MONTH: JAPAN, 1920 - 1949 - Cont'd
1A/ DIARRHEA, ENTERITIS AND ULCERATION OF THE INTESTINES (ALL AGES) (119-120)

Year	Total Deaths	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Un- known
1949	72,1	4,824	4,046	4,755	4,857	5,375	6,767	9,328	9,603	7,424	5,327	5,044	4,836	-
1948	83,3	5,416	5,429	6,133	5,746	6,016	7,172	11,681	11,479	8,034	5,628	5,512	5,017	1
1947	31,4	6,204	5,782	6,580	6,219	6,269	7,661	15,363	18,168	10,288	6,901	5,925	6,092	2
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1943	105,811	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1942	96,801	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1941	56,044	4,683	5,021	5,166	5,166	6,574	9,555	15,929	15,385	10,545	7,351	5,740	5,534	-
1940	108,367	5,119	5,697	5,837	5,643	6,779	9,353	17,980	19,071	11,979	8,079	6,043	5,687	-
1939	112,828	6,316	5,343	6,106	5,722	6,976	9,486	16,942	19,183	13,517	9,848	6,863	6,526	-
1938	116,956	6,841	5,846	5,063	5,077	7,377	10,660	18,033	18,995	13,966	9,969	7,105	6,224	-
1937	119,991	6,086	5,401	6,044	5,877	7,635	10,132	18,355	22,335	14,256	9,418	7,205	7,247	-
1936	121,133	6,745	6,167	6,693	6,456	7,527	9,993	18,687	20,604	14,622	10,247	6,920	6,471	1
1935	112,113	6,290	5,457	5,838	6,073	7,749	11,271	17,682	17,548	12,042	8,846	6,772	6,544	1
1934	127,853	6,879	5,858	6,855	6,751	8,193	12,378	21,739	21,005	14,129	10,292	7,326	6,518	-
1933	130,751	6,968	6,345	6,817	6,590	8,408	12,816	22,090	22,066	14,438	9,704	7,469	7,100	-
1932	137,351	7,470	6,840	7,123	6,876	8,522	13,097	21,533	23,695	16,295	10,473	7,693	7,824	-
1931	140,062	7,824	7,220	7,773	7,575	9,200	12,094	19,532	21,990	16,969	12,719	8,901	8,265	-
1930	142,383	8,270	7,138	7,705	7,636	9,662	13,085	21,895	24,017	15,626	10,795	8,707	8,047	-
1929	156,736	8,451	7,530	8,456	8,210	9,507	13,421	24,017	28,102	17,693	13,138	9,897	8,305	-
1928	146,359	7,532	7,515	7,552	7,272	9,196	12,573	21,244	22,767	18,263	13,465	9,605	8,975	-
1927	149,793	7,755	7,371	8,322	8,243	9,471	12,206	23,201	26,554	17,629	12,303	8,688	8,050	-
1926	141,200	8,015	7,037	8,015	7,945	9,588	13,342	18,898	23,005	16,859	11,824	8,425	8,246	1
1925	142,161	8,109	7,704	8,447	8,205	10,092	13,422	21,345	20,745	15,848	11,824	8,398	8,022	-
1924	151,718	9,111	8,172	8,969	8,210	10,099	12,378	21,648	24,777	17,905	13,104	9,442	8,499	3
1923	165,082	8,507	7,685	8,548	8,564	9,801	13,796	21,931	26,536	22,012	16,775	11,217	9,710	-
1922	153,321	7,995	7,528	8,619	8,097	10,377	14,170	22,277	25,741	17,868	12,292	9,353	8,308	1
1921	154,679	7,715	6,965	7,840	7,396	10,192	12,181	22,346	26,749	19,829	14,278	10,435	8,753	-
1920	142,091	7,826	7,561	7,779	7,138	8,305	11,508	22,236	24,007	17,202	11,782	8,560	8,187	-

See footnotes at end of table.

TABLE 15. - 1/ DEATHS FROM SELECTED CAUSES BY MONTH: JAPAN, 1920 - 1949 - Cont'd
15/ DIARRHEA, ENTERITIS AND ULCERATION OF THE INTESTINES, UNDER 2 YEARS OF AGE (119)

Year	Total Deaths	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Un-known
*1949	41,822	2,825	2,295	2,696	2,824	3,263	4,424	5,676	5,371	3,734	2,631	3,069	3,014	-
1948	48,444	2,996	2,799	3,024	2,895	3,308	4,587	2,943	7,108	4,439	2,931	3,369	3,045	-
1947	55,485	2,609	2,400	2,983	2,943	2,981	4,300	9,596	11,850	5,627	3,348	3,279	3,569	-
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1943	53,057	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1942	48,657	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1941	47,098	2,450	2,145	2,264	2,421	3,477	5,579	8,624	7,320	4,544	3,000	2,618	2,656	-
1940	49,212	2,768	2,490	2,523	2,405	2,984	4,616	9,131	8,662	5,107	3,192	2,680	2,654	-
1939	52,965	3,074	2,501	2,852	2,605	3,562	5,069	9,145	9,150	5,524	3,707	2,705	3,071	-
1938	58,465	3,647	2,926	2,995	2,949	3,962	6,112	9,475	9,392	6,426	4,296	3,191	3,094	-
1937	66,698	3,394	2,955	3,198	3,159	4,457	6,401	11,294	12,881	7,275	4,172	3,563	3,949	-
1936	70,539	3,850	3,359	3,672	3,660	4,448	6,433	12,186	12,452	7,988	5,227	3,625	3,638	1
1935	64,298	3,395	2,942	3,157	3,441	4,639	7,446	11,410	10,165	6,187	4,159	3,589	3,767	1
1934	74,812	3,917	3,218	3,745	3,774	4,843	8,380	14,541	12,442	7,456	5,124	3,730	3,642	-
1933	79,914	4,042	3,617	3,893	3,726	5,213	8,767	15,337	14,063	8,018	4,900	4,080	4,258	-
1932	82,422	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1931	83,482	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1930	85,652	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1929	93,500	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1928	86,978	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1927	91,087	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1926	87,280	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1925	84,857	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1924	87,597	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1923	93,172	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1922	87,273	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1921	87,991	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1920	79,325	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-

See footnotes at end of table.

TABLE 15. - 1/ DEATHS FROM SELECTED CAUSES BY MONTH: JAPAN, 1920 - 1949 - Cont'd
16/ DIARRHEA, ENTERITIS, AND ULCERATION OF THE INTESTINES, 2 YEARS OF AGE AND OVER (120)

Year	Total Deaths	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Un-known
1949	30,364	1,999	1,751	2,059	2,033	2,112	2,343	3,652	4,232	3,690	2,666	1,975	1,822	-
1948	34,820	2,420	2,630	3,109	2,851	2,708	2,585	3,738	4,371	3,595	2,697	2,143	1,972	1
1947	45,969	3,595	3,392	3,597	3,276	3,288	3,361	5,767	6,318	4,661	3,553	2,646	2,523	2
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1943	52,754	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1942	48,144	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1941	49,626	2,791	2,538	2,757	2,745	3,097	3,976	7,305	8,065	6,001	4,351	3,122	2,878	-
1940	59,155	3,451	3,207	3,314	3,238	3,795	4,737	8,849	10,409	6,872	4,887	3,363	3,033	-
1939	59,863	3,242	2,842	3,254	3,117	3,414	4,417	7,797	10,033	7,953	6,141	4,158	3,455	-
1938	58,491	3,194	2,920	2,968	3,028	3,415	4,548	8,558	9,603	7,540	5,673	3,914	3,130	-
1937	53,293	2,692	2,446	2,846	2,718	3,178	3,731	7,061	9,454	6,981	5,246	3,642	3,298	-
1936	50,594	2,895	2,808	3,021	2,796	3,079	3,560	6,501	8,152	6,634	5,020	3,295	2,833	-
1935	47,815	2,895	2,515	2,681	2,632	3,110	3,825	6,272	7,383	5,855	4,687	3,183	2,777	-
1934	53,041	2,962	2,640	3,110	2,977	3,280	3,998	7,198	8,563	6,673	5,168	3,596	2,876	-
1933	50,837	2,926	2,728	2,924	2,664	3,195	4,049	6,753	7,943	6,420	4,804	3,389	2,842	-
1932	54,929	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1931	56,580	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1930	56,931	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1929	63,236	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1928	59,381	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1927	58,706	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1926	53,920	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1925	57,304	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1924	64,121	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1923	71,910	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1922	66,048	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1921	66,682	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1920	62,766	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	

See footnotes at end of table.

TABLE 15. - 1/ DEATHS FROM SELECTED CAUSES BY MONTH: JAPAN, 1920 - 1949 - Cont'd
APPENDICITIS (121)

Year	Total Deaths	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Un- known
*1949	3,267	266	231	288	270	290	255	334	348	305	238	227	215	-
1948	3,808	276	338	331	342	293	346	342	393	356	290	256	230	-
1947	4,424	300	262	288	328	354	391	507	528	429	406	302	329	-
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1943	2,291	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1942	2,360	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1941	2,141	170	176	177	171	188	184	204	232	192	187	164	192	-
1940	2,577	173	195	218	228	198	227	239	281	222	209	195	192	-
1939	2,701	226	170	208	215	211	232	264	265	268	238	197	207	-
1938	2,672	203	176	223	199	221	199	249	286	234	271	219	192	-
1937	2,695	229	168	206	191	225	226	244	284	275	233	194	220	-
1936	2,553	222	164	205	207	187	202	242	267	244	235	208	170	-
1935	2,440	170	171	172	185	187	208	217	244	230	237	204	215	-
1934	2,609	205	167	206	204	195	215	238	293	266	223	210	187	-
1933	2,532	202	157	192	179	203	205	245	295	281	211	177	185	-
1932	2,648	208	179	197	187	206	214	241	296	251	245	219	205	-
1931	2,530	171	192	194	184	202	236	235	255	250	247	193	171	-
1930	2,598	205	188	213	192	210	200	252	282	266	235	180	175	-
1929	2,661	172	166	212	209	216	196	262	311	266	223	192	200	-
1928	2,586	187	152	201	183	199	208	254	268	266	227	210	231	-
1927	2,646	186	158	204	211	225	207	257	266	275	263	190	204	-
1926	2,559	216	162	193	194	178	207	241	274	272	251	202	159	-
1925	2,626	214	186	197	194	198	206	252	261	246	281	188	203	-
1924	2,692	203	177	193	178	186	209	249	297	277	267	250	206	-
1923	2,763	236	184	177	197	182	191	222	291	294	330	246	213	-
1922	2,817	210	135	175	170	178	220	277	345	340	300	243	224	-
1921	2,879	234	162	175	174	202	245	302	319	333	289	235	209	-
1920	2,624	146	143	180	168	190	209	236	326	279	293	254	200	-

See footnotes at end of table.

TABLE 15. - 1/ DEATHS FROM DETECTED CAUSES BY MONTH: JAPAN, 1920 - 1949 - Cont'd
17/ NEPHRITIS (ACUTE, CHRONIC AND UNSPECIFIED) (130-132)

Year	Total Deaths	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Un- known
+ 1949	33,859	3,265	2,910	3,330	2,897	2,587	2,421	2,514	2,473	2,591	2,896	2,926	3,069	-
1948	36,561	3,806	3,427	3,312	2,919	3,005	2,766	2,739	2,564	2,820	3,059	3,039	3,105	-
1947	45,017	4,439	4,351	4,371	3,920	3,644	3,324	3,273	3,297	3,280	3,811	3,561	3,745	1
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1943	49,708	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1942	51,108	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1941	48,900	5,009	4,581	4,277	3,931	3,655	3,436	3,605	3,752	4,111	4,447	4,441	4,655	-
1940	54,703	5,714	5,396	5,107	4,384	4,220	3,842	3,859	4,017	4,181	4,734	4,589	4,670	-
1939	58,799	6,204	5,177	5,595	4,574	4,313	4,034	4,132	4,326	4,513	5,171	5,085	5,675	-
1938	61,996	6,413	5,542	5,341	4,666	4,407	4,096	4,043	4,043	5,369	5,447	5,529	5,817	-
1937	56,285	5,219	4,364	4,732	4,288	4,080	3,978	3,789	4,426	4,820	5,513	5,250	5,827	-
1936	57,771	6,243	5,433	5,353	4,405	4,207	3,989	4,041	4,345	4,454	5,273	5,081	5,007	-
1935	56,026	5,358	4,784	4,783	4,277	4,119	3,851	4,010	4,320	4,676	5,317	5,095	5,436	-
1934	59,553	6,154	5,248	5,547	4,628	4,348	4,165	4,243	4,226	4,699	5,677	5,358	5,260	-
1933	58,986	5,663	5,291	5,532	4,578	4,322	4,169	4,238	4,484	5,156	5,734	5,412	5,407	-
1932	61,360	5,654	5,507	5,747	5,034	4,616	4,284	4,408	4,684	5,026	5,589	5,362	5,656	1
1931	64,241	6,162	5,666	5,666	5,015	4,799	4,677	4,675	4,722	5,428	6,081	5,722	5,831	-
1930	63,435	5,889	5,120	5,542	4,891	5,003	4,646	4,665	5,011	5,533	5,869	5,651	5,615	-
1929	65,391	6,039	5,526	5,876	5,065	4,903	4,655	4,975	5,299	5,559	5,981	5,945	5,608	-
1928	62,884	5,650	5,144	5,338	4,602	4,394	4,119	4,558	5,026	5,651	6,565	5,972	5,865	-
1927	60,034	5,454	5,061	5,492	4,455	4,524	4,121	4,126	4,804	5,361	5,815	5,396	5,335	-
1926	59,581	5,440	4,785	5,284	4,829	4,606	4,314	4,301	4,588	4,952	5,719	5,410	5,341	2
1925	59,873	5,770	5,481	5,380	4,574	4,476	4,312	4,532	4,428	4,728	5,653	5,119	5,419	1
1924	62,227	5,588	5,129	5,455	4,397	4,564	4,377	4,593	5,610	5,553	5,881	5,588	5,492	-
1923	62,574	5,560	4,716	4,956	4,367	4,261	4,405	4,805	5,142	6,198	6,915	5,784	5,517	8
1922	60,870	5,791	4,941	5,056	4,198	4,119	3,960	4,361	5,115	5,809	6,089	5,813	5,618	-
1921	59,788	4,806	4,512	4,665	3,908	3,945	3,915	4,495	5,188	6,221	6,678	6,126	5,609	-
1920	55,546	4,891	4,405	4,358	3,786	3,641	3,632	4,061	5,010	5,567	5,685	5,184	5,325	-

See footnotes at end of table.

TABLE 15. - $\frac{1}{2}$ DEATHS FROM SELECTED CAUSES BY MONTH: JAPAN, 1920 - 1949 - Cont'd
12/ DISEASE OF PREGNANCY, CHILD BIRTH AND THE PUERPERIUM (140-150)

Year	Total Deaths	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Un-known
* 1949	4,495	404	346	406	413	373	342	345	415	361	369	354	367	-
1948	4,437	425	425	411	354	331	318	332	395	391	363	312	360	-
1947	4,482	413	386	343	346	353	338	406	391	394	351	376	390	1
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1943	4,542	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1942	4,586	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1941	4,929	489	436	456	366	324	335	405	401	438	429	385	466	-
1940	5,070	516	533	459	381	332	340	435	401	418	372	410	473	-
1939	4,818	500	413	412	338	346	293	380	418	423	416	414	465	-
1938	4,877	508	477	453	353	336	318	385	476	429	376	384	382	-
1937	5,444	498	492	442	443	380	349	415	487	487	465	467	519	-
1936	5,384	563	482	505	422	348	392	442	428	471	449	437	445	-
1935	5,498	551	505	495	439	391	374	437	497	478	537	488	506	-
1934	5,709	608	503	520	435	437	419	422	488	471	468	450	488	-
1933	5,763	557	473	578	452	405	391	439	479	490	506	529	464	-
1932	5,530	488	502	449	474	401	377	442	503	452	496	446	460	-
1931	5,667	529	452	511	427	457	434	449	514	487	521	458	428	-
1930	5,681	507	459	502	398	447	391	493	530	505	478	461	510	-
1929	5,867	518	477	555	474	446	404	539	618	502	478	427	429	-
1928	5,997	504	502	524	457	440	449	469	534	560	561	504	493	-
1927	5,765	518	502	550	478	438	420	427	514	498	544	430	446	-
1926	5,721	494	471	520	434	440	429	485	501	510	514	464	458	1
1925	6,309	559	591	599	486	477	478	523	545	523	556	481	491	-
1924	6,273	525	507	530	472	454	404	540	627	630	575	502	507	-
1923	6,897	589	570	569	497	430	479	646	695	717	646	548	511	-
1922	6,565	623	569	563	482	461	414	549	603	617	640	510	534	-
1921	7,181	653	573	579	491	516	466	619	714	708	702	599	561	-
1920	7,158	715	654	612	539	497	468	588	614	679	633	556	603	-

See footnotes at end of table.

TABLE 15. - 1/ DEATHS FROM SELECTED CAUSES BY MONTH: JAPAN, 1920 - 1949 - Cont'd
CONGINITAL MAIFORMATION (157)

Year	Total Deaths	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Un-known
* 1949	6,556	641	573	678	560	493	463	459	411	495	536	605	642	-
1948	5,751	546	532	580	448	388	373	448	427	458	460	505	586	-
1947	4,990	419	396	430	361	369	351	418	411	363	427	517	528	-
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1943	3,781	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1942	3,897	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1941	3,670	344	353	340	316	280	266	278	298	249	318	307	321	-
1940	3,507	364	376	331	300	278	228	270	216	240	265	294	345	-
1939	3,718	374	351	383	291	254	225	261	282	271	294	303	429	-
1938	3,916	397	400	380	326	312	249	304	292	291	290	320	355	-
1937	4,295	431	380	420	362	332	301	318	304	321	336	358	432	-
1936	4,308	462	402	432	357	295	282	335	334	334	343	336	396	-
1935	4,333	422	384	395	367	339	289	287	318	347	359	374	452	-
1934	4,198	421	348	444	361	325	278	314	307	293	353	348	386	-
1933	4,381	470	431	424	328	339	305	305	314	349	356	349	411	-
1932	2,785	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1931	2,823	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1930	2,815	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1929	3,018	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1928	3,070	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1927	3,002	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1926	3,353	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1925	3,075	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1924	2,797	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1923	2,177	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1922	1,284	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1921	1,306	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1920	1,251	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	

See footnotes at end of table.

TABLE 15. - 1/ DEATHS FROM UNIDENTIFIED CAUSES BY MONTH: JAPAN, 1920 - 1945 - Cont'd
CONGENITAL DEBILITY (158)

Year	Total Deaths	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Unknown
* 1949	37,285	4,885	4,024	4,664	3,397	2,524	2,224	2,385	2,174	2,000	2,298	3,009	3,701	-
1948	38,204	5,471	5,176	4,405	2,944	2,361	2,145	2,186	2,111	2,153	2,319	2,820	3,913	-
1947	41,751	4,811	5,390	5,013	3,346	2,509	2,188	2,653	2,325	2,169	2,673	3,595	5,079	-
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1943	51,941	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1942	52,096	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1941	52,376	6,192	6,265	5,416	4,137	3,337	3,278	3,653	3,517	3,426	3,794	3,999	5,362	-
1940	54,939	7,321	6,923	6,407	4,396	3,585	3,194	3,772	3,509	3,262	3,557	3,894	5,119	-
1939	57,124	7,973	6,633	6,207	4,213	3,484	3,032	3,528	3,529	3,431	3,987	4,615	6,492	-
1938	60,568	8,669	7,952	6,496	4,158	3,627	3,345	3,720	3,596	3,578	3,984	4,792	6,151	-
1937	63,387	7,340	6,358	6,393	4,803	3,940	3,865	4,151	4,169	4,239	4,946	5,467	7,676	-
1936	67,986	9,534	8,578	7,992	5,440	4,200	3,995	4,401	4,173	3,912	4,573	5,073	6,115	-
1935	64,369	7,732	7,128	6,543	4,749	3,997	3,675	4,109	4,143	4,300	4,880	5,658	7,455	-
1934	68,604	8,831	8,012	7,568	5,615	4,441	3,989	4,323	4,221	4,181	5,055	5,451	6,517	-
1933	67,498	8,798	7,916	7,921	5,340	4,244	4,021	4,276	4,139	3,965	4,818	5,361	6,799	-
1932	59,389	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1931	61,448	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1930	59,288	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1929	65,424	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1928	63,901	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1927	64,132	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1926	67,227	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1925	67,019	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1924	69,551	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1923	73,748	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1922	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1921	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1920	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-

See footnotes at end of table.

TABLE 15. - 1/ DEATHS FROM SELECTED CAUSES BY MONTH: TATAN, 1920 - 1949 - Cont'd
PREMATURE BIRTH (159)

Year	Total Deaths	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Un- known
*1949	13,667	1,484	1,272	1,362	1,112	1,015	972	1,003	1,002	989	963	1,117	1,376	-
1948	13,720	1,602	1,461	1,311	987	952	935	914	943	940	1,075	1,215	1,385	-
1947	14,597	1,696	1,387	1,363	1,075	979	955	1,078	1,077	905	1,019	1,337	1,576	-
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1943	5,084	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1942	5,152	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1941	5,284	589	576	459	346	361	398	370	390	401	448	433	513	-
1940	5,267	672	570	497	398	387	325	364	382	357	388	429	498	-
1939	5,165	624	524	465	328	344	288	355	381	346	453	462	595	-
1938	5,410	732	626	522	397	349	341	353	337	357	381	459	556	-
1937	5,929	653	548	510	403	423	389	422	476	412	509	563	621	-
1936	6,328	765	725	634	385	365	415	430	404	402	502	547	680	-
1935	5,685	648	550	430	377	342	397	398	418	478	484	513	650	-
1934	5,654	667	590	544	437	329	332	376	385	397	470	501	626	-
1933	5,071	643	498	470	360	335	324	340	354	369	414	416	548	-
1932	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1931	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1930	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1929	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1928	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1927	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1926	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1925	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1924	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1923	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1922	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1921	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-
1920	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	-

See footnotes at end of table.

TABLE 15. - 1/ DEATHS FROM SELECTED CAUSES BY MONTH: JAPAN, 1920 - 1949 - Continued
INJURY AT BIRTH (160)

Year	Total Deaths	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Un- known
*1949	1,082	89	91	92	96	75	85	95	90	91	101	85	92	-
** 1948	1,030	108	111	99	79	82	72	83	91	77	55	69	104	-
1947	885	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1943	447	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1942	459	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1941	443	35	55	41	40	28	24	30	43	35	33	35	44	-
1940	291	32	18	41	22	19	24	28	23	24	18	18	24	-
1939	314	20	31	17	27	24	23	32	23	19	28	26	44	-
1938	381	41	39	35	31	29	27	35	35	29	27	26	27	-
1937	441	35	38	43	36	42	36	39	43	27	36	30	36	-
1936	407	34	40	45	34	17	19	36	41	43	33	28	37	-
1935	425	40	37	48	35	28	29	39	33	32	29	40	35	-
1934	401	44	30	51	27	27	29	30	25	37	34	37	30	-
1933	361	31	34	38	22	18	23	42	37	26	23	27	40	-
1932	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1931	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1930	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1929	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1928	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1927	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1926	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1925	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1924	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1923	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1922	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1921	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1920	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	

See footnotes at end of table.

TABLE 15. - $\frac{1}{2}$ DEATHS FROM SELECTED CAUSES BY MONTH: JAPAN, 1920 - 1949 - Cont'd
15/ SENILITY (162)

Year	Total Deaths	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Un-known
*1949	66,699	6,635	5,893	6,815	5,927	4,749	4,268	4,598	4,784	4,861	5,545	5,853	6,811	-
1948	64,074	8,127	7,206	7,139	5,644	4,286	4,019	4,125	4,308	4,168	4,662	4,859	5,528	3
1947	78,953	9,277	9,784	8,937	6,613	5,596	4,902	4,896	5,098	4,619	5,404	6,064	7,755	8
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1944	N	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1943	99,636	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1942	96,428	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1941	90,215	10,134	9,625	8,299	6,755	5,915	5,615	6,660	6,868	7,149	7,625	7,150	8,460	-
1940	90,098	9,988	9,688	8,762	6,898	5,982	5,400	6,384	7,550	7,129	7,124	6,880	8,313	-
1939	95,466	10,419	9,155	9,886	7,177	6,092	5,849	6,626	7,789	7,749	7,942	7,542	9,240	-
1938	98,772	11,242	10,635	9,099	6,892	5,914	5,651	7,284	8,522	8,835	8,375	7,902	8,461	-
1937	84,766	7,837	7,133	7,098	6,129	5,469	5,258	5,854	7,648	7,634	8,084	7,231	9,381	-
1936	91,936	11,687	11,560	9,524	6,788	5,688	5,278	6,079	6,855	6,931	7,549	6,931	7,066	-
1935	79,240	8,099	7,237	6,898	5,938	5,118	5,035	5,732	6,978	6,679	7,007	6,608	7,910	1
1934	87,323	10,443	9,295	9,201	6,631	5,669	5,151	5,570	6,533	6,806	7,595	7,256	7,173	-
1933	83,197	8,448	8,367	8,267	6,177	5,444	5,069	5,787	7,111	6,959	7,266	6,805	7,497	-
1932	77,529	7,320	7,289	7,666	6,233	5,259	4,679	5,559	6,526	6,567	6,999	6,379	7,053	-
1931	85,650	9,169	9,115	8,975	7,071	6,012	5,431	5,747	6,329	6,724	7,407	6,582	7,088	-
1930	76,591	7,794	6,828	6,593	5,583	5,538	5,087	5,731	6,625	6,767	6,650	6,277	7,118	-
1929	79,181	8,167	7,640	7,983	6,172	5,518	4,850	5,880	7,012	6,902	6,778	6,301	5,978	-
1928	77,251	7,451	7,105	7,469	6,071	4,780	4,444	5,218	6,370	6,985	7,664	6,562	7,131	1
1927	74,664	7,225	7,648	7,713	5,663	5,281	4,659	5,085	6,425	6,761	6,298	5,768	6,138	-
1926	76,576	6,772	5,819	6,097	5,161	4,546	4,171	4,493	5,563	5,855	6,541	5,958	6,596	4
1925	70,065	7,242	6,612	8,033	5,332	4,635	4,072	4,740	5,111	5,419	5,756	5,756	5,854	2
1924	74,003	7,459	7,133	7,714	5,523	4,987	4,408	4,996	6,716	6,247	6,546	6,054	6,209	1
1923	76,579	7,083	6,354	6,596	5,323	4,682	4,681	5,067	7,067	7,852	8,056	6,516	6,561	3
1922	75,957	8,382	7,382	7,894	5,668	4,751	4,237	5,032	6,343	6,868	6,565	5,860	6,553	1
1921	75,654	6,798	6,425	6,526	5,126	4,446	4,456	5,258	6,982	7,966	7,645	7,227	6,809	-
1920	73,468	7,754	7,628	7,101	5,441	4,731	4,215	4,928	6,536	6,989	6,321	5,432	6,392	-

See footnotes at end of table.

TABLE 15. - 1/ DEATHS FROM SELECTED CAUSES BY MONTH: JAPAN, 1920 - 1949 - Cont'd
SUICIDE (163-164)

Year	Total Deaths	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Un-known
* 1949	14,211	864	860	1,118	1,342	1,593	1,250	1,396	1,288	1,169	1,140	1,091	1,100	-
1948	12,753	805	829	1,112	1,240	1,301	1,255	1,339	1,198	1,073	944	818	827	12
1947	12,262	698	625	981	1,208	1,431	1,219	1,412	1,175	1,004	911	824	765	9
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	8,784	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	9,393	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1941	9,713	674	655	850	928	980	913	964	912	717	771	705	644	-
1940	9,877	633	635	825	917	1,096	885	1,036	885	800	772	715	678	-
1939	10,785	746	722	937	1,058	1,189	987	1,064	979	860	812	749	660	22
1938	12,223	799	818	1,134	1,267	1,385	1,167	1,159	1,179	944	881	745	738	7
1937	14,295	1,035	1,019	1,268	1,512	1,531	1,327	1,483	1,310	1,086	987	904	804	9
1936	15,423	975	1,009	1,300	1,527	1,713	1,515	1,512	1,385	1,286	1,157	1,062	996	6
1935	14,172	970	987	1,206	1,385	1,514	1,288	1,379	1,233	1,072	1,100	1,052	977	9
1934	14,554	1,054	938	1,222	1,452	1,539	1,379	1,415	1,348	1,092	1,043	1,000	1,068	4
1933	14,805	1,039	997	1,240	1,518	1,667	1,422	1,545	1,313	1,145	1,002	1,002	908	7
1932	14,746	1,072	909	1,222	1,343	1,578	1,426	1,608	1,445	1,194	1,075	945	926	3
1931	14,353	923	893	1,259	1,362	1,490	1,452	1,361	1,376	1,081	1,063	999	1,088	6
1930	13,942	906	879	1,171	1,324	1,418	1,316	1,516	1,321	1,170	982	966	963	4
1929	12,740	842	758	1,077	1,232	1,320	1,191	1,339	1,158	985	1,000	886	948	4
1928	13,032	892	858	1,128	1,332	1,407	1,177	1,366	1,161	1,071	977	843	819	1
1927	12,845	825	773	999	1,269	1,313	1,250	1,381	1,179	1,013	1,033	885	909	6
1926	12,484	859	770	1,030	1,148	1,395	1,174	1,357	1,212	993	903	846	783	74
1925	12,249	747	741	966	1,194	1,378	1,287	1,321	1,064	985	941	842	763	20
1924	11,261	725	707	927	1,082	1,221	1,057	1,218	1,082	875	862	770	723	12
1923	11,488	730	708	1,037	1,086	1,169	1,170	1,243	1,193	944	797	716	679	16
1922	11,546	796	859	984	1,129	1,221	1,128	1,218	1,101	872	860	704	702	10
1921	11,358	781	885	710	1,080	1,194	1,157	1,157	1,157	900	907	772	788	8
1920	10,630	658	653	775	975	1,089	1,050	1,186	1,110	928	800	715	679	12

See footnotes at end of table.

TABLE 15. - 1/ DEATHS FROM SELECTED CAUSES BY MONTH: JAPAN, 1920 - 1949 - Cont'd
HOMICIDE (165-168)

Year	Total Deaths	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Un- Known
* 1949	1,669	109	119	145	132	150	115	144	171	164	135	143	142	-
1948	1,641	120	118	153	185	159	121	161	150	114	118	115	127	-
1947	1,314	65	77	105	126	128	131	124	121	108	113	101	114	1
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1943	226	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1942	282	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1941	238	16	15	17	21	15	21	18	30	25	18	20	22	-
1940	259	11	22	20	22	22	21	31	28	29	17	18	18	-
1939	253	21	16	30	12	33	22	23	22	11	22	19	22	-
1938	435	29	32	53	41	78	44	29	20	27	28	24	30	-
1937	450	41	40	33	42	58	44	43	31	31	31	33	25	-
1936	468	24	42	52	57	47	30	38	47	40	34	28	29	-
1935	427	34	29	33	27	29	55	50	28	27	39	40	36	-
1934	423	36	22	33	38	46	47	42	38	37	35	27	22	-
1933	461	39	32	44	32	38	51	41	41	36	47	34	26	-
1932	451	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1931	491	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1930	395	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1929	395	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1928	422	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1927	444	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1926	457	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1925	429	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1924	419	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1923	371	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1922	441	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1921	330	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1920	365	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	

See footnotes at end of table.

TABLE 15. - $\frac{1}{2}$ DEATHS FROM SELECTED CAUSES BY MONTH: JAPAN, 1920 - 1949 - Cont'd
ACCIDENTAL DEATHS (189-195)

Year	Total Deaths	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Un-known
* 1949	34,498	2,253	1,984	2,720	2,537	2,871	3,125	4,320	4,463	3,166	2,528	2,240	2,491	-
1948	38,960	2,700	2,262	2,574	2,501	2,714	6,699	4,458	4,183	3,717	2,553	2,177	2,463	9
1947	38,471	2,640	2,592	2,643	2,500	2,957	2,811	5,106	5,024	4,245	2,619	2,349	2,573	12
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1943	33,476	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1942	31,087	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1941	28,742	2,185	1,843	2,208	2,230	2,285	2,491	3,178	3,618	2,339	2,420	1,871	2,074	-
1940	28,328	2,541	1,990	2,073	2,182	2,284	2,408	3,547	3,041	2,336	2,014	1,876	2,036	-
1939	29,262	2,529	2,061	2,223	2,262	2,382	2,373	3,826	3,161	2,329	2,163	1,916	2,032	5
1938	31,637	2,443	2,098	2,269	2,274	2,382	2,764	4,105	3,470	2,855	2,743	2,044	2,185	5
1937	30,095	1,957	1,997	2,218	2,292	2,274	2,433	3,618	3,952	2,441	2,253	2,190	2,425	5
1936	30,080	2,390	2,218	2,269	2,521	2,310	2,573	3,622	3,148	2,519	2,275	2,203	2,029	3
1935	28,874	2,000	1,817	2,266	2,195	2,477	2,573	3,607	3,180	2,618	2,326	1,912	2,031	12
1934	31,830	2,162	1,922	3,934	2,141	2,222	2,424	3,196	2,890	4,885	2,083	1,933	2,033	5
1933	29,953	1,971	1,866	4,676	2,030	2,253	2,461	3,686	3,134	2,212	2,109	1,754	1,795	6
1932	26,637	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1931	25,165	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1930	26,176	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1929	26,529	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1928	25,939	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1927	29,215	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1926	25,526	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1925	24,893	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1924	25,699	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1923	71,298	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1922	25,436	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1921	24,438	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1920	26,056	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	

See footnotes at end of table.

TABLE 15. - 1/ DEATHS FROM SELECTED CAUSES BY MONTH: JAPAN, 1920 - 1949 - Cont'd
20/ ILL-DEFINED AND UNKNOWN (200a and 200c)

Year	Total Deaths	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Un-known
*1949	16,900	1,652	1,503	1,716	1,595	1,333	1,281	1,298	1,236	1,115	1,248	1,358	1,565	-
**1948	17,188	1,835	1,752	1,840	1,496	1,407	1,313	1,277	1,312	1,149	1,222	1,210	1,375	-
**1947	20,147	1,984	2,082	2,174	1,834	1,574	1,447	1,571	1,592	1,281	1,361	1,421	1,826	-
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1943	31,638	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1942	31,738	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1941	30,931	3,599	3,608	3,150	2,605	2,421	2,097	2,094	2,005	1,977	2,235	2,316	2,824	-
1940	33,008	4,066	3,869	3,538	2,953	2,636	2,293	2,379	2,148	2,031	2,131	2,204	2,720	-
1939	35,180	4,155	3,740	4,051	3,146	2,694	2,364	2,286	2,280	2,210	2,382	2,481	3,388	3
1938	36,255	4,636	4,472	3,872	3,113	2,604	2,278	2,295	2,356	2,448	2,364	2,626	3,189	2
1937	32,819	3,480	2,991	3,254	2,753	2,593	2,310	2,193	2,208	2,203	2,528	2,542	3,763	1
1936	37,838	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1935	33,600	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1934	37,339	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1933	37,955	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1932	40,992	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1931	44,611	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1930	41,137	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1929	45,486	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1928	45,485	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1927	42,958	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1926	39,115	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1925	40,787	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1924	43,243	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
1923	45,867	8,076	7,423	8,056	5,627	5,019	4,621	5,090	5,448	5,529	5,673	5,551	6,490	8
1922	72,611	6,403	6,462	6,781	5,367	5,144	4,765	5,362	6,130	6,090	6,484	6,470	6,783	9
1921	72,450	8,426	8,389	7,550	6,084	5,157	4,623	5,116	5,496	5,544	5,670	5,256	6,525	13
1920	73,899	8,426	8,389	7,550	6,084	5,157	4,623	5,116	5,496	5,544	5,670	5,256	6,525	13

See footnotes at end of table.

FOOTNOTES:

- * Data for 1949 are provisional.
- ** The 1947 and 1948 monthly death data for syphilis (30) and ill-defined and unknown cause (200a and c) and the 1948 monthly death data for simple meningitis (81a), pleurisy (110), and injury at birth (160) are estimates based on preliminary figures. The annual totals are final.
- 1/ 1920-1941: All data for 1920-1941 refer to deaths of Japanese nationals in Japan Proper and Okinawa, including military personnel in the country.
- 1942-1943: The annual totals for 1942 refer to deaths of Japanese nationals in Japan Proper and Okinawa, including military personnel in the country. The annual totals for 1943 refer to deaths of Japanese nationals in Japan Proper, Okinawa and Karafuto, including military personnel in the country. However, the monthly data for 1942-1943 refer to deaths of Japanese nationals and foreigners in Japan Proper and are limited to the civilian population.
- 1944-1946: All data refer to deaths among Japanese nationals and foreigners in Japan Proper and are limited to the civilian population only.
- 1947-1949: All data refer to deaths of Japanese nationals in Japan Proper,
- 1947-1949: All data for 1920-1941, 1947-1949, and annual totals for 1942 and 1943 are based upon tabulations of death certificates.
- All data for 1944-1946 (except tuberculosis) and monthly data 1942 and 1943 are based upon follow-up reports on communicable disease cases. Data for tuberculosis deaths 1944-1946 are estimates prepared by the Ministry of Welfare.
- 2/ Typhoid fever. 1920-1943: includes typhoid pneumonia. 1944-1946, unknown whether typhoid pneumonia is included. 1947-1949 excludes typhoid pneumonia.
- 3/ Scarlet fever. 1920-1943: includes puerperal scarlet fever. 1944-1946: unknown whether puerperal scarlet fever is included. 1947-1949: excludes puerperal scarlet fever.
- 4/ Diphtheria. 1920-1922 and 1933-1943: excludes diphtheric endocarditis; includes croupous bronchitis and laryngeal group. 1923-1932: excludes diphtheric endocarditis and croupous bronchitis; includes laryngeal group. 1944-1946: unknown whether diphtheric endocarditis, croupous bronchitis, and laryngeal group are included. 1947-1949: includes diphtheric endocarditis; excludes croupous bronchitis and laryngeal group.
- 5/ Tuberculosis (all forms). 1920-1943: includes chronic pleuro pneumonia and all adenopathy; excludes spondylitis. 1947-1949: excludes chronic pleuro pneumonia and nontubercular adenopathy; includes spondylitis. 1944-1946: are estimates of the Ministry of Welfare.
- 6/ Syphilis. 1920-1922: excludes general paralysis of the insane and aneurysm of the aorta. 1923-1932: includes general paralysis of the insane, a small amount of non-syphilitic paralytic psychosis, aneurysm of the aorta and a small amount of non-syphilitic aneurysm. 1933-1943: includes general paralysis of the insane, aneurysm of the aorta and a small amount of non-syphilitic aneurysm. 1947-1949: includes general paralysis of the insane and aneurysm of the aorta.
- 7/ Typhus fever and typhus-like diseases. 1920-1943 and 1947-1949: includes typhus-like diseases. 1944-1946: excludes typhus-like diseases.
- 8/ Simple meningitis. 1920-1943: includes acute cerebrospinal meningitis not due to meningococcus. 1947-1949: excludes acute cerebrospinal meningitis not due to meningococcus.

TABLE 15. - 2/ DEATHS FROM SELECTED CAUSES BY MONTH: JAPAN, 1920 - 1949 - Cont'd

FOOTNOTES - Cont'd:

- 2/ Intracranial lesions of vascular origin. 1920-1932: includes alcoholic cerebral atrophy, congestion and a small amount of paralysis not due to intracranial lesions of vascular origin. 1933-1943: includes alcoholic cerebral atrophy and congestion. 1947-1949: excludes alcoholic atrophy and congestion.
- 10/ Diseases of the heart. 1920-1943: includes diphtheric endocarditis. 1947-1949: excludes diphtheric endocarditis.
- 11/ Bronchitis. 1920-1922 and 1933-1943: excludes croupous bronchitis. 1923-1932 and 1947-1949: includes croupous bronchitis.
- 12/ Pneumonia (all forms). 1920-1943: excludes typhoid pneumonia. 1947-1949: includes typhoid pneumonia.
- 13/ Ulcer of stomach or duodenum. 1920-1922: excludes ulcer of duodenum. 1923-1943; 1947-1949: includes ulcer of duodenum.
- 14/ Diarrhea, enteritis and ulceration of the intestines (all ages). 1920-1943: includes malnutrition 1 year of age and over. 1947-1949: excludes malnutrition 1 year of age and over.
- 15/ Diarrhea, enteritis and ulceration of the intestines (under 2 years). 1920-1943: includes malnutrition 1 to 2 years of age. 1947-1949: excludes malnutrition 1 to 2 years of age.
- 16/ Diarrhea, enteritis and ulceration of the intestines (2 years and over). 1920-1943: includes malnutrition, 2 years of age and over. 1947-1949: excludes malnutrition, 2 years of age and over.
- 17/ Nephritis. 1920-1943: includes alcoholic nephritis. 1947-1949: excludes alcoholic nephritis.
- 18/ Diseases of pregnancy, childbirth and the puerperium. 1920-1943: excludes puerperal scarlet fever. 1947-1949: includes puerperal scarlet fever.
- 19/ Senility. 1920-1922: excludes senility with mention of senile dementia, psychosis and paresis. 1923-1943 and 1947-1949: includes senility with mention of senile dementia.
- 20/ Ill-defined and unknown. 1920-1922: excludes malnutrition 1 year of age and over. Includes sudden death and found dead. 1923-1943, excludes malnutrition 1 year of age and over. Excludes sudden death and found dead. 1947-1949: includes malnutrition 1 year of age and over. Excludes sudden death and found dead.

"NA" indicates data are not available.

A dash (-) indicates no deaths were reported.

Source:

- 1920-1941: Annual and Monthly data-Annual Reports, Cabinet Bureau of Statistics.
- 1942-1943: Annual data-Annual Reports, Cabinet Bureau of Statistics.
- Monthly data-Unpublished Annual Reports, Ministry of Welfare.
- 1944-1945: Annual and Monthly data-Unpublished Annual Reports, Ministry of Welfare.
- 1946: Annual and Monthly data-Weekly Reports, Ministry of Welfare.
- 1947-1948: Annual and Monthly data-Final Annual Schedule Report, Ministry of Welfare.
- 1949: Annual and Monthly data-Final Annual Schedule Reports, Ministry of Welfare.

TABLE 16. - 1/ DEATH RATES FOR SELECTED CAUSES BY MONTH, JAPAN, 1920 - 1949
TYPHOID FEVER (1)

Year	Annual	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
• 1949	1.2	0.8	0.9	1.0	0.8	1.0	1.2	1.4	2.1	1.8	1.3	0.9	0.9
1948	1.8	1.2	1.4	1.0	1.1	1.9	1.9	2.9	3.3	2.5	1.7	1.5	1.2
1947	3.8	3.6	2.6	2.2	2.6	3.4	4.5	8.2	8.2	6.5	4.4	2.8	1.6
1946	7.2	7.9	7.4	7.9	6.6	7.0	6.2	7.1	8.6	9.1	6.9	6.3	5.1
1945	11.4	3.8	4.2	4.5	4.9	5.3	7.1	9.0	15.7	23.3	23.6	18.6	15.7
1944	10.8	4.8	6.2	6.2	6.3	8.5	10.4	13.7	18.5	17.9	13.6	12.0	11.0
1943	11.9	3.6	4.7	4.8	6.0	7.9	8.1	11.7	15.0	18.1	14.0	10.0	10.1
1942	8.7	3.6	5.5	4.9	5.3	5.7	6.4	9.4	11.7	11.5	10.2	7.7	9.8
1941	9.4	8.3	7.3	6.9	6.9	8.3	10.2	12.2	13.7	12.8	10.6	8.1	7.9
1940	9.9	7.4	6.5	5.7	5.2	6.9	8.4	11.3	14.6	16.1	14.2	12.1	9.8
1939	9.7	7.2	7.1	5.3	6.1	7.8	9.3	10.7	13.3	16.0	13.2	11.8	8.9
1938	11.0	7.3	6.6	5.8	6.3	8.2	9.4	12.7	17.2	19.9	15.6	13.1	9.6
1937	10.0	7.1	7.9	7.1	7.0	8.2	9.6	10.6	14.4	14.9	13.2	10.9	9.3
1936	9.8	6.0	5.4	4.9	5.3	6.4	8.9	11.2	15.1	17.6	15.6	11.6	9.4
1935	10.5	9.5	6.9	6.6	5.3	6.7	9.0	11.7	15.0	17.9	15.5	12.3	8.6
1934	11.9	9.3	7.9	6.6	6.4	7.0	10.5	13.5	18.7	18.7	18.9	13.9	11.3
1933	11.4	8.8	7.7	5.6	5.2	6.6	10.5	12.1	16.9	19.5	16.8	13.6	12.7
1932	10.5	9.3	7.6	6.4	5.3	6.9	8.5	10.8	14.2	16.7	16.5	12.6	10.7
1931	12.5	11.2	8.3	7.6	7.0	8.1	12.7	14.4	17.2	19.7	18.2	14.6	10.9
1930	13.0	9.8	9.1	7.8	7.2	8.4	11.1	15.5	20.8	23.1	18.3	12.5	11.4
1929	12.6	9.8	8.8	7.9	7.3	9.4	12.0	13.6	19.3	21.6	16.9	14.0	10.9
1928	14.0	9.7	10.0	8.4	7.4	9.5	12.0	15.2	21.6	23.5	21.3	18.8	12.5
1927	12.7	9.7	7.9	6.4	6.6	7.5	11.1	14.4	18.2	22.2	20.4	16.3	11.6
1926	16.1	21.0	20.7	11.7	9.0	9.6	11.4	14.3	20.3	25.2	21.2	16.8	12.1
1925	16.8	18.3	14.0	10.9	8.9	9.0	12.6	16.6	23.3	28.4	26.8	20.1	14.1
1924	23.9	24.7	23.4	17.7	14.0	14.4	16.8	24.2	38.4	36.8	30.7	23.6	21.6
1923	22.2	13.8	11.5	8.8	8.2	10.0	14.4	20.5	31.7	37.6	43.3	36.8	29.2
1922	21.7	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1921	21.1	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1920	21.6	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA

See footnotes at end of table.

TABLE 16. - 1/ DEATH RATES FOR SELECTED CAUSES BY MONTH: JAPAN, 1920 - 1949 Cont'd
PARATYPHOID FEVER (2)

Year	Annual	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
• 1949	0.1	0.1	0.0	0.2	0.1	0.1	0.2	0.2	0.2	0.3	0.1	0.1	0.1
1948	0.2	0.1	0.2	0.1	0.2	0.2	0.3	0.2	0.4	0.2	0.3	0.2	0.1
1947	0.4	0.3	0.3	0.2	0.3	0.3	0.3	0.5	0.7	0.8	0.5	0.3	0.2
1946	0.6	0.3	0.3	0.5	0.5	0.7	0.7	0.8	0.8	1.2	0.7	0.6	0.5
1945	0.8	0.2	0.4	0.3	0.4	0.4	0.6	0.8	0.9	1.4	1.3	1.4	1.1
1944	0.8	0.2	0.5	0.4	0.4	0.7	0.7	1.0	1.5	1.2	1.1	0.9	0.7
1943	0.7	0.2	0.2	0.4	0.3	0.4	0.6	0.8	1.1	1.4	1.2	0.9	0.8
1942	0.4	0.1	0.1	0.2	0.2	0.3	0.4	0.5	0.7	0.6	0.5	0.5	0.5
1941	0.4	0.2	0.3	0.2	0.5	0.3	0.5	0.4	0.7	0.6	0.5	0.4	0.4
1940	0.4	0.1	0.2	0.3	0.2	0.2	0.5	0.5	0.9	0.8	0.6	0.4	0.4
1939	0.4	0.2	0.2	0.3	0.2	0.3	0.3	0.6	0.8	0.6	0.8	0.3	0.3
1938	0.4	0.2	0.2	0.2	0.2	0.3	0.3	0.7	0.7	0.8	0.6	0.6	0.4
1937	0.4	0.1	0.2	0.1	0.2	0.4	0.4	0.6	0.8	0.6	0.6	0.3	0.2
1936	0.4	0.2	0.2	0.1	0.2	0.2	0.5	0.4	0.6	0.9	0.7	0.5	0.4
1935	0.3	0.2	0.2	0.2	0.2	0.3	0.3	0.5	0.6	0.5	0.4	0.4	0.3
1934	0.5	0.2	0.2	0.2	0.3	0.4	0.7	0.6	1.1	0.5	0.5	0.7	0.3
1933	0.5	0.3	0.2	0.1	0.2	0.1	0.5	0.8	0.7	1.4	0.7	0.6	0.3
1932	0.6	0.5	0.2	0.3	0.3	0.2	0.5	0.7	0.8	1.2	0.9	0.6	0.4
1931	0.5	0.3	0.5	0.2	0.2	0.2	0.5	0.5	0.7	0.7	0.9	0.6	0.5
1930	0.6	0.4	0.4	0.3	0.2	0.3	0.7	0.9	0.9	1.2	0.8	0.5	0.3
1929	0.6	0.3	0.2	0.2	0.3	0.5	0.5	0.7	0.7	1.2	0.9	0.7	0.5
1928	0.7	0.4	0.4	0.2	0.3	0.4	0.6	0.7	1.4	1.5	1.4	0.8	0.5
1927	0.8	0.4	0.4	0.3	0.3	0.4	0.6	0.7	1.4	1.7	1.7	0.8	0.5
1926	0.7	0.6	0.5	0.6	0.3	0.3	0.5	0.8	0.9	1.4	1.3	0.7	0.6
1925	0.7	0.6	0.3	0.2	0.4	0.4	0.5	0.8	0.9	1.8	1.4	0.8	0.7
1924	1.0	0.7	0.6	0.6	0.4	0.7	1.1	1.5	1.7	1.8	1.4	1.1	0.7
1923	1.2	0.7	0.6	0.3	0.3	0.5	0.7	1.5	2.2	2.4	2.0	1.5	1.1
1922	1.3	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1921	1.2	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1920	1.4	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA

See footnotes at end of table.

TABLE 16. $\frac{1}{2}$ DEATH RATES FOR SELECTED CAUSES BY MONTH; JAPAN, 1920 - 1949 - Cont'd
PLAGUE (3)

Year	Annual	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
* 1949	-	-	-	-	-	-	-	-	-	-	-	-	-
1948	-	-	-	-	-	-	-	-	-	-	-	-	-
1947	-	-	-	-	-	-	-	-	-	-	-	-	-
1946	-	-	-	-	-	-	-	-	-	-	-	-	-
1945	-	-	-	-	-	-	-	-	-	-	-	-	-
1944	-	-	-	-	-	-	-	-	-	-	-	-	-
1943	0.0	-	-	-	-	-	-	-	-	-	-	-	-
1942	-	-	-	-	-	-	-	-	-	-	-	-	-
1941	-	-	-	-	-	-	-	-	-	-	-	-	-
1940	-	-	-	-	-	-	-	-	-	-	-	-	-
1939	-	-	-	-	-	-	-	-	-	-	-	-	-
1938	-	-	-	-	-	-	-	-	-	-	-	-	-
1937	-	-	-	-	-	-	-	-	-	-	-	-	-
1936	-	-	-	-	-	-	-	-	-	-	-	-	-
1935	-	-	-	-	-	-	-	-	-	-	-	-	-
1934	-	-	-	-	-	-	-	-	-	-	-	-	-
1933	-	-	-	-	-	-	-	-	-	-	-	-	-
1932	-	-	-	-	-	-	-	-	-	-	-	-	-
1931	-	-	-	-	-	-	-	-	-	-	-	-	-
1930	-	-	-	-	-	-	-	-	-	-	-	-	-
1929	0.0	-	-	-	-	0.0	0.0	-	-	-	-	-	-
1928	-	-	-	-	-	-	-	-	-	-	-	-	-
1927	-	-	-	-	-	-	-	-	-	-	-	-	-
1926	0.0	-	-	-	-	-	-	0.1	0.0	-	-	-	-
1925	-	-	-	-	-	-	-	-	-	-	-	-	-
1924	0.0	0.0	-	-	-	-	0.0	0.0	-	-	-	-	-
1923	0.0	-	-	-	-	-	-	-	-	0.0	-	-	-
1922	0.1	-	-	-	-	-	-	0.2	0.0	0.0	0.6	0.5	0.1
1921	0.0	-	-	0.0	-	-	-	-	0.0	-	-	-	-
1920	0.0	-	-	-	-	-	-	-	-	0.1	0.1	0.1	0.0

See footnotes at end of table.

TABLE 16. 1/ DEATH RATES FOR SELECTED CAUSES BY MONTH: JAPAN, 1920 - 1949 Cont'd
CHOLERA (4)

Year	Annual	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
* 1949	-	-	-	-	-	-	-	-	-	-	-	-	-
1948	-	-	-	-	-	-	-	-	-	-	-	-	-
1947	-	-	-	-	-	-	-	-	-	-	-	-	-
1946	0.7	-	-	-	-	-	0.8	1.6	3.7	1.7	0.0	0.0	0.2
1945	-	-	-	-	-	-	-	-	-	-	-	-	-
1944	-	-	-	-	-	-	-	-	-	-	-	-	-
1943	0.0	-	-	-	-	-	-	-	-	-	-	-	-
1942	-	-	-	-	-	-	-	-	-	-	-	-	-
1941	-	-	-	-	-	-	-	-	-	-	-	-	-
1940	0.0	-	-	-	-	-	-	0.0	-	0.0	-	0.0	-
1939	0.0	-	-	-	-	-	0.0	-	-	-	-	-	-
1938	0.0	-	-	-	-	-	-	0.1	0.0	0.0	0.0	-	-
1937	0.0	-	-	-	-	-	-	-	-	0.1	0.0	0.0	0.0
1936	-	-	-	-	-	-	-	-	-	-	-	-	-
1935	-	-	-	-	-	-	-	-	-	-	-	-	-
1934	-	-	-	-	-	-	-	-	-	-	-	-	-
1933	-	-	-	-	-	-	-	-	-	-	-	-	-
1932	0.0	-	-	-	-	-	0.0	-	-	-	-	-	-
1931	-	-	-	-	-	-	-	-	-	-	-	-	-
1930	0.0	-	-	-	-	0.0	-	-	0.0	-	-	-	-
1929	0.2	-	-	-	-	-	-	-	0.4	1.6	0.1	0.1	0.0
1928	0.0	-	-	-	-	-	-	-	-	-	0.0	-	-
1927	0.0	-	-	-	-	-	-	-	0.0	0.0	-	-	-
1926	0.0	0.2	-	-	-	-	-	-	-	-	0.0	0.0	-
1925	0.6	-	-	-	-	-	0.0	0.1	0.1	3.3	2.6	1.0	0.3
1924	-	-	-	-	-	-	-	-	-	-	-	-	-
1923	0.1	-	-	0.1	0.0	0.1	0.0	0.1	0.1	0.1	0.0	0.1	-
1922	0.9	-	-	-	-	-	-	0.1	0.1	3.2	7.8	0.1	-
1921	0.1	0	0.0	0.0	0.0	0.0	0.1	0.1	0.1	0.1	0.1	0.0	-
1920	6.1	0.1	0.1	0.1	-	0.0	6.9	27.1	21.4	13.5	3.2	0.5	0.1

See footnotes at end of table.

TABLE 16. - 1/ DEATH RATES FOR SELECTED CAUSES BY MONTH: JAPAN, 1920 - 1949 - Cont'd
 CEREBROSPINAL (MENINGOCOCCUS) MENINGITIS (6)

Year	Annual	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
* 1949	0.6	0.5	0.6	0.8	0.7	0.5	0.7	0.4	0.7	1.2	0.5	0.3	0.3
1948	0.8	0.7	0.9	1.3	1.1	0.8	0.7	0.7	1.3	0.8	0.5	0.4	0.6
1947	1.5	0.8	1.7	3.1	2.9	2.3	1.5	1.1	1.4	1.4	1.0	0.6	0.6
1946	0.6	0.2	0.6	0.6	0.9	0.8	0.7	0.5	0.5	0.5	0.5	0.4	0.6
1945	1.5	0.3	2.1	3.4	4.1	3.3	1.9	1.3	0.4	0.7	0.5	0.4	0.3
1944	0.5	0.3	0.7	0.7	1.0	0.8	0.6	0.5	0.4	0.5	0.3	0.4	0.4
1943	0.8	0.3	0.5	0.9	0.7	1.1	0.7	0.3	0.6	0.4	0.4	0.3	0.4
1942	0.6	0.3	0.5	0.6	0.5	0.7	0.5	0.4	0.2	0.4	0.5	0.4	0.4
1941	0.7	0.6	0.9	1.2	1.1	1.0	0.7	0.6	0.7	0.6	0.5	0.6	0.4
1940	1.0	0.8	1.0	1.6	1.9	1.5	1.2	0.7	0.7	0.6	0.6	0.8	0.6
1939	1.3	0.6	1.2	1.8	2.3	2.1	1.4	1.2	1.5	1.3	0.6	0.4	0.6
1938	0.8	0.5	1.0	1.0	1.3	1.4	0.9	0.7	0.7	1.0	0.6	0.4	0.3
1937	0.7	0.5	1.0	0.7	0.9	0.5	0.8	0.6	1.2	0.7	0.5	0.3	0.5
1936	0.9	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1935	1.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1934	0.9	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1933	0.4	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1932	0.2	0.1	0.2	0.2	0.3	0.3	0.2	0.2	0.4	0.3	0.4	0.2	0.2
1931	0.3	0.1	0.2	0.4	0.5	0.2	0.2	0.2	0.2	0.2	0.3	0.3	0.2
1930	0.3	0.1	0.2	0.2	0.2	0.3	0.2	0.2	0.7	0.3	0.3	0.2	0.2
1929	0.3	0.1	0.2	0.5	0.4	0.5	0.3	0.3	0.7	0.4	0.1	0.2	0.2
1928	0.3	0.2	0.2	0.2	0.3	0.4	0.3	0.3	0.3	0.4	0.2	0.1	0.3
1927	0.3	0.3	0.2	0.4	0.4	0.5	0.4	0.3	0.5	0.5	0.5	0.3	0.2
1926	0.4	0.2	0.3	0.3	0.3	0.5	0.4	0.4	0.5	0.8	0.4	0.2	0.3
1925	0.5	0.3	0.5	0.6	0.8	0.8	0.5	0.4	0.4	0.5	0.3	0.2	0.2
1924	2.3	0.5	0.6	0.7	0.6	0.4	0.4	0.5	13.5	8.0	1.1	0.4	0.5
1923	0.7	0.4	0.6	1.1	0.9	0.8	0.9	0.7	0.4	0.8	0.8	0.3	0.3
1922	1.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1921	0.8	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1920	1.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA

See footnotes at end of table.

TABLE 16. - 1/ DEATH RATES FOR SELECTED CAUSES BY MONTH: JAPAN, 1920 - 1949 - Cont'd
ANTRAX (7)

Year	Annual	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
* 1949	-	-	-	-	-	-	-	-	-	-	-	-	-
1948	-	-	-	-	-	-	-	-	-	-	-	-	-
1947	0.0	-	-	-	0.0	0.0	-	-	-	-	-	-	-
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	0.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	0.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1941	0.0	-	-	-	0.0	-	-	0.1	0.0	0.0	-	-	0.0
1940	0.0	0.0	0.0	0.1	0.0	0.0	-	0.0	0.0	-	-	0.0	-
1939	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.1	-	0.0	0.0	0.0	-
1938	0.0	0.0	0.0	0.0	-	0.0	0.0	0.0	0.0	0.0	-	-	0.0
1937	0.0	0.0	0.1	0.0	0.1	-	-	-	0.0	-	0.0	-	-
1936	0.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1935	0.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1934	0.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1933	0.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1932	0.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1931	0.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1930	0.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1929	0.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1928	0.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1927	0.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1926	0.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1925	0.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1924	0.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1923	0.1	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1922	0.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1921	0.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1920	0.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA

See footnotes at end of table.

TABLE 16. - $\frac{1}{L}$ DEATH RATES FOR SELECTED CAUSES BY MONTH: JAPAN, 1920 - 1949 - Cont'd
3/ SCARLET (8)

Year	Annual	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
+1949	0.1	0.1	0.1	0.0	0.1	0.1	0.1	0.0	0.1	0.0	0.0	0.1	0.0
1948	0.1	0.0	0.0	0.1	0.1	0.1	0.0	0.0	0.1	0.0	0.0	0.1	0.1
1947	0.1	0.1	0.2	0.1	0.1	0.2	0.1	0.1	0.1	0.0	0.1	0.1	0.1
1946	0.1	0.1	0.2	0.3	0.3	0.2	0.1	0.1	0.0	0.1	0.0	0.1	0.1
1945	0.1	0.1	0.2	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.2	0.1
1944	0.2	0.2	0.2	0.2	0.4	0.1	0.1	0.1	0.1	0.2	0.1	0.1	0.1
1943	0.3	0.3	0.3	0.3	0.2	0.2	0.3	0.2	0.2	0.2	0.2	0.1	0.2
1942	0.3	0.3	0.4	0.2	0.3	0.3	0.3	0.3	0.4	0.2	0.2	0.3	0.5
1941	0.4	0.5	0.6	0.4	0.5	0.3	0.2	0.2	0.2	0.3	0.3	0.4	0.5
1940	0.5	0.6	0.7	0.8	0.7	0.7	0.5	0.3	0.2	0.3	0.3	0.4	0.7
1939	0.7	0.7	0.9	0.8	0.7	0.6	0.6	0.6	0.4	0.5	0.3	0.9	1.0
1938	0.6	0.8	0.8	0.6	0.4	0.5	0.6	0.2	0.4	0.3	0.6	0.8	0.7
1937	0.6	0.5	0.8	0.7	0.9	0.8	0.7	0.6	0.3	0.4	0.4	0.7	0.8
1936	0.7	0.7	1.0	0.8	0.7	0.9	0.7	0.2	0.3	0.4	0.7	0.9	0.8
1935	0.7	1.0	1.0	0.9	0.7	0.6	0.7	0.5	0.5	0.6	0.6	0.8	0.7
1934	0.7	0.9	0.9	1.1	0.8	0.8	0.7	0.5	0.4	0.4	0.6	0.8	0.9
1933	0.6	0.6	0.6	0.8	0.6	0.6	0.4	0.5	0.5	0.4	0.5	0.9	0.7
1932	0.5	0.7	0.6	0.6	0.7	0.5	0.5	0.3	0.3	0.3	0.3	0.6	0.8
1931	0.5	0.7	0.6	0.7	0.6	0.4	0.5	0.5	0.3	0.3	0.4	0.5	0.5
1930	0.5	0.4	0.5	0.4	0.5	0.5	0.5	0.2	0.2	0.2	0.4	1.0	0.8
1929	0.4	0.4	0.6	0.5	0.5	0.5	0.3	0.4	0.1	0.2	0.3	0.5	0.4
1928	0.4	0.4	0.6	0.4	0.4	0.3	0.3	0.3	0.3	0.3	0.4	0.2	0.4
1927	0.4	0.4	0.4	0.3	0.5	0.4	0.4	0.3	0.1	0.3	0.4	0.4	0.5
1926	0.5	0.7	0.6	0.5	0.5	0.5	0.8	0.8	0.4	0.2	0.3	0.3	0.4
1925	0.5	0.4	0.3	0.5	0.9	0.6	0.6	0.5	0.3	0.2	0.2	0.7	0.6
1924	0.2	0.3	0.1	0.3	0.3	0.3	0.2	0.1	0.1	0.1	0.2	0.3	0.4
1923	0.2	0.3	0.2	0.2	0.2	0.2	0.2	0.1	0.1	0.1	0.1	0.2	0.2
1922	0.2	0.2	0.2	0.2	0.3	0.3	0.2	0.1	0.1	0.1	0.3	0.0	0.2
1921	0.2	0.3	0.2	0.2	0.1	0.2	0.2	0.1	-	0.1	0.2	0.2	0.1
1920	0.2	0.2	0.3	0.1	0.2	0.2	0.2	0.1	0.1	0.1	0.2	0.2	0.1
See footnotes at end of table.													

TABLE 16. - $\frac{1}{2}$ DEATH RATES FOR SELECTED CAUSES BY MONTH: JAF/N, 1920 - 1949 - Cont'd
WHOPING COUGH (9)

Year	Annual	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
*1949	11.1	8.1	8.8	7.6	8.6	10.4	13.7	15.3	17.0	13.6	8.5	8.7	12.9
1948	5.9	6.9	5.9	4.2	4.7	5.2	6.3	8.3	8.1	6.4	3.8	4.5	6.6
1947	21.8	30.0	46.2	33.0	25.3	24.4	25.1	24.1	21.0	13.3	6.6	5.5	6.9
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	12.7	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	12.9	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1941	14.7	13.1	14.7	12.5	11.8	13.0	16.0	21.6	16.9	16.9	11.1	11.0	14.2
1940	12.2	14.6	13.6	10.3	10.4	11.3	13.6	16.6	17.6	12.6	7.5	7.6	10.5
1939	14.1	12.7	13.6	14.2	14.0	13.9	16.1	19.3	21.8	14.5	9.5	8.2	11.5
1938	12.5	14.7	14.8	11.2	9.6	9.7	12.9	16.0	18.0	14.5	9.5	7.9	10.7
1937	15.5	14.8	14.5	14.5	14.3	16.5	19.9	22.2	22.9	16.6	9.8	7.5	12.1
1936	13.9	14.6	15.6	12.1	10.0	11.1	15.1	20.8	22.0	17.0	9.9	8.1	10.4
1935	17.6	16.3	18.2	16.3	17.4	18.7	23.3	28.0	24.6	17.3	10.7	8.8	12.1
1934	12.5	8.3	9.1	8.3	7.3	8.8	13.8	19.6	22.7	16.9	10.8	9.9	13.8
1933	9.1	12.7	12.1	9.8	8.4	9.2	10.7	11.2	11.9	9.1	5.3	4.0	5.6
1932	22.1	22.1	24.3	21.5	22.0	24.2	32.7	33.2	31.7	21.2	11.1	9.1	11.8
1931	12.6	8.4	9.0	7.7	8.4	9.1	12.9	17.0	19.1	19.5	12.8	10.8	16.6
1930	11.5	13.0	11.8	11.4	10.9	11.1	14.5	17.2	17.4	12.6	6.2	5.4	6.9
1929	15.5	13.8	14.1	14.1	15.1	15.1	18.6	23.7	26.9	17.8	8.9	8.0	9.2
1928	18.2	17.9	20.4	18.0	16.9	17.0	22.0	25.1	27.1	19.9	12.0	9.6	12.9
1927	14.8	12.0	11.2	10.1	11.0	13.3	17.2	22.6	25.2	19.9	10.6	10.2	14.3
1926	12.0	9.5	10.6	11.8	11.4	12.3	13.5	17.3	17.9	15.0	9.3	6.5	9.1
1925	14.2	17.5	18.5	14.2	11.4	12.5	17.1	20.6	19.6	14.6	8.8	6.8	8.3
1924	17.4	17.1	15.3	15.7	15.7	18.7	22.2	25.1	28.3	18.8	10.4	8.5	13.1
1923	14.6	7.1	8.5	8.7	10.0	13.3	19.5	24.7	28.1	21.1	11.9	9.6	11.9
1922	10.9	10.8	12.9	10.0	7.7	8.3	12.4	17.2	18.9	13.1	7.3	5.8	6.7
1921	12.3	10.2	11.5	9.7	10.0	10.9	14.1	16.3	21.6	16.8	9.6	7.2	9.2
1920	14.2	13.8	15.0	13.5	12.0	13.3	17.6	23.1	22.7	14.8	7.5	7.1	9.5

See footnotes at end of table.

TABLE 16. - 1/ DEATH RATES FOR SELECTED CAUSES BY MONTH: JAPAN, 1920 - 1949 - Cont'd
1/ DIPHTHERIA (10)

Year	Annual	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
1949	2.0	3.5	3.1	2.6	2.4	1.6	1.1	1.2	0.7	0.9	1.4	2.6	3.0
1948	2.4	4.1	3.7	3.0	2.0	1.4	1.2	0.7	0.9	1.4	2.1	3.7	4.3
1947	4.3	8.2	7.2	6.5	5.0	4.1	2.8	1.9	1.4	1.6	3.3	4.6	5.8
1946	5.2	11.8	9.5	6.8	5.1	3.6	3.2	2.4	1.9	2.7	4.6	5.2	6.2
1945	11.2	14.4	17.4	15.1	12.9	11.1	8.7	6.0	4.5	6.0	9.2	14.7	15.0
1944	8.5	11.4	11.2	10.4	7.9	5.4	4.9	3.9	4.5	4.6	8.3	11.7	17.8
1943	9.6	10.6	10.6	10.5	9.3	7.7	6.1	4.1	2.3	2.9	5.9	8.7	12.6
1942	7.0	7.5	9.1	7.8	6.2	3.8	3.1	2.6	3.1	3.3	5.6	8.3	14.3
1941	6.8	10.9	11.1	7.8	6.9	4.9	4.0	2.7	3.1	4.2	6.7	9.1	10.9
1940	6.6	10.2	10.5	8.8	6.5	4.8	3.6	2.7	2.2	3.9	6.2	8.6	11.1
1939	7.4	10.7	11.2	11.7	9.4	6.4	5.6	2.7	2.6	3.0	5.8	8.0	10.1
1938	5.8	9.5	9.4	7.9	6.5	4.4	3.5	2.7	2.1	2.8	4.2	6.9	10.1
1937	6.0	8.9	9.1	8.4	6.7	5.1	4.7	2.6	2.1	2.9	4.9	7.4	9.1
1936	6.2	10.3	10.4	9.3	6.9	4.9	3.1	2.4	1.9	2.8	4.9	8.4	9.1
1935	6.4	10.1	10.6	8.7	7.6	5.8	4.3	2.6	3.0	3.0	5.1	7.4	9.4
1934	7.7	13.9	12.5	12.0	8.9	6.3	4.6	2.7	2.6	3.5	6.4	8.6	10.3
1933	8.1	10.7	12.7	12.3	9.9	7.4	5.3	3.3	2.6	3.9	7.0	9.7	12.3
1932	6.8	9.6	9.6	10.0	7.5	5.5	4.6	3.1	2.6	2.9	5.8	9.5	11.0
1931	7.0	10.9	11.8	10.4	8.4	6.4	5.1	3.4	2.6	3.0	5.1	7.7	9.7
1930	6.3	9.4	9.7	8.5	6.1	4.8	3.4	2.6	2.4	3.7	7.1	8.6	9.8
1929	7.4	12.3	13.8	13.0	9.4	6.0	4.3	2.4	1.9	2.7	5.6	8.7	9.3
1928	7.0	10.6	10.9	10.7	7.3	5.7	4.7	3.3	2.8	3.2	6.0	8.8	10.4
1927	6.4	8.9	9.9	9.0	7.4	6.0	4.5	2.9	1.7	3.2	5.1	7.9	10.0
1926	6.0	8.1	8.6	8.3	6.8	5.6	4.0	2.9	2.1	2.0	4.5	8.4	9.5
1925	6.0	10.0	10.1	10.2	6.8	5.9	3.9	2.3	1.9	2.8	4.5	6.1	8.2
1924	6.0	9.3	8.2	9.2	6.7	5.0	4.2	2.8	2.1	3.1	4.7	8.0	9.5
1923	5.8	10.9	8.8	7.9	6.3	4.4	3.8	2.8	1.8	3.1	4.5	6.8	8.8
1922	6.2	10.1	10.7	8.7	5.2	4.2	3.8	2.3	1.9	2.7	4.9	8.6	11.2
1921	6.8	11.2	10.4	8.7	6.6	4.4	3.7	3.3	2.7	3.6	6.5	10.0	10.1
1920	6.8	10.8	9.1	7.6	5.9	5.1	4.6	2.8	2.7	3.6	6.8	10.4	12.8

See footnotes at end of table.

TABLE 16. - 1/ DEATH RATES FOR SELECTED CAUSES BY MONTH: JAPAN, 1920 - 1949 - Cont'd
TETANUS (12)

Year	Annual	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
* 1949	2.4	1.9	2.3	1.9	2.3	2.2	2.5	2.7	3.0	3.2	2.4	2.1	1.9
1948	2.7	2.2	1.8	2.0	3.1	2.6	2.8	3.5	3.4	3.0	2.2	2.8	2.5
1947	2.8	2.9	2.3	2.1	2.7	2.5	2.7	3.7	3.5	4.1	2.9	2.9	1.9
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	2.7	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	2.5	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1941	2.4	1.9	1.9	1.7	1.9	2.3	2.2	3.7	3.0	3.1	2.6	2.1	2.6
1940	2.6	2.2	2.0	2.0	2.5	2.5	2.6	3.1	3.2	2.9	3.0	2.2	2.4
1939	2.4	2.0	1.8	2.1	2.4	2.0	2.3	2.8	2.9	3.1	2.8	2.1	2.2
1938	2.5	2.2	2.2	2.1	2.3	2.6	2.6	2.7	3.0	3.2	2.4	2.3	2.3
1937	2.7	2.2	2.4	2.4	2.6	2.7	2.8	3.4	3.5	3.3	2.7	2.6	2.1
1936	2.8	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1935	3.1	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1934	3.2	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1933	3.4	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1932	3.7	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1931	3.7	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1930	3.9	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1929	4.2	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1928	4.4	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1927	4.5	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1926	4.7	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1925	4.9	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1924	5.1	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1923	5.5	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1922	5.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1921	5.4	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1920	5.5	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA

See footnotes at end of table.

TABLE 16. - 1/ DEATH RATES FOR SELECTED CAUSES BY MONTH: JAPAN, 1920 - 1949 - Cont'd
2/ TUBERCULOSIS (ALL FORMS) (13-22)

Year	Annual	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
*1949	168.8	159.5	160.3	175.9	184.9	186.3	181.9	173.2	169.5	170.6	161.0	155.8	146.7
1948	179.4	161.1	169.0	182.9	195.7	193.5	192.8	186.1	183.8	181.2	173.9	168.9	162.1
1947	187.5	182.8	189.1	190.3	200.6	208.2	197.3	198.4	190.1	183.7	177.7	167.9	164.1
1946	264.2	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	280.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	240.2	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	230.8	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	218.8	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1941	210.9	197.1	200.8	197.7	207.4	208.4	212.9	221.9	227.7	224.3	216.5	208.8	206.0
1940	212.4	206.4	208.9	210.9	205.6	209.9	210.4	221.1	225.0	223.2	216.0	208.5	202.4
1939	215.9	199.4	201.5	201.8	205.2	206.1	214.9	229.9	232.5	231.2	226.4	216.6	215.4
1938	209.0	195.9	199.1	201.8	204.4	205.9	212.7	224.1	224.3	225.0	210.8	205.8	197.8
1937	203.7	193.3	195.1	202.6	206.7	208.1	213.7	210.6	215.9	212.8	200.7	194.1	190.3
1936	206.8	200.7	200.5	199.1	203.2	210.8	213.2	218.4	215.1	216.2	209.2	200.0	195.0
1935	190.8	179.0	178.4	182.1	183.8	191.9	195.9	195.7	204.3	202.4	198.4	187.5	185.3
1934	192.6	182.2	186.9	192.4	193.4	201.7	200.3	198.8	202.2	196.8	194.9	185.5	175.5
1933	188.0	172.1	182.8	187.5	189.2	193.3	195.0	195.5	203.0	201.5	184.1	178.1	175.7
1932	179.5	165.3	178.5	178.7	179.9	182.3	184.2	188.9	190.7	190.3	180.3	170.1	165.0
1931	186.4	172.6	181.9	184.6	184.2	189.5	192.1	193.9	199.1	197.5	189.9	179.3	171.4
1930	185.6	180.7	180.8	180.5	194.7	198.1	191.5	192.7	192.4	190.5	178.1	171.2	167.0
1929	194.5	176.7	183.8	190.7	195.4	197.4	202.2	207.0	206.4	206.6	197.0	189.3	180.8
1928	190.8	177.0	187.9	192.2	195.3	197.0	188.8	198.4	195.9	203.8	190.7	183.9	174.8
1927	193.6	166.1	175.3	181.5	194.4	200.6	202.8	205.6	210.3	210.5	200.8	184.2	174.8
1926	185.9	177.2	180.1	184.2	187.2	196.6	195.6	192.7	197.8	197.5	197.1	175.5	162.3
1925	194.1	179.2	194.8	188.0	195.3	198.8	199.2	208.2	208.0	203.0	200.8	179.6	174.5
1924	193.9	179.1	183.2	191.5	196.8	202.6	198.5	207.8	214.0	210.1	199.3	180.7	172.8
1923	203.1	185.7	191.4	202.0	205.0	204.8	211.3	219.7	221.6	221.6	208.1	189.1	176.5
1922	218.7	161.5	213.3	229.4	226.5	233.3	227.7	227.4	242.4	233.4	215.1	197.8	185.8
1921	212.9	186.9	201.7	204.8	207.8	215.3	213.9	225.6	230.3	237.2	227.2	205.1	189.3
1920	223.7	222.2	219.3	223.7	224.6	225.5	222.1	234.6	243.5	244.7	224.4	205.9	193.1

See footnotes at end of table.

TABLE 16. - 1/ DEATH RATES FOR SELECTED CAUSES BY MONTH: JAPAN, 1920 - 1949 - Cont'd
TIFPROSY (23)

Year	Annual	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
1949	0.2	0.2	0.2	0.2	0.3	0.2	0.2	0.2	0.1	0.1	0.2	0.3	0.2
1948	0.2	0.2	0.3	0.3	0.1	0.3	0.1	0.1	0.2	0.1	0.1	0.2	0.2
1947	0.2	0.3	0.3	0.3	0.4	0.3	0.1	0.2	0.2	0.3	0.3	0.2	0.2
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	0.2	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	0.2	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1941	0.3	0.2	0.2	0.3	0.4	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.2
1940	0.4	0.5	0.5	0.5	0.5	0.2	0.4	0.3	0.3	0.4	0.2	0.3	0.3
1939	0.4	0.6	0.6	0.6	0.5	0.3	0.4	0.4	0.3	0.4	0.4	0.4	0.4
1938	0.5	0.5	0.8	0.4	0.4	0.4	0.5	0.5	0.4	0.5	0.2	0.5	0.5
1937	0.6	0.7	0.4	0.8	0.6	0.5	0.5	0.4	0.5	0.6	0.6	0.4	0.5
1936	0.6	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1935	0.7	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1934	0.7	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1933	0.8	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1932	0.8	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1931	1.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1930	1.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1929	1.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1928	1.1	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1927	1.2	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1926	1.3	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1925	1.3	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1924	1.4	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1923	1.6	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1922	1.7	2.0	1.8	2.3	1.5	1.7	1.8	1.4	1.8	1.5	1.6	1.5	1.4
1921	1.9	1.9	2.4	2.0	1.9	1.5	1.8	1.9	1.8	2.4	2.0	1.8	1.8
1920	2.0	2.2	2.8	2.3	2.1	1.7	1.7	1.7	1.8	2.2	1.8	1.7	2.1

See footnote at end of table.

TABLE 16. - 2/ DEATH RATES FOR SELECTED CAUSES BY MONTH: JAPAN, 1920 - 1949 - Cont'd
GIARDENS (26*)

Year	Annual	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
*1949	0.0	-	-	-	-	0.0	-	-	-	-	-	-	-
1948	-	-	-	-	-	-	-	-	-	-	-	-	-
1947	-	-	-	-	-	-	-	-	-	-	-	-	-
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	-	-	-	-	-	-	-	-	-	-	-	-	-
1942	-	-	-	-	-	-	-	-	-	-	-	-	-
1941	-	-	-	-	-	-	-	-	-	-	-	-	-
1940	-	-	-	-	-	-	-	-	-	-	-	-	-
1939	-	-	-	-	-	-	-	-	-	-	-	-	-
1938	-	-	-	-	-	-	-	-	-	-	-	-	-
1937	-	-	-	-	-	-	-	-	-	-	-	-	-
1936	-	-	-	-	-	-	-	-	-	-	-	-	-
1935	0.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1934	0.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1933	-	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1932	-	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1931	0.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1930	0.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1929	-	-	-	-	-	-	-	-	-	-	-	-	-
1928	-	-	-	-	-	-	-	-	-	-	-	-	-
1927	-	-	-	-	-	-	-	-	-	-	-	-	-
1926	0.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1925	0.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1924	0.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1923	0.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1922	0.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1921	0.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1920	-	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA

See footnotes at end of table.

TABLE 16. - 1/ DEATH RATES FOR SELECTED CAUSES BY MONTH: JAPAN, 1920 - 1949 - Cont'd
DYSENTERY (27)

Year	Annual	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
1949	9.5	0.7	0.9	0.9	1.6	3.2	7.9	26.5	34.8	24.2	8.2	2.8	1.6
1948	6.4	0.6	0.9	1.0	1.9	2.9	6.9	20.5	21.9	12.6	5.1	1.6	1.0
1947	12.3	1.7	1.5	1.5	2.4	4.3	9.1	39.6	48.8	24.7	8.5	2.7	1.0
1946	17.6	2.3	0.8	1.2	1.3	2.6	5.5	26.1	56.5	52.6	37.7	15.6	6.8
1945	28.8	0.8	0.9	1.0	1.7	3.2	7.4	36.1	81.7	107.0	57.3	32.3	13.4
1944	15.4	0.6	1.3	1.3	1.9	3.7	9.2	30.2	51.1	39.9	24.0	13.3	7.5
1943	17.0	1.8	2.1	2.4	3.5	5.9	13.0	33.9	43.6	31.2	18.3	6.9	4.6
1942	19.4	2.8	4.0	4.8	6.5	11.9	16.9	37.3	37.7	37.3	22.4	13.9	8.5
1941	22.3	5.1	5.1	6.1	9.2	14.9	28.9	62.2	62.3	39.6	20.6	8.6	5.2
1940	30.6	8.2	9.0	10.1	12.8	20.8	40.3	82.9	83.8	50.5	26.9	12.7	6.9
1939	34.8	7.2	6.7	8.8	10.8	18.3	34.2	75.7	98.9	72.1	48.0	23.9	10.8
1938	30.9	5.4	5.6	7.7	11.4	16.7	35.2	77.1	89.1	61.9	33.2	15.2	9.3
1937	27.8	6.0	6.5	7.8	11.4	16.9	28.0	59.0	70.0	59.6	39.9	18.2	7.7
1936	23.8	4.4	4.5	7.0	9.3	13.2	24.7	56.3	71.3	48.1	25.9	12.7	7.1
1935	23.0	6.7	5.6	7.2	9.5	15.3	26.7	58.8	61.0	41.6	23.9	11.9	5.9
1934	22.7	3.8	4.5	6.4	9.4	14.0	24.9	57.3	62.4	43.1	26.6	11.7	6.2
1933	22.1	4.6	5.1	5.3	7.3	12.6	25.5	51.3	61.6	45.2	26.9	11.5	6.3
1932	20.4	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1931	19.9	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1930	20.2	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1929	21.1	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1928	19.2	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1927	16.6	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1926	14.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1925	12.6	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1924	14.7	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1923	16.2	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1922	15.9	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1921	13.9	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1920	14.6	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA

See footnotes at end of table.

TABLE 16. - $\frac{1}{2}$ / DEATH RATES FOR SELECTED CAUSES BY MONTH: JAPAN, 1920 - 1949 - Cont'd
MALARIA (28)

Year	Annual	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
* 1949	0.1	0.2	0.2	0.3	0.1	0.1	0.1	0.1	0.1	0.3	0.1	0.1	0.1
1948	0.3	0.3	0.4	0.3	0.3	0.4	0.2	0.2	0.3	0.4	0.3	0.2	0.0
1947	0.6	0.6	0.6	0.6	0.6	0.6	0.7	0.5	0.5	0.6	0.5	0.5	0.5
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	0.3	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	0.3	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1941	0.2	0.3	0.2	0.2	0.2	0.1	0.2	0.1	0.2	0.4	0.2	0.2	0.2
1940	0.3	0.4	0.5	0.3	0.3	0.4	0.2	0.3	0.3	0.3	0.3	0.3	0.2
1939	0.3	0.6	0.4	0.2	0.2	0.3	0.2	0.2	0.3	0.4	0.4	0.4	0.2
1938	0.3	0.1	0.1	0.1	0.1	0.1	0.1	0.2	0.5	0.4	0.6	0.4	0.6
1937	0.1	0.0	0.1	0.0	0.1	0.1	0.1	0.1	0.2	0.1	0.1	0.1	0.1
1936	0.1	-	0.1	0.1	-	0.1	0.2	0.2	0.2	0.1	0.1	0.1	0.1
1935	0.1	0.1	0.0	0.0	0.0	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
1934	0.1	0.1	0.2	0.1	0.1	0.1	0.1	0.0	0.1	0.1	0.1	0.1	0.1
1933	0.1	0.1	0.2	0.1	0.1	0.1	0.1	0.1	0.2	0.1	0.1	0.1	0.1
1932	0.1	0.1	0.1	0.0	0.0	0.1	0.1	0.1	0.1	0.2	0.2	0.1	0.1
1931	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.0	0.1	0.1	0.1	0.0	0.1
1930	0.1	0.1	0.1	0.1	0.1	0.1	0.2	0.1	0.1	0.0	0.1	0.1	0.1
1929	0.2	0.2	0.1	0.2	0.1	0.1	0.4	0.1	0.2	0.2	0.2	0.1	0.1
1928	0.3	0.1	0.1	0.1	0.1	0.1	0.1	0.3	0.6	0.7	0.4	0.3	0.2
1927	0.3	0.2	0.1	0.1	0.2	0.1	0.3	0.3	0.7	0.4	0.4	0.4	0.1
1926	0.3	0.2	0.2	0.1	0.1	0.2	0.2	0.2	0.3	0.5	0.8	0.5	0.5
1925	0.2	0.2	0.1	0.1	0.2	0.1	0.3	0.2	0.2	0.5	0.3	0.1	0.1
1924	0.3	0.3	0.2	0.1	0.2	0.1	0.5	0.2	0.3	0.5	0.2	0.3	0.2
1923	0.3	0.2	0.1	0.1	0.1	0.4	0.4	0.4	0.5	0.4	0.3	0.4	0.2
1922	0.3	0.2	0.2	0.1	0.2	0.1	0.3	0.5	0.8	0.6	0.4	0.2	0.3
1921	0.4	0.4	0.1	0.2	0.2	0.4	0.4	0.6	0.8	0.4	0.5	0.2	0.2
1920	0.4	0.5	0.2	0.2	0.2	0.2	0.3	0.8	1.0	0.7	0.5	0.4	0.4

See footnotes at end of table.

TABLE 16. - 1/ DEATH RATES FOR SELECTED CAUSES BY MONTH: JAPAN, 1920 - 1949 - Cont'd
6/ SYPHILIS (30)

Year	Annual	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
*1949	6.7	7.4	7.4	7.4	6.9	5.7	5.8	5.8	5.6	5.9	6.8	7.6	7.8
**1948	5.4	5.3	6.3	6.1	5.1	4.7	4.7	4.7	4.7	5.1	5.3	5.9	7.3
**1947	5.7	7.9	7.0	6.2	5.4	5.3	5.8	5.5	4.5	4.5	5.1	5.6	5.7
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	10.6	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	10.7	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1941	10.6	11.9	12.4	11.1	10.4	9.4	9.2	9.4	8.8	10.2	11.0	10.8	12.9
1940	10.7	14.3	11.9	12.2	10.3	9.0	9.0	8.8	9.3	10.6	10.5	10.7	11.6
1939	11.3	13.7	12.7	11.5	10.5	9.4	9.9	10.0	10.3	9.9	12.5	11.9	13.2
1938	12.3	15.8	14.6	13.1	11.1	11.1	9.6	11.2	11.2	11.9	12.7	13.0	12.2
1937	13.3	14.9	14.2	14.9	13.3	11.9	11.2	11.8	12.2	12.8	12.7	14.0	15.8
1936	13.4	17.3	16.3	15.1	13.1	10.9	11.6	11.5	11.6	11.8	13.5	13.3	15.0
1935	13.5	15.1	15.8	14.6	12.4	11.8	11.2	11.3	12.0	13.2	13.2	14.7	16.4
1934	13.7	16.8	14.9	15.4	14.3	11.6	11.3	11.8	11.2	11.8	14.7	15.9	15.1
1933	14.2	16.6	17.8	15.4	12.9	12.3	13.0	12.2	12.6	13.2	13.6	15.1	15.6
1932	14.4	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1931	15.1	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1930	15.2	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1929	16.2	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1928	17.5	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1927	17.6	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1926	18.3	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1925	17.9	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1924	18.5	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1923	19.6	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1922	16.2	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1921	18.4	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1920	18.2	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA

See footnotes at end of table.

TABLE 16. - 1/ DEATH RATES FOR SELECTED CAUSES BY MONTH: JAPAN, 1920 - 1940 - Continued
INFLUENZA (33)

Year	Annual	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
*1949	0.6	0.8	1.2	0.9	0.8	0.7	0.4	0.2	0.2	0.3	0.3	0.5	1.1
1948	0.6	1.3	1.0	1.3	0.8	0.4	0.5	0.3	0.2	0.3	0.4	0.4	0.7
1947	2.4	6.5	11.4	5.2	2.0	1.2	0.7	0.5	0.2	0.3	0.5	0.4	1.0
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	4.6	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	5.4	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1941	5.9	14.7	23.7	13.4	5.5	2.8	1.7	1.1	1.0	1.0	1.3	2.3	4.2
1940	4.4	10.4	12.5	9.2	5.1	2.7	2.0	10.0	0.9	0.6	1.1	1.8	5.5
1939	8.2	13.8	21.5	30.4	15.5	4.8	2.0	0.9	1.0	0.7	1.4	1.9	5.1
1938	10.7	24.6	45.9	31.0	11.0	4.8	1.8	0.9	0.9	1.4	1.4	1.9	5.7
1937	4.2	7.3	9.8	7.4	5.3	3.2	2.1	1.2	0.9	1.4	1.7	2.5	8.2
1936	12.4	43.9	58.4	24.3	8.2	3.7	1.5	1.1	0.6	0.9	1.3	1.9	4.0
1935	4.4	8.7	9.8	9.5	6.0	3.7	1.7	1.4	1.5	1.1	1.6	2.1	6.3
1934	14.8	45.2	58.0	40.8	15.0	6.3	2.5	1.2	0.9	0.9	1.7	2.9	5.7
1933	7.1	10.6	14.8	19.0	13.5	6.5	2.7	1.3	1.1	1.4	1.9	3.0	9.6
1932	8.1	10.3	12.5	20.2	22.2	11.9	3.9	1.8	1.4	1.7	2.0	3.2	6.1
1931	24.0	54.6	89.9	64.2	34.0	20.2	9.2	3.5	1.7	1.4	2.3	3.6	7.8
1930	8.1	11.2	14.6	18.1	16.4	11.6	5.4	2.3	2.1	2.1	2.2	3.4	8.1
1929	13.4	19.2	29.0	42.6	28.1	14.4	5.9	3.1	2.0	2.6	2.8	3.7	7.9
1928	15.4	12.9	26.0	62.1	40.1	15.6	5.0	2.1	1.8	1.8	2.8	4.2	10.9
1927	13.2	26.1	40.7	28.7	18.3	17.0	9.0	3.4	2.5	2.3	2.7	3.2	6.6
1926	7.6	11.7	14.0	14.9	11.8	9.7	5.0	2.3	1.8	2.2	3.2	4.8	10.0
1925	18.1	25.1	90.4	59.1	18.5	6.8	3.7	2.0	1.4	2.2	2.7	3.7	7.1
1924	11.1	15.1	25.0	26.0	19.5	12.8	7.6	3.0	3.3	3.1	3.2	4.9	6.8
1923	11.2	14.6	27.3	29.1	20.5	12.1	6.9	3.8	2.3	2.9	3.8	4.4	8.1
1922	22.1	23.9	92.5	85.3	29.7	11.1	5.2	3.2	2.4	3.0	2.9	4.3	7.1
1921	18.2	25.2	34.4	44.4	39.1	19.9	9.3	6.0	4.8	6.0	6.8	10.1	13.4
1920	193.7	834.6	699.2	374.0	235.0	105.2	27.5	9.5	6.4	7.7	9.7	12.7	20.3

See footnotes at end of table.

TABLE 16. - 1/ DEATH RATES FOR SELECTED CAUSES BY MONTH: JAPAN, 1920 - 1949 - Cont'd
SMALLPOX (34)

Year	Annual	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
1949	0.0	-	0.0	0.0	0.1	0.1	0.0	0.0	0.0	-	0.0	-	-
1948	0.0	-	0.1	0.0	-	-	0.0	0.0	-	-	-	-	-
1947	0.1	0.2	0.1	0.2	0.2	0.1	0.2	0.0	0.1	0.0	0.0	0.0	0.0
1946	3.8	1.9	6.5	10.3	12.0	8.4	3.3	0.7	0.1	0.1	0.1	0.2	1.7
1945	0.4	0.1	0.1	0.1	0.4	0.9	0.7	0.3	0.4	0.3	0.1	0.5	0.5
1944	0.1	0.0	0.1	0.2	0.1	0.2	0.1	0.0	0.0	-	0.0	-	0.0
1943	0.1	0.0	0.2	0.1	0.4	0.2	0.1	-	-	0.0	0.0	0.0	0.0
1942	0.1	0.0	0.4	0.3	0.1	0.0	0.1	-	-	-	-	-	-
1941	0.1	0.2	0.2	0.3	0.1	0.2	0.3	0.1	0.0	0.0	0.0	0.0	0.0
1940	0.1	0.0	0.0	0.3	0.4	0.1	0.1	0.0	-	0.0	-	-	0.0
1939	0.0	0.0	0.1	0.1	0.1	0.1	0.0	0.0	-	-	-	-	-
1938	0.0	-	-	-	-	-	-	-	-	-	-	0.0	0.1
1937	0.0	-	0.1	0.0	0.0	0.0	-	-	-	-	-	-	-
1936	0.0	0.0	-	-	0.1	0.1	0.1	0.0	0.0	0.1	0.0	-	0.0
1935	0.0	0.1	0.0	0.0	0.0	0.1	-	-	-	-	-	0.0	-
1934	0.1	0.1	0.0	0.1	0.1	0.1	0.0	-	-	-	-	0.0	0.2
1933	0.1	0.0	0.2	0.1	0.2	0.2	0.1	0.1	-	-	-	-	-
1932	0.1	0.0	0.0	0.1	0.2	0.4	0.1	-	-	-	0.0	-	-
1931	0.0	0.0	-	0.0	0.0	0.0	-	-	-	-	-	-	0.0
1930	0.0	-	-	0.0	0.0	0.0	-	-	-	0.0	-	-	-
1929	0.0	-	-	0.1	0.0	0.0	0.0	-	-	-	-	-	-
1928	0.2	0.2	0.6	0.3	0.5	0.2	0.1	-	-	-	0.0	0.0	-
1927	0.1	0.2	0.2	0.1	0.1	0.2	0.2	0.2	0.1	0.1	0.1	0.0	0.4
1926	0.3	0.1	0.2	0.6	0.6	0.9	0.3	0.1	0.0	0.0	-	-	0.2
1925	0.1	0.0	0.0	0.1	0.2	0.4	0.4	0.1	0.0	-	0.0	-	0.0
1924	0.5	1.8	1.5	0.8	1.0	0.4	0.2	0.2	0.1	0.0	-	-	-
1923	0.7	0.5	1.7	1.2	0.8	1.0	1.2	0.5	0.1	0.1	0.0	0.2	1.1
1922	0.2	0.2	0.4	0.2	0.5	0.4	0.5	0.1	0.0	0.0	-	0.0	0.1
1921	0.4	0.1	0.4	0.2	0.8	0.8	0.5	0.4	0.0	0.0	-	0.0	0.1
1920	1.3	1.2	1.8	2.6	3.5	3.4	2.0	0.8	0.2	0.1	0.0	-	0.0

See footnotes at end of table.

TABLE 16. - 1/ DEATH RATES FOR SELECTED CAUSES BY MONTH: JAPAN, 1920 - 1949 - Cont'd
MEASLES (35)

Year	Annual	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
*1949	15.1	8.2	10.8	20.1	26.4	39.5	32.7	20.7	9.7	3.4	2.4	3.4	3.9
1948	7.0	6.0	6.5	10.4	8.5	10.8	11.3	9.0	5.5	2.8	2.4	4.2	6.3
1947	26.8	18.8	35.6	45.8	48.9	50.6	48.1	33.3	17.3	6.5	4.5	5.6	7.8
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	29.4	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	9.2	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1941	20.0	7.2	12.3	24.4	36.2	52.5	44.7	25.7	11.3	5.4	5.3	6.2	8.1
1940	7.1	6.2	5.7	8.0	8.9	12.8	12.9	11.6	5.8	2.6	2.5	3.2	4.5
1939	20.5	8.1	16.8	31.0	41.0	51.1	44.2	27.2	10.4	4.2	2.8	4.2	5.5
1938	7.0	5.3	5.7	7.4	8.7	15.3	14.5	10.6	4.7	2.1	2.0	3.2	4.6
1937	15.3	8.1	12.7	19.7	30.6	36.6	30.0	22.6	9.2	3.1	3.0	3.6	4.7
1936	7.5	5.3	5.8	7.2	9.5	14.4	16.6	10.6	5.5	2.2	2.3	4.2	6.3
1935	14.2	11.3	15.5	23.8	28.0	29.8	25.3	14.7	6.9	2.6	3.2	3.9	5.3
1934	14.3	10.9	13.7	21.7	26.6	32.6	24.6	16.1	7.3	3.2	2.8	5.1	7.3
1933	12.5	6.7	8.2	17.0	23.0	30.3	24.4	16.1	6.9	2.5	2.3	5.2	7.4
1932	9.4	10.2	11.3	15.3	15.8	18.9	13.8	11.2	5.1	2.1	1.5	2.5	4.7
1931	20.9	13.0	18.3	25.3	34.5	46.0	48.0	26.3	14.6	5.4	4.0	6.4	9.6
1930	9.3	9.5	9.1	11.2	13.0	16.0	16.8	12.3	6.4	2.1	3.0	4.5	7.2
1929	25.3	25.8	36.3	50.0	45.3	48.4	40.9	24.6	10.7	4.2	4.1	6.8	7.6
1928	17.7	17.1	19.4	23.6	22.5	30.0	30.5	23.0	11.0	5.4	4.5	9.6	16.2
1927	21.8	14.3	18.1	26.7	31.8	44.2	42.3	31.3	15.1	6.2	6.6	10.5	14.7
1926	17.3	12.0	13.6	15.7	23.3	35.5	35.7	25.7	13.4	5.6	5.2	9.2	12.3
1925	25.8	23.8	29.2	38.5	46.6	54.4	47.1	27.6	13.7	6.3	5.0	8.2	10.1
1924	20.3	13.2	16.6	25.1	31.0	39.8	36.5	28.5	13.7	6.2	6.1	11.4	16.1
1923	22.8	16.9	17.7	23.4	32.7	45.7	47.2	33.9	20.6	8.0	7.4	8.9	11.0
1922	16.8	21.1	18.1	19.0	20.1	28.4	29.7	21.1	12.4	4.5	4.4	8.3	14.9
1921	24.5	15.7	19.9	25.6	34.0	44.9	44.0	37.5	23.3	9.7	8.8	14.7	17.8
1920	13.4	20.5	18.5	15.2	16.3	17.7	20.1	15.4	8.6	4.5	4.3	7.5	12.5

See footnotes at end of table.

TABLE 16. - 1/ DEATH RATES FOR SELECTED CAUSES BY MONTH: JAPAN, 1920 - 1949 - Cont'd
ACUTE POLIOMYELITIS AND ACUTE POLIOENCEPHALITIS (36)

Year	Annual	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
*1949	1.2	1.1	1.0	1.1	1.0	1.2	1.3	1.6	1.2	1.6	1.6	1.2	1.2
1948	1.0	0.9	0.8	0.9	0.8	0.7	0.7	1.2	1.4	1.3	1.1	0.8	1.0
1947	1.3	1.0	1.5	1.4	1.1	1.4	1.3	1.9	1.9	1.2	0.9	0.8	1.0
1920 - 1946	Not Available.												

JAPANESE "B" ENCEPHALITIS (ENCEPHALITIS JAPONICA) (37b)

Year	Annual	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
* 1949	1.4	0.0	0.1	0.1	0.0	0.1	0.2	0.3	2.2	11.5	2.4	0.3	0.0
1948	3.3	0.0	0.0	0.0	0.0	-	0.1	0.4	24.4	12.0	1.5	0.4	0.2
1947	0.3	0.0	0.1	0.0	0.1	0.1	0.1	0.2	0.8	1.3	0.6	0.0	-
1920 - 1946	Not Available.												

See footnotes at end of table.

TABLE 16. - 1/ DEATH RATES FOR SELECTED CAUSES BY MONTH: JAPAN, 1930 - 1949 - Cont'd
YELLOW FEVER (38a)

Year	Annual	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
*1949	-	-	-	-	-	-	-	-	-	-	-	-	-
1948	-	-	-	-	-	-	-	-	-	-	-	-	-
1947	-	-	-	-	-	-	-	-	-	-	-	-	-
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	0.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	0.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1941	-	-	-	-	-	-	-	-	-	-	-	-	-
1940	-	-	-	-	-	-	-	-	-	-	-	-	-
1939	-	-	-	-	-	-	-	-	-	-	-	-	-
1938	-	-	-	-	-	-	-	-	-	-	-	-	-
1937	0.0	-	-	-	-	-	-	-	-	-	-	0.0	-
1936	-	-	-	-	-	-	-	-	-	-	-	-	-
1935	-	-	-	-	-	-	-	-	-	-	-	-	-
1934	-	-	-	-	-	-	-	-	-	-	-	-	-
1933	0.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1932	-	-	-	-	-	-	-	-	-	-	-	-	-
1931	0.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1930	0.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1929	0.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1928	-	-	-	-	-	-	-	-	-	-	-	-	-
1927	0.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1926	-	-	-	-	-	-	-	-	-	-	-	-	-
1925	-	-	-	-	-	-	-	-	-	-	-	-	-
1924	0.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1923	0.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1922	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1921	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1920	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA

See footnotes at end of table.

TABLE 16. - 1/ DEATH RATES FOR SELECTED CAUSES BY MONTH: JAPAN, 1920 - 1949 - Cont'd
RABIES (38b)

Year	Annual	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
* 1949	0.1	0.0	0.1	0.0	0.1	0.1	0.1	0.1	0.2	0.1	0.1	0.1	0.1
1948	0.0	0.1	0.1	-	0.0	0.0	0.0	0.0	0.1	0.1	0.0	0.1	0.1
1947	0.0	0.0	0.0	-	0.0	0.0	-	0.0	-	-	-	0.0	0.0
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	0.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	0.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1941	0.0	-	-	-	-	0.0	0.0	0.0	0.0	0.0	-	-	-
1940	0.0	-	-	0.0	-	-	-	0.0	0.0	-	0.0	0.0	-
1939	0.0	-	-	0.0	-	-	-	-	-	-	-	-	0.0
1938	0.0	-	-	-	0.0	0.0	-	-	-	-	-	-	-
1937	0.0	-	-	-	0.0	0.0	-	0.0	0.0	-	-	-	-
1936	0.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1935	0.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1934	0.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1933	0.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1932	0.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1931	0.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1930	0.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1929	0.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1928	0.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1927	0.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1926	0.1	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1925	0.2	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1924	0.4	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1923	0.3	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1922	0.1	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1921	0.1	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1920	0.1	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA

See footnote at end of table.

TABLE 16. - 1/ DEATH RATES FOR SELECTED CAUSES BY MONTH: JAPAN, 1920 - 1949 - Cont'd
2/ TYPHUS FEVER AND TYPHUS LIKE DISEASES (39)

Year	Annual	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
* 1949	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.0	0.1	0.1	0.0	0.0	0.0
1948	0.1	0.2	0.2	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
1947	0.2	0.4	0.2	0.3	0.4	0.2	0.2	0.2	0.2	0.1	0.1	0.0	0.3
1946	3.9	0.6	2.7	10.2	12.5	10.2	5.4	2.0	0.5	0.2	1.4	0.2	0.3
1945	0.4	0.1	0.2	0.2	0.6	1.1	1.0	0.3	0.4	0.1	0.1	0.1	0.3
1944	0.9	0.8	1.0	1.2	1.1	1.9	2.1	0.8	0.8	0.3	0.1	0.0	0.0
1943	0.2	0.0	0.0	0.0	0.1	0.2	0.6	0.1	0.0	0.1	0.2	0.3	0.4
1942	0.1	0.0	0.0	-	0.1	0.0	0.1	0.0	-	0.0	-	-	0.0
1941	0.1	0.0	-	0.0	0.0	0.0	0.1	0.1	0.2	0.1	-	0.0	0.0
1940	0.1	-	-	-	0.0	0.0	0.0	0.2	0.3	0.1	0.0	0.0	0.0
1939	0.1	-	-	-	-	-	-	0.1	0.2	0.2	0.1	0.0	0.0
1938	0.1	-	-	-	-	-	-	0.2	0.3	0.1	0.1	-	-
1937	0.0	-	-	-	-	-	0.0	0.1	0.2	0.1	0.1	0.0	-
1936	0.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1935	0.1	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1934	0.1	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1933	0.1	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1932	0.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1931	0.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1930	0.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1929	0.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1928	0.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1927	0.1	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1926	0.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1925	0.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1924	0.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1923	0.1	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1922	0.1	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1921	0.1	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1920	0.2	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA

See footnotes at end of table.

TABLE 16. - 1/ DEATH RATES FOR SELECTED CAUSES BY MONTH: JAPAN, 1920 - 1949 - Cont'd
CANCER AND OTHER MALIGNANT TUMORS (45-55)

Year	Annual	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
* 1949	71.6	67.9	67.2	67.9	68.8	70.3	71.6	73.6	76.2	76.3	73.4	72.8	73.4
1948	69.4	63.2	62.9	64.4	66.5	67.5	70.6	72.4	71.9	75.2	74.7	72.9	70.8
1947	68.0	65.5	63.0	64.4	65.3	64.2	69.2	70.9	71.1	72.5	72.8	69.6	67.3
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	70.9	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	71.7	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1941	71.0	66.2	66.2	66.6	66.9	68.5	72.7	75.7	75.8	78.1	74.0	69.6	71.1
1940	70.6	67.5	64.6	65.4	67.0	66.6	70.6	77.6	79.2	76.6	71.5	71.4	69.4
1939	71.5	67.8	65.4	68.7	66.8	70.2	73.2	77.6	75.7	76.9	74.0	71.2	70.1
1938	70.9	65.7	66.6	64.1	66.5	67.4	72.4	76.0	78.6	78.3	75.5	71.7	67.2
1937	71.3	66.4	67.0	65.6	67.5	71.0	73.2	74.8	76.2	77.4	76.0	71.7	68.9
1936	70.1	69.1	64.3	63.1	64.4	68.2	70.1	73.8	75.6	76.3	76.4	72.7	67.0
1935	70.9	65.0	66.3	62.8	67.0	69.6	73.0	76.6	76.6	75.9	76.9	72.0	68.8
1934	70.2	66.4	64.0	62.5	65.2	67.3	72.8	75.6	76.0	78.5	76.4	70.8	66.9
1933	69.6	63.2	65.3	63.5	65.2	69.9	71.6	74.0	76.3	77.5	76.4	67.1	64.4
1932	68.2	60.9	60.5	64.0	64.1	65.3	69.6	72.3	72.3	76.7	73.1	70.1	66.7
1931	68.1	63.7	65.0	64.8	65.5	67.2	68.6	69.4	73.1	73.4	73.1	68.3	65.0
1930	69.7	64.9	64.3	65.0	66.0	68.7	71.2	75.4	76.0	77.2	72.4	68.7	65.7
1929	68.9	64.0	62.5	63.2	67.5	66.3	71.9	73.4	75.9	74.6	72.7	67.9	66.2
1928	71.0	64.6	62.5	65.8	66.9	68.9	71.8	76.1	78.1	78.3	72.7	72.4	66.5
1927	69.4	62.7	65.4	64.9	65.6	69.0	72.0	74.5	76.0	76.5	72.5	69.3	65.7
1926	69.0	65.0	62.7	63.3	66.4	69.2	71.3	75.0	78.1	78.8	74.7	68.6	67.0
1925	70.9	62.3	64.3	62.5	65.6	70.0	72.1	75.3	77.2	76.3	77.1	69.2	66.2
1924	70.1	62.4	64.1	65.0	65.3	68.7	72.1	78.3	78.3	79.0	74.5	68.8	65.0
1923	71.8	64.4	62.8	64.8	68.4	72.1	76.7	78.1	79.9	82.3	79.1	68.5	64.5
1922	71.1	63.5	63.5	66.1	65.4	70.9	72.3	76.9	79.9	80.2	78.8	71.1	64.9
1921	71.6	63.4	63.0	64.5	65.8	71.9	70.1	76.9	82.3	82.0	80.9	72.2	65.1
1920	72.1	66.1	64.4	64.5	65.9	70.8	72.9	75.5	84.7	85.7	77.4	69.9	66.7

See footnotes at end of table.

TABLE 16. - 1/ DEATH RATE FOR SELECTED CAUSES BY MONTH: JAPAN, 1920 - 1949 - Cont'd
DIABETES MELLITUS (61)

Year	Annual	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
* 1949	2.3	2.9	2.8	2.9	2.6	2.2	1.9	2.0	1.8	1.9	2.2	2.1	2.3
1948	2.2	3.2	3.1	2.1	2.6	2.0	1.8	2.0	2.0	1.8	1.9	2.0	2.3
1947	2.3	2.8	3.4	3.0	2.3	2.2	2.1	2.2	1.6	2.1	1.9	1.9	2.8
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	3.3	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	3.5	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1941	3.6	4.8	5.5	3.8	3.5	3.4	2.5	3.3	3.2	3.3	2.8	3.7	3.9
1940	3.8	4.4	5.1	4.4	4.2	3.8	3.5	2.9	3.3	3.5	3.3	3.4	4.2
1939	3.9	4.9	4.7	4.6	4.0	3.6	3.1	3.3	3.6	3.6	3.9	3.6	4.0
1938	4.3	5.5	4.8	5.2	4.3	4.0	3.7	3.7	3.8	3.6	4.0	4.1	4.7
1937	4.0	4.4	4.5	4.2	3.5	3.4	3.5	3.6	3.7	3.8	4.0	4.1	4.8
1936	3.7	5.2	5.5	4.7	3.9	3.2	3.2	3.1	3.1	3.4	3.5	3.2	3.0
1935	3.6	4.3	4.6	3.8	3.8	3.4	3.2	3.3	3.7	3.2	3.0	3.4	4.2
1934	4.0	5.1	5.3	4.9	4.3	3.7	3.5	3.3	3.2	3.2	3.7	3.8	3.7
1933	3.8	4.8	4.8	4.7	4.0	3.5	3.1	3.1	3.4	3.3	3.7	4.0	4.0
1932	3.5	3.8	4.2	3.7	4.1	2.9	3.3	3.1	3.1	3.6	3.4	3.2	3.7
1931	3.4	4.5	4.2	4.3	3.4	4.0	3.0	2.8	2.6	2.3	2.9	2.8	3.7
1930	3.5	4.7	4.2	3.8	3.4	3.6	3.2	3.1	2.6	3.1	3.2	3.5	3.6
1929	3.6	4.1	4.3	4.7	3.9	3.2	3.3	2.9	3.2	3.6	3.1	3.6	3.5
1928	3.5	4.0	4.1	3.9	4.0	3.2	3.2	3.0	2.7	3.0	3.2	3.0	4.3
1927	3.5	4.1	5.0	4.2	4.3	3.3	2.9	3.2	2.7	3.1	3.3	3.0	3.3
1926	3.4	3.6	3.9	3.7	3.4	3.1	3.1	3.3	2.9	3.4	3.3	3.4	3.4
1925	3.3	4.4	4.3	4.0	3.4	3.2	3.1	2.7	2.7	2.7	3.5	2.8	3.0
1924	3.3	3.7	3.4	3.7	3.5	3.0	3.1	3.2	3.1	3.0	3.4	3.3	3.5
1923	3.2	3.9	4.2	3.1	3.4	3.2	2.8	3.3	2.7	2.8	3.4	2.7	3.2
1922	3.3	4.1	4.5	3.6	3.3	2.9	3.3	2.4	2.7	3.1	3.0	3.3	3.6
1921	3.3	3.8	3.8	3.4	3.3	3.0	3.1	3.3	3.0	3.6	3.4	2.9	3.3
1920	3.1	3.6	4.1	3.5	3.1	2.5	2.9	2.6	2.7	2.8	2.6	3.4	3.4

See footnotes at end of table.

TABLE 16. - 1/ DEATH RATES FOR SELECTED BY MONTH: JAPAN, 1920 - 1949 - Cont'd
BERIBERI (68)

Year	Annual	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
*1949	6.8	8.6	8.5	8.6	8.2	6.8	6.1	5.3	5.9	5.3	5.5	5.8	7.1
1948	7.8	10.5	11.3	9.7	8.6	6.9	7.2	6.7	7.2	6.4	6.0	6.5	7.2
1947	11.0	12.2	12.8	13.5	12.8	11.3	9.9	10.2	9.6	8.8	9.1	9.6	12.5
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	8.7	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	9.1	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1941	9.1	11.1	10.7	10.8	9.3	7.9	7.3	7.9	7.3	8.8	8.2	9.2	11.1
1940	10.0	13.5	12.0	11.5	9.7	8.3	8.8	8.6	9.9	9.1	8.6	9.4	10.2
1939	13.4	16.1	14.3	12.3	11.3	10.1	9.9	12.0	15.1	15.5	15.4	14.0	14.6
1938	17.9	15.7	13.7	12.5	11.7	12.4	13.5	17.1	24.3	29.1	26.2	20.8	17.0
1937	15.0	12.0	12.0	11.0	10.5	9.8	10.2	12.3	17.4	20.9	24.0	21.4	18.0
1936	15.8	12.5	12.4	13.3	11.1	10.3	12.3	16.0	20.4	25.3	25.3	16.9	13.9
1935	14.5	11.2	10.8	10.9	9.5	8.3	9.3	12.2	17.3	23.0	27.8	18.9	15.0
1934	16.8	12.3	11.4	11.4	11.0	11.0	12.1	17.3	20.3	26.2	31.8	22.7	13.9
1933	20.5	14.1	12.3	12.0	9.9	13.5	15.2	23.1	27.6	41.1	37.7	23.0	16.3
1932	17.8	13.0	12.4	11.7	10.5	10.7	11.6	18.8	24.8	29.9	32.3	22.2	15.2
1931	27.2	11.7	12.5	14.4	16.2	20.8	18.7	30.1	44.6	53.0	54.9	29.8	18.0
1930	23.9	14.8	13.1	13.9	16.2	20.9	20.7	35.0	39.9	45.9	33.8	19.1	13.0
1929	24.3	15.5	14.5	13.0	14.6	17.7	22.9	34.0	41.7	37.5	35.4	26.9	17.6
1928	30.4	11.2	10.5	9.8	11.3	13.3	17.0	31.3	54.4	76.9	65.1	41.2	22.1
1927	19.5	10.8	9.4	9.3	9.9	12.3	15.3	24.5	31.8	36.1	36.7	23.3	14.3
1926	19.9	13.1	10.5	12.2	10.6	13.0	17.8	22.6	32.7	37.3	36.1	19.7	12.5
1925	23.3	13.5	12.2	11.1	11.8	15.5	20.8	31.3	35.4	43.0	42.9	25.0	16.0
1924	31.1	15.8	14.6	15.3	13.7	20.0	24.8	45.8	59.3	61.9	51.6	31.4	18.5
1923	46.0	12.8	12.6	13.8	19.5	27.3	32.2	57.0	103.2	118.5	86.9	43.2	22.7
1922	33.4	13.4	10.8	12.9	13.9	21.8	25.7	43.0	67.2	63.1	37.2	32.2	17.3
1921	40.0	8.1	8.0	7.4	7.4	13.3	18.2	45.4	94.8	112.1	95.2	47.4	20.3
1920	25.4	7.0	6.0	5.4	8.0	10.3	12.2	30.1	57.2	80.8	52.2	23.7	11.9

See footnotes at end of table.

TABLE 16. - 1/ DEATH RATES FOR SELECTED CAUSES BY MONTH: JAPAN, 1920 - 1949 - Cont'd
g/ SIMPLE MENINGITIS (81a)

Year	Annual	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
* 1919	11.1	11.0	11.2	12.5	13.6	12.7	11.2	10.9	11.0	11.1	9.2	9.1	9.4
** 1948	13.0	14.5	14.0	14.8	16.2	15.3	14.1	14.5	12.7	9.9	9.7	9.9	10.3
1947	18.1	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1941	41.0	34.1	37.6	40.1	41.6	44.4	45.9	53.8	50.2	40.1	36.4	33.8	33.6
1940	43.8	38.7	41.4	45.1	45.2	47.9	49.2	59.4	53.4	43.7	36.3	33.2	32.1
1939	49.8	39.8	42.8	49.3	49.0	49.8	55.7	61.5	74.7	57.3	43.9	38.1	35.5
1938	51.6	44.1	47.3	49.6	51.2	52.2	54.5	67.2	69.6	62.4	43.9	39.4	37.8
1937	51.0	40.0	42.1	46.2	50.0	51.2	52.7	65.7	81.0	57.7	45.3	39.8	39.1
1936	52.6	43.7	47.7	52.0	55.2	52.5	58.2	65.8	69.4	61.6	46.0	38.2	40.3
1935	54.3	45.6	47.1	52.0	54.2	54.7	58.1	69.5	84.4	56.4	47.1	40.8	41.0
1934	58.4	50.6	51.7	57.9	58.5	62.4	65.7	77.8	72.7	61.6	50.4	46.2	45.2
1933	63.1	53.0	55.3	59.2	61.3	66.7	68.4	85.4	90.5	67.4	55.3	48.0	45.9
1932	65.6	54.8	58.6	63.2	61.5	66.1	70.5	89.1	94.6	72.5	55.8	49.5	50.8
1931	70.8	56.5	63.1	65.6	69.0	73.3	75.7	87.3	97.7	84.6	66.7	56.9	52.3
1930	73.8	62.4	65.0	68.0	70.3	71.3	81.2	99.6	114.6	80.4	62.7	54.7	53.7
1929	85.2	67.0	71.2	80.3	80.2	79.8	91.0	118.8	143.7	93.3	72.8	63.8	58.5
1928	85.1	69.8	72.4	78.3	79.2	83.4	88.3	115.4	115.9	108.3	79.1	65.9	65.0
1927	89.8	68.3	75.4	79.4	82.4	85.9	91.5	125.7	147.0	110.8	79.6	64.5	65.5
1926	93.4	72.9	74.4	81.2	87.5	94.1	95.8	116.1	148.8	127.3	82.9	71.2	66.5
1925	98.0	81.0	88.7	92.7	95.4	99.1	106.6	126.6	133.5	120.0	88.5	72.7	70.4
1924	116.2	91.8	94.7	101.9	104.2	104.8	117.2	164.1	243.6	127.4	88.9	76.8	77.1
1923	126.3	93.0	94.9	104.9	107.7	109.0	125.4	164.0	213.0	197.0	125.0	91.6	87.2
1922	121.2	93.7	107.4	105.4	101.8	105.4	115.6	152.4	228.5	172.5	106.9	83.9	79.7
1921	120.4	87.0	98.2	103.0	106.9	105.0	113.5	155.8	199.5	174.3	116.1	93.8	90.2
1920	123.7	105.8	113.2	118.5	109.3	109.8	115.6	165.7	208.4	160.6	100.0	87.5	88.3

See footnotes at end of table.

TABLE 16. - 1/ DEATH RATES FOR SELECTED CAUSES BY MONTH: JAPAN, 1920 - 1949 - Cont'd
2/ INTRACRANIAL LESIONS OF VASCULAR ORIGIN (83)

Year	Annual	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
* 1949	122.4	137.2	133.9	140.5	133.3	113.8	108.9	102.2	96.3	114.2	122.0	129.6	138.4
1948	117.6	148.1	148.2	139.2	119.5	109.9	101.7	90.5	95.7	106.6	106.8	119.9	126.3
1947	129.6	163.1	181.4	161.7	136.3	121.2	110.0	98.8	90.8	101.5	116.9	128.1	149.0
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	162.8	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	169.8	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1941	170.9	195.9	203.1	186.1	177.3	160.7	153.7	149.6	145.5	163.5	166.1	168.9	183.3
1940	177.3	219.3	224.7	207.5	183.9	170.7	154.8	143.4	153.2	162.8	162.9	164.5	181.5
1939	183.0	217.6	207.7	203.4	180.8	168.7	159.7	151.1	174.7	177.9	179.2	176.4	199.7
1938	178.2	214.8	216.2	188.0	172.6	157.3	153.4	152.8	156.7	189.8	172.2	179.4	188.1
1937	167.3	182.9	176.2	178.3	163.7	154.1	147.8	135.2	166.9	166.5	170.6	166.9	198.2
1936	168.3	212.5	204.8	194.0	172.2	153.8	144.5	137.8	137.1	161.1	167.1	163.7	172.4
1935	165.4	185.9	184.6	176.4	161.6	152.5	146.0	138.0	172.2	163.1	158.2	163.2	184.2
1934	167.6	207.2	204.0	193.4	174.7	151.9	145.3	133.9	138.6	149.8	169.6	170.8	174.1
1933	164.3	195.6	197.5	189.3	162.9	150.3	145.7	130.6	150.7	153.7	158.7	160.9	177.3
1932	161.7	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1931	164.1	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1930	162.8	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1929	170.8	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1928	164.3	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1927	164.8	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1926	162.3	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1925	161.2	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1924	174.5	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1923	162.6	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1922	159.3	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1921	159.7	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1920	157.6	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA

See footnotes at end of table.

TABLE 16. --1/ DEATH RATES FOR SELECTED CAUSES BY MONTH: JAPAN, 1920 - 1949 - Cont'd
10/ DISEASES OF THE HEART (90-95)

Year	Annual	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
*1949	64.3	73.2	72.5	74.9	72.2	60.0	58.5	55.3	52.7	54.1	60.5	65.2	72.6
1948	63.2	70.5	74.6	72.3	62.9	57.9	57.0	54.0	50.7	52.7	56.2	62.0	63.6
1947	62.3	69.5	72.6	73.7	63.3	61.6	57.5	53.3	50.9	49.9	57.0	60.4	72.4
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	61.1	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	58.9	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1941	58.1	74.9	77.1	65.7	59.8	54.0	50.6	45.3	45.7	50.4	52.5	55.6	63.2
1940	63.2	83.9	85.6	75.2	67.0	59.6	57.1	53.0	49.5	52.2	54.1	55.4	66.0
1939	66.4	89.0	83.5	85.8	67.2	59.3	56.8	55.6	53.4	53.4	58.3	60.8	74.0
1938	66.7	86.4	90.7	75.4	64.4	58.6	55.1	57.3	55.4	58.3	61.7	66.3	72.0
1937	60.3	71.5	68.6	63.8	61.6	56.5	53.8	52.0	50.3	53.8	58.8	59.3	74.0
1936	61.1	87.1	84.6	70.5	58.9	53.8	54.3	51.6	48.0	50.3	56.6	57.0	61.5
1935	57.6	65.8	69.4	62.8	56.3	53.0	54.2	48.5	48.0	52.1	56.6	57.4	68.0
1934	62.3	81.6	81.0	74.4	63.5	56.2	52.7	50.5	49.7	52.2	61.3	61.4	63.8
1933	59.5	70.4	73.3	72.2	60.5	54.5	53.0	50.0	47.2	53.8	56.6	59.3	64.2
1932	58.7	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1931	64.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1930	63.8	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1929	65.4	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1928	63.6	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1927	63.2	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1926	62.4	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1925	66.8	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1924	69.1	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1923	72.5	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1922	65.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1921	66.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1920	63.5	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA

See footnotes at end of table.

TABLE 16. - 1/ DEATH RATES FOR SELECTED CAUSES BY MONTH: JAPAN, 1920 - 1949 - Cont'd
11/ BRONCHITIS (106)

Year	Annual	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
1949	31.3	44.5	48.5	42.3	42.1	30.8	25.0	19.9	16.3	16.9	21.1	25.9	37.0
1948	32.3	54.9	48.4	48.4	37.5	28.9	24.1	18.6	16.7	20.7	23.6	26.5	34.6
1947	44.8	48.7	67.8	81.1	63.3	46.4	38.2	30.6	25.7	22.4	27.9	36.7	50.0
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	31.3	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	30.4	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1941	30.1	41.8	54.6	44.7	34.7	27.4	23.7	20.9	17.3	19.6	22.2	25.2	31.0
1940	31.3	49.6	54.4	45.9	35.6	28.0	23.7	20.9	19.4	20.6	21.9	23.5	32.9
1939	35.1	51.8	54.9	56.0	44.9	29.7	26.9	22.6	21.4	21.8	25.3	28.6	38.9
1938	36.8	56.9	64.4	57.2	41.7	32.8	24.9	25.0	22.3	24.1	25.4	30.4	38.0
1937	33.8	43.8	47.4	45.1	37.8	31.9	26.6	23.4	22.2	23.8	27.9	30.9	46.1
1936	37.2	63.6	70.7	57.8	41.6	31.6	26.8	25.1	19.8	21.6	25.9	28.6	34.6
1935	35.0	50.1	53.7	47.7	40.0	32.3	27.3	23.4	22.5	24.4	27.4	31.7	40.7
1934	39.6	61.1	66.1	65.7	47.6	35.4	28.2	24.3	21.4	24.1	29.0	32.6	40.9
1933	37.7	51.3	63.7	57.8	45.7	34.0	30.2	23.8	21.6	24.0	28.9	32.8	40.4
1932	41.6	52.7	60.6	63.8	52.5	42.2	35.0	29.6	26.0	28.5	31.2	35.0	42.8
1931	46.4	67.4	78.1	68.6	55.0	44.5	38.1	32.9	28.4	29.2	33.4	38.7	45.1
1930	43.3	62.5	65.5	60.3	49.3	40.8	35.9	30.1	27.5	29.6	34.6	39.0	46.4
1929	50.2	75.6	78.2	78.0	62.3	48.5	38.6	34.9	29.0	33.9	35.4	42.0	48.0
1928	52.1	69.5	82.9	80.1	61.5	45.3	38.2	34.8	31.7	32.1	41.0	46.4	62.8
1927	54.0	73.6	90.2	84.6	61.9	52.5	43.7	36.7	33.1	36.6	39.5	43.7	54.1
1926	51.5	69.9	73.2	74.0	62.4	52.3	43.8	34.7	30.6	31.0	40.6	46.3	61.3
1925	59.5	86.7	121.2	100.3	69.0	52.0	43.6	37.9	30.7	32.3	42.3	44.4	57.4
1924	60.8	87.7	95.4	97.8	68.4	56.5	47.3	37.5	35.1	39.1	45.9	54.9	64.9
1923	62.2	90.7	97.1	87.4	69.8	52.8	47.9	42.6	37.5	44.1	53.5	54.9	71.0
1922	90.2	150.4	168.7	156.6	101.6	74.2	60.5	49.2	45.3	51.1	61.3	70.6	97.7
1921	91.8	122.5	134.3	132.7	104.3	81.4	69.8	58.0	54.1	62.1	73.4	96.4	115.7
1920	94.1	153.5	168.7	146.5	109.5	82.4	65.4	56.9	50.2	56.8	64.3	72.6	104.8

See footnotes at end of table.

TABLE 16. - 1/ DEATH RATES FOR SELECTED CAUSES BY MONTH: JAPAN, 1920 - 1949 - Cont'd
12/ PNEUMONIA, ALL FORMS (107-109)

Year	Annual	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
*1949	68.9	107.8	108.1	104.2	94.4	70.4	51.1	40.5	29.0	33.3	38.6	59.8	92.1
1948	66.1	118.6	120.4	108.6	80.9	56.7	46.5	33.9	31.0	31.3	37.5	51.8	76.1
1947	130.3	235.7	317.5	243.8	181.5	129.4	94.5	62.5	43.2	40.7	47.5	70.0	113.5
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	156.8	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	143.7	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1941	142.2	217.2	288.4	252.9	190.1	149.9	111.1	79.8	63.2	60.4	70.2	93.5	139.9
1940	154.0	272.7	288.9	275.2	223.6	165.4	122.5	79.1	61.4	60.3	71.0	85.1	137.6
1939	184.0	264.0	318.4	375.7	275.6	189.3	155.0	97.9	75.5	68.5	83.1	112.0	199.6
1938	165.9	292.2	347.0	298.2	210.6	156.4	110.6	83.6	64.8	68.5	81.0	110.7	179.9
1937	182.5	217.4	247.4	240.5	217.9	167.6	134.2	95.6	71.8	73.3	86.1	104.1	180.5
1936	199.8	305.0	348.3	265.7	196.0	146.5	115.4	84.5	64.9	63.5	80.6	103.8	150.7
1935	151.7	229.2	253.9	242.2	204.4	163.1	130.3	88.8	73.2	75.4	80.9	110.2	176.1
1934	181.7	311.5	349.9	345.6	250.7	190.8	133.4	86.1	69.7	68.5	90.7	116.5	178.5
1933	197.6	225.8	271.3	283.4	216.5	169.0	128.5	83.1	68.4	69.8	91.3	115.9	176.5
1932	169.7	229.7	267.1	306.5	281.5	202.9	138.0	98.5	74.5	73.1	88.1	111.0	169.4
1931	197.8	296.8	379.6	366.2	276.3	218.1	172.8	109.7	85.4	77.0	94.2	127.5	183.5
1930	156.8	234.6	260.0	248.5	217.9	172.7	130.8	91.0	75.2	77.7	91.3	115.3	173.7
1929	190.8	272.6	306.3	353.5	289.1	220.5	157.7	110.5	86.3	86.8	96.9	129.1	178.9
1928	200.6	339.5	391.5	391.5	303.4	197.7	140.2	108.3	93.2	86.5	111.2	147.6	217.3
1927	194.4	281.6	333.3	325.7	253.6	228.2	172.8	117.9	95.3	95.2	105.1	135.5	198.4
1926	177.8	241.4	264.5	273.7	255.0	210.0	159.2	113.1	84.6	82.0	107.2	138.2	210.7
1925	216.2	323.8	492.3	417.8	277.1	212.6	170.1	116.8	85.6	85.9	107.8	135.2	189.4
1924	209.5	298.0	327.1	364.0	292.4	233.0	186.5	131.2	110.9	100.6	111.7	148.0	214.8
1923	204.9	296.5	326.0	323.4	258.4	212.4	176.0	141.5	113.9	117.7	132.6	152.7	215.8
1922	196.2	292.5	428.6	383.7	232.8	170.9	138.5	101.5	96.0	93.7	109.2	132.8	191.0
1921	200.5	245.5	286.0	319.8	279.9	212.2	174.2	132.8	113.2	118.5	134.5	174.0	222.5
1920	313.9	940.8	691.8	532.0	395.3	252.7	160.1	123.3	111.1	107.4	118.8	144.5	204.8

See footnotes at end of table.

TABLE 16. - 1/ DEATH RATES FOR SELECTED CAUSES BY MONTH: JAPAN, 1920 - 1949 - Cont'd
PLEURISY (110)

Year	Annual	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
* 1949	7.3	7.4	8.0	8.8	8.1	7.7	7.7	7.7	7.2	6.8	6.7	6.2	5.7
** 1948	5.6	11.0	10.9	11.1	11.0	10.5	10.1	9.0	9.2	9.1	7.7	7.9	7.9
1947	4.3	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	24.1	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	24.5	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1941	24.7	23.6	24.0	24.8	27.1	26.6	26.9	24.8	24.5	25.3	24.0	22.4	22.5
1940	27.8	29.3	30.3	30.9	31.5	29.7	28.2	29.2	26.4	26.4	24.6	23.5	23.9
1939	30.3	28.1	28.1	32.6	33.4	32.2	33.1	31.8	30.3	29.7	28.6	27.9	28.0
1938	29.5	28.0	29.3	28.8	33.0	30.7	31.2	30.1	30.3	28.3	28.4	27.7	26.8
1937	26.5	22.5	26.4	26.1	26.3	29.7	28.1	27.1	28.7	29.3	27.0	26.0	26.2
1936	25.7	25.5	25.0	27.7	28.7	27.9	27.1	26.4	26.3	24.5	23.8	23.0	23.6
1935	24.6	23.9	24.8	27.9	25.9	26.0	26.6	24.4	24.0	24.1	24.5	23.1	21.9
1934	26.1	25.3	26.4	28.9	29.1	28.9	28.0	26.4	23.8	25.2	24.3	23.5	23.3
1933	26.1	23.9	24.7	27.6	28.3	28.9	28.4	27.0	24.7	24.7	24.6	24.5	23.6
1932	26.4	26.9	26.9	28.3	29.5	30.2	27.1	26.7	25.6	26.2	24.8	24.4	22.7
1931	27.0	24.5	26.2	28.8	30.5	29.7	28.8	27.7	26.4	26.2	26.6	23.0	25.1
1930	25.7	26.2	25.5	28.5	28.1	27.1	26.2	26.5	25.4	26.4	24.4	21.2	23.2
1929	27.1	25.3	27.0	28.3	28.8	29.1	28.8	28.9	26.5	25.8	25.2	24.8	23.5
1928	25.7	23.3	24.4	27.1	27.7	25.9	26.3	26.8	26.3	26.5	26.0	24.3	24.3
1927	25.9	23.4	24.5	26.3	27.7	27.1	27.7	28.4	27.5	26.6	25.0	23.4	23.6
1926	25.0	22.3	24.9	26.9	26.8	27.4	25.9	25.1	26.0	24.8	25.2	22.6	21.6
1925	27.2	25.5	28.8	30.4	30.6	30.0	27.7	28.4	26.7	26.8	25.8	23.5	22.8
1924	28.4	27.5	28.7	30.1	30.7	29.3	30.0	29.5	28.7	29.4	25.9	26.1	24.6
1923	27.7	24.6	27.5	27.3	27.5	27.4	29.5	28.4	28.9	30.8	28.2	26.0	25.3
1922	28.1	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1921	26.3	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1920	25.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA

See footnotes at end of table.

TABLE 16. - 1/ DEATH RATES FOR SELECTED CAUSES BY MONTH: JAPAN, 1920 - 1949 - Cont'd
13/ULCER OF STOMACH OR DUODENUM (117)

Year	Annual	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
1919	23.7	26.3	25.2	26.3	25.0	23.3	21.8	21.0	19.4	19.9	23.1	26.0	27.6
1918	25.8	33.8	31.2	29.2	26.8	23.6	23.6	21.1	19.8	22.0	24.5	27.0	27.4
1917	33.2	36.0	38.3	36.9	33.8	31.6	30.4	30.2	26.6	28.5	32.7	35.8	38.1
1916	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1915	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1914	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1913	23.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1912	20.2	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1911	17.6	19.1	18.4	17.3	16.5	16.9	16.7	15.2	15.2	16.7	17.8	20.2	21.5
1910	17.9	22.0	20.8	19.0	17.8	16.3	16.8	14.9	14.7	15.6	18.0	19.2	20.2
1909	18.5	21.4	21.3	20.2	17.7	17.0	15.8	16.5	14.5	16.7	18.3	21.4	21.3
1908	18.7	21.2	20.2	18.0	17.5	16.9	16.6	16.0	16.4	18.1	20.0	22.2	20.7
1907	17.4	19.6	16.7	17.3	16.4	16.4	16.6	14.7	14.2	15.4	19.0	19.0	22.8
1906	17.1	20.3	19.2	18.2	17.2	15.4	14.0	15.1	14.1	14.7	18.4	19.9	18.3
1905	16.3	18.1	16.2	16.5	15.3	15.1	14.6	14.9	14.0	15.5	17.5	18.7	19.0
1904	16.6	18.3	18.3	18.4	16.0	14.2	14.5	14.6	13.4	15.1	17.9	19.9	18.7
1903	15.6	17.6	17.3	16.1	14.1	14.5	13.8	13.6	13.3	14.9	16.3	17.8	18.3
1902	15.1	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1901	14.8	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1900	14.3	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1909	13.7	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1908	13.7	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1907	13.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1906	18.9	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1905	13.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1904	12.5	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1903	12.3	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1902	11.1	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1901	11.8	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1900	11.2	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA

See footnotes at end of table.

TABLE 16. - 1/ DEATH RATES FOR SELECTED CAUSES BY MONTH: JAPAN, 1920 - 1949 - Cont'd
15/ DIARRHEA, ENTERITIS AND ULCERATION OF THE INFECTIONS, UNDER 2 YEARS OF AGE (119)

Year	Annual	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
* 1949	50.9	40.5	36.4	38.6	41.8	46.7	65.5	81.3	76.9	55.3	37.7	45.4	43.2
1948	60.4	44.1	44.0	44.5	44.0	48.7	69.8	116.9	104.6	67.5	43.1	51.2	44.8
1947	71.1	39.4	40.1	45.0	45.9	45.0	67.1	144.9	178.9	87.8	50.5	51.1	53.9
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	71.4	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	65.9	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1941	64.3	39.4	38.2	36.4	40.2	55.9	92.7	138.7	117.7	75.5	48.3	43.5	42.7
1940	68.2	45.3	43.6	41.3	40.7	48.9	78.1	149.5	141.8	86.4	52.3	45.3	43.5
1939	74.1	50.6	45.6	47.0	44.3	58.7	86.3	150.6	150.7	94.0	61.0	46.0	50.6
1938	82.1	60.3	53.6	49.5	50.4	65.5	104.4	156.7	155.3	109.8	71.0	54.5	51.2
1937	93.9	56.3	54.3	53.0	54.1	73.9	109.7	187.3	213.6	124.7	69.2	61.1	65.5
1936	100.5	64.8	60.4	61.8	63.6	74.8	111.8	204.9	209.4	138.8	87.9	63.0	61.2
1935	92.8	57.7	55.4	53.7	60.5	78.9	130.8	194.0	172.8	108.7	70.7	63.1	64.0
1934	109.5	67.5	61.4	64.6	67.2	83.5	149.3	250.7	214.5	132.8	88.3	66.4	62.8
1933	118.6	70.6	70.0	68.0	67.3	91.1	158.3	267.9	245.7	144.7	85.6	73.6	74.4
1932	124.1	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1931	127.6	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1930	132.9	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1929	147.2	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1928	138.7	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1927	147.6	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1926	143.6	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1925	142.1	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1924	148.7	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1923	160.1	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1922	152.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1921	155.2	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1920	141.7	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA

See footnotes at end of table.

TABLE 16. · 1/ DEATH RATES FOR SELECTED CAUSES BY MONTH: JAPAN, 1920 - 1949 - Cont'd
14/ DIARRHEA, ENTERITIS AND ITCERATION OF THE INTESTINES (119-120)

Year	Annual	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
* 1949	87.8	69.1	64.2	68.1	71.9	77.0	100.2	133.6	137.6	109.9	76.3	74.7	69.3
1948	103.8	79.7	85.4	90.3	87.4	88.6	109.1	172.0	169.0	122.2	82.9	83.8	73.9
1947	130.1	93.7	96.6	99.3	97.0	94.6	119.5	231.9	274.3	160.5	104.2	92.4	92.0
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	142.4	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	131.2	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1941	132.1	84.3	83.4	80.8	85.9	105.7	158.8	256.2	247.5	175.3	118.2	65.4	89.0
1940	150.3	101.8	95.7	95.6	95.5	111.0	158.2	284.4	312.2	202.7	132.3	102.2	93.1
1939	157.8	104.0	97.4	100.5	97.4	114.9	161.4	279.0	315.9	230.0	162.2	116.8	107.5
1938	164.3	113.1	107.0	98.6	102.1	122.0	182.2	298.2	314.1	238.7	164.9	121.4	102.9
1937	169.0	100.9	95.2	100.2	100.7	126.6	173.6	304.4	370.4	244.3	156.2	123.5	120.2
1936	172.6	113.4	110.9	112.6	112.2	126.6	173.7	314.3	346.5	254.1	172.3	120.3	108.3
1935	161.9	106.9	102.7	99.3	106.7	131.7	198.0	300.6	298.3	211.6	150.4	119.0	111.3
1934	187.2	118.6	111.8	118.2	120.3	140.0	220.5	374.8	362.1	251.7	177.4	130.5	112.4
1933	194.0	121.7	122.7	115.1	119.0	146.9	231.3	385.9	384.4	260.6	169.5	134.8	124.0
1932	206.9	132.8	130.0	126.7	126.3	151.5	239.0	380.9	421.3	299.4	186.2	141.3	130.1
1931	214.2	140.9	143.9	136.9	140.9	165.6	225.0	351.6	395.9	315.7	226.0	165.6	148.8
1930	221.2	151.1	144.4	140.8	144.2	176.5	247.0	400.0	438.8	295.0	197.2	164.4	147.0
1929	246.8	156.7	154.8	156.8	157.3	176.3	257.1	445.3	521.1	330.0	243.6	189.6	154.0
1928	233.4	149.4	151.3	147.2	141.5	173.2	244.6	400.0	428.7	355.4	253.5	186.9	169.0
1927	242.8	148.0	155.7	158.8	165.5	180.7	240.7	442.7	442.7	347.6	234.8	171.3	153.6
1926	232.2	155.2	150.9	152.2	159.0	185.7	267.0	366.0	445.5	337.4	229.0	168.6	156.7
1925	238.0	155.8	148.1	166.5	167.1	198.9	273.4	420.7	408.9	322.8	233.1	171.0	158.1
1924	257.6	182.6	175.1	179.8	170.1	202.4	256.4	433.9	484.6	370.9	262.7	195.6	170.4
1923	283.6	172.1	177.1	172.9	179.0	198.3	288.4	443.7	536.8	460.2	339.4	234.5	196.4
1922	262.1	176.3	171.0	176.8	171.6	212.9	300.4	457.0	528.0	378.7	252.1	198.2	170.4
1921	272.8	160.2	160.1	162.8	158.7	211.6	361.4	464.0	555.5	425.5	296.5	223.9	181.8
1920	253.9	145.1	170.5	164.1	155.6	175.2	250.9	465.1	506.5	375.0	248.6	186.6	172.7

See footnotes at end of table.

TABLE 16. - 1/ DEATH RATES FOR SELECTED CAUSES BY MONTH: JAPAN, 1920 - 1949 - Cont'd
16/ DIARRHEA, ENTERITIS AND ULCERATIONS OF THE INTESTINES, 2 YEAR OF AGE AND OVER (120)

Year	Annual	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
1949	36.9	28.6	27.6	29.5	30.1	30.3	34.7	52.3	60.6	54.6	38.6	29.2	26.1
1948	43.4	35.6	41.4	45.8	43.4	39.9	39.3	55.0	64.3	54.7	39.7	32.6	29.0
1947	56.9	54.3	56.5	54.3	51.1	45.6	52.4	87.1	95.4	72.7	53.6	41.3	38.1
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	71.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	65.2	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1941	67.8	44.9	45.2	44.3	45.6	49.8	66.1	117.5	129.7	99.7	70.0	51.9	46.3
1940	82.0	56.5	56.1	54.3	54.8	62.1	80.1	144.9	170.4	116.3	80.0	56.9	49.7
1939	83.7	53.4	51.8	53.6	53.0	56.2	75.2	128.4	165.2	136.0	101.1	70.8	56.9
1938	82.2	52.8	53.5	49.1	51.7	56.5	77.7	141.5	158.8	128.8	93.8	66.9	51.8
1937	75.1	44.6	44.9	47.2	46.6	52.7	63.9	117.1	156.8	119.6	87.0	62.4	54.7
1936	72.1	48.7	50.5	50.8	48.6	51.8	61.9	109.3	137.1	115.3	84.4	57.3	47.6
1935	69.0	49.2	47.3	45.6	46.2	52.9	67.2	106.6	125.5	102.9	79.7	55.9	47.2
1934	77.7	51.1	50.4	53.6	53.0	56.5	71.2	124.1	147.6	118.9	89.1	64.1	49.6
1933	75.4	51.1	52.8	51.1	51.7	55.8	73.1	118.0	136.8	115.9	83.9	61.2	49.6
1932	82.7	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1931	86.5	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1930	88.3	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1929	99.6	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1928	94.7	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1927	95.1	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1926	88.7	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1925	95.9	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1924	108.9	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1923	123.6	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1922	115.1	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1921	117.6	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1920	122.2	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA

See footnotes at end of table.

TABLE 16. - 1/ DEATH RATES FOR SELECTED CAUSES BY MONTH: JAPAN, 1920 - 1949 - Cont'd
APPENDICITIS (121)

Year	Annual	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
1949	4.0	3.8	3.7	4.1	4.0	4.2	3.8	4.8	5.0	4.5	3.4	3.4	3.1
1948	4.7	4.1	5.3	4.9	5.2	4.3	5.3	5.3	5.8	5.4	4.3	3.9	3.4
1947	5.7	4.5	4.4	4.3	5.1	5.3	6.1	7.7	8.0	6.7	6.1	4.7	5.0
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	3.1	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	3.2	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1941	3.1	2.7	3.1	2.8	2.8	3.0	3.1	3.3	3.7	3.2	3.0	2.7	3.1
1940	3.6	2.8	3.4	3.6	3.9	3.2	3.8	3.9	4.6	3.8	3.4	3.3	3.1
1939	3.8	3.7	3.1	3.4	3.7	3.5	3.9	4.3	4.4	4.6	3.9	3.4	3.4
1938	3.8	3.4	3.2	3.7	3.4	3.7	3.4	4.1	4.7	4.0	4.5	3.7	3.2
1937	3.8	3.8	3.1	3.4	3.3	3.7	3.9	4.0	4.7	4.7	3.9	3.3	3.6
1936	3.6	3.7	2.9	3.4	3.6	3.1	3.5	4.1	4.5	4.2	4.0	3.6	2.9
1935	3.5	2.9	3.2	2.9	3.3	3.2	3.7	3.7	4.1	4.0	4.0	3.6	3.7
1934	3.8	3.5	3.2	3.6	3.6	3.4	3.8	4.1	5.1	4.7	3.8	3.7	3.2
1933	3.8	3.5	3.0	3.4	3.2	3.5	3.7	4.3	5.2	5.1	3.7	3.2	3.2
1932	4.0	3.7	3.4	3.5	3.4	3.7	3.9	4.3	5.3	4.6	4.4	4.0	3.6
1931	3.9	3.1	3.8	3.5	3.4	3.6	4.4	4.2	4.6	4.7	4.4	3.6	3.1
1930	4.0	3.7	3.8	3.9	3.6	3.8	3.8	4.6	5.2	5.0	4.3	3.4	3.2
1929	4.2	3.2	3.4	3.9	4.0	4.0	3.8	4.9	5.8	5.1	4.8	3.7	3.7
1928	4.1	3.5	3.1	3.8	3.6	3.7	4.0	4.8	5.0	5.2	4.3	4.1	4.1
1927	4.3	3.5	3.3	3.9	4.2	4.3	4.1	4.9	5.1	5.4	5.0	3.7	3.9
1926	4.2	4.4	3.5	3.7	3.9	3.4	4.1	4.7	5.3	5.4	4.9	4.0	3.1
1925	4.4	4.2	4.1	3.9	4.0	3.9	4.2	5.0	5.1	5.0	5.5	3.8	4.0
1924	4.6	4.1	3.8	3.9	3.7	3.7	4.3	5.0	6.0	5.7	5.4	5.2	4.1
1923	4.7	4.8	4.1	3.6	4.1	3.7	4.0	4.5	5.9	6.1	6.7	5.1	4.3
1922	4.9	4.3	3.1	3.6	3.6	3.7	4.7	5.7	7.1	7.2	6.2	5.2	4.6
1921	5.1	4.9	3.7	3.6	3.7	4.2	5.3	6.3	6.6	7.1	6.0	5.0	4.3
1920	4.7	3.1	3.2	3.8	3.7	4.0	4.6	5.0	6.9	6.1	6.2	5.5	4.2

See footnotes at end of table.

TABLE 16. - 1/ DEATH RATES FOR SELECTED CAUSES BY MONTH: JAPAN, 1920 - 1949 - Cont'd
17/ NEPHRITIS (ACUTE, CHRONIC AND UNSPECIFIED) (130-132)

Year	Annual	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
1949	41.2	46.8	46.1	47.7	42.7	37.1	35.8	36.0	35.4	38.4	41.3	43.3	44.0
1948	45.6	56.0	53.9	48.8	44.4	44.2	42.1	40.3	37.7	42.9	45.0	46.2	45.7
1947	57.7	67.0	72.7	66.0	61.1	55.0	51.9	49.4	49.8	51.2	57.5	55.5	56.5
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	66.9	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	69.3	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1941	68.2	80.6	81.6	68.8	65.3	58.8	57.1	58.0	60.4	68.3	71.5	73.8	74.9
1940	75.9	93.5	94.3	83.6	74.2	69.1	65.0	63.2	65.8	70.7	77.5	77.6	76.5
1939	82.2	102.2	94.4	92.1	77.8	71.0	68.6	68.0	71.2	76.8	85.2	86.5	93.5
1938	87.1	106.1	101.5	88.3	79.7	72.9	70.0	72.7	81.7	92.3	90.1	94.5	96.2
1937	79.3	86.5	80.1	78.5	73.5	67.7	68.2	62.8	73.4	82.6	91.4	90.0	96.6
1936	82.3	105.0	97.7	90.0	76.6	70.8	69.3	68.0	71.4	78.1	88.7	88.3	84.2
1935	80.9	91.1	90.0	81.3	75.1	70.0	67.7	68.2	73.4	82.1	90.4	89.5	92.4
1934	87.2	106.1	100.2	95.6	82.4	75.0	74.2	73.1	72.9	83.7	97.9	95.4	90.7
1933	89.0	98.9	102.3	96.6	82.6	75.5	75.3	74.0	78.3	93.1	100.2	97.7	94.5
1932	92.4	100.5	104.7	102.2	92.5	82.1	78.7	78.4	82.1	89.7	99.4	98.5	100.6
1931	98.2	110.9	108.9	102.0	93.3	86.4	87.0	84.2	85.0	101.0	109.5	106.4	105.0
1930	98.4	107.6	103.6	101.2	92.3	91.4	87.7	85.2	91.5	104.5	107.2	106.7	102.6
1929	103.0	112.0	113.4	109.0	97.0	90.9	89.2	92.2	97.5	106.5	110.9	113.9	104.0
1928	100.3	106.4	103.5	100.5	89.5	82.7	80.1	85.8	94.6	110.0	123.6	116.2	110.4
1927	97.3	104.1	106.9	104.8	87.8	86.3	82.6	78.7	91.7	106.1	111.0	106.4	101.8
1926	98.0	105.3	102.6	102.5	96.6	89.2	86.3	83.3	88.3	99.1	111.3	108.3	103.4
1925	100.2	113.7	119.6	106.0	93.2	88.2	87.8	89.3	87.3	96.3	111.4	104.3	106.8
1924	105.6	112.0	109.9	109.3	91.1	91.5	90.7	92.1	112.5	115.0	117.9	115.7	110.1
1923	107.5	112.5	105.6	96.0	91.3	86.2	92.1	97.2	104.0	109.6	139.9	100.9	111.6
1922	106.0	118.8	112.2	103.7	89.0	84.5	83.9	89.5	104.9	123.1	124.9	123.2	115.2
1921	105.4	96.8	103.7	96.9	83.9	82.3	84.0	87.1	107.7	133.5	138.7	131.5	116.5
1920	99.3	103.2	99.3	92.0	82.5	76.8	79.2	85.7	105.7	121.4	119.9	113.0	112.3

See footnotes at end of table.

TABLE 16. - 1/ DEATH RATES FOR SELECTED CAUSES BY MONTH: JAPAN, 1920 - 1949 - Cont'd
18/ DISEASE OF PREGNANCY, CHILDBIRTH AND THE PUERPERIUM (140-J 90)

Year	Annual	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
* 1949	5.5	5.8	5.5	5.8	6.1	5.3	5.1	4.9	5.9	5.3	5.3	5.2	5.3
1948	5.5	6.3	6.7	6.1	5.4	4.9	4.8	5.2	5.8	5.9	5.3	4.7	5.3
1947	5.8	6.2	6.5	5.2	5.4	5.3	5.3	6.1	5.8	6.1	5.3	5.9	5.9
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	6.1	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	6.2	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1941	7.9	7.9	7.8	7.3	6.1	5.2	5.6	6.5	6.5	7.3	6.9	6.4	7.5
1940	7.0	8.4	9.3	7.5	6.4	5.4	5.8	7.1	6.6	7.1	6.1	6.9	7.7
1939	6.7	8.2	7.5	6.8	5.8	5.7	5.0	6.3	6.9	7.2	6.9	7.0	7.7
1938	6.8	8.4	8.7	7.5	6.0	5.6	5.4	6.4	7.9	7.3	6.2	6.6	6.3
1937	7.7	8.3	9.0	7.3	7.6	6.3	6.0	6.9	8.1	8.3	7.7	8.0	8.6
1936	7.7	9.5	8.7	8.5	7.3	5.9	6.8	7.4	7.2	8.2	7.6	7.6	7.5
1935	8.2	9.4	9.5	8.4	7.7	6.6	6.6	7.4	8.4	8.4	9.1	8.6	8.6
1934	8.4	10.5	9.6	9.0	7.7	7.5	7.5	7.3	8.4	8.4	8.1	8.0	8.4
1933	8.6	9.7	9.1	10.1	8.2	7.1	7.1	7.7	8.4	8.8	8.8	9.5	8.1
1932	8.3	8.7	9.5	8.7	8.7	7.1	6.9	7.9	8.9	8.3	8.8	8.2	8.2
1931	8.5	9.0	9.0	9.2	7.9	8.2	8.1	8.1	9.3	9.1	9.4	8.5	7.7
1930	8.8	9.3	9.3	9.2	7.5	8.2	7.4	9.0	9.7	9.5	8.7	8.7	9.3
1929	9.2	9.6	9.8	10.3	9.1	8.3	7.7	10.0	11.5	9.6	8.9	8.2	8.0
1928	9.6	9.5	10.1	9.9	8.9	8.3	8.7	8.8	10.1	10.9	10.6	9.8	9.3
1927	9.3	10.6	10.6	10.5	9.4	8.3	8.3	8.8	9.8	9.8	10.4	8.5	8.5
1926	9.4	9.6	10.1	10.1	8.7	8.5	8.6	8.4	9.7	10.2	10.0	9.3	8.9
1925	10.6	11.0	12.9	11.8	9.9	9.4	9.7	10.3	10.7	10.7	11.0	9.8	9.7
1924	10.7	10.5	10.9	10.6	9.8	9.1	8.4	10.8	12.6	13.0	11.5	10.4	10.2
1923	11.9	11.9	12.8	11.5	10.4	8.7	10.0	13.1	14.1	15.0	13.1	11.5	10.3
1922	12.8	12.8	12.9	11.5	10.2	9.5	8.8	11.3	12.4	13.1	13.1	10.8	11.0
1921	12.7	13.6	13.2	12.0	10.5	10.7	10.0	12.9	14.8	15.2	14.6	12.9	11.6
1920	12.8	15.1	14.7	12.9	11.8	10.5	10.2	12.4	13.0	14.8	13.4	12.1	12.7

See footnotes at end of table.

TABLE 16. - 1/ DEATH RATES FOR SELECTED CAUSES BY MONTH: JAPAN, 1920 - 1949 - Cont'd
CONGENITAL MALFORMATION (157)

Year	Annual	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
1949	8.0	9.2	9.1	9.7	8.3	7.1	6.9	6.6	5.9	7.3	7.7	9.0	9.2
1948	7.2	8.0	8.4	8.5	6.8	5.7	5.7	6.6	6.3	7.0	6.8	7.7	8.6
1947	6.4	6.3	6.6	6.5	5.6	5.6	5.5	6.3	6.2	5.7	6.4	8.1	8.0
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	5.1	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	5.3	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1941	5.0	5.5	6.3	5.5	5.3	4.5	4.4	4.5	4.8	4.1	5.1	5.1	5.2
1940	4.9	6.0	6.6	5.4	5.1	4.6	3.9	4.4	3.5	4.1	4.3	5.0	5.6
1939	5.2	6.2	6.4	6.3	5.0	4.2	3.8	4.3	4.6	4.6	4.8	5.2	7.1
1938	5.5	6.6	7.3	6.3	5.6	5.2	4.3	5.0	4.8	5.0	5.6	5.5	5.9
1937	6.0	7.1	7.0	7.0	6.2	5.5	5.2	5.3	5.0	5.5	5.6	6.1	7.2
1936	6.1	7.8	7.2	7.3	6.2	5.0	4.9	5.6	5.6	5.8	6.1	5.8	6.7
1935	6.3	7.2	7.2	6.7	6.4	5.8	5.1	4.9	5.4	6.1	6.1	6.6	7.7
1934	6.1	7.6	6.6	7.7	6.4	5.6	5.0	5.4	5.3	5.2	6.1	6.2	6.7
1933	6.5	8.2	8.3	7.4	5.9	5.9	5.5	5.3	5.5	6.3	6.2	6.3	7.2
1932	4.2	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1931	4.3	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1930	4.4	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1929	4.8	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1928	4.9	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1927	4.9	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1926	5.5	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1925	5.1	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1924	4.7	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1923	3.7	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1922	2.2	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1921	2.3	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1920	2.2	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA

See footnotes at end of table.

TABLE 16. - 1/ DEATH RATES FOR SELECTED CAUSES BY MONTH: JAPAN, 1920 - 1949 - Cont'd
CONGENITAL DEBILITY (158)

Year	Annual	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
* 1949	45.4	70.0	63.8	66.8	50.3	36.2	32.9	34.2	31.1	29.6	32.9	44.5	53.0
1948	47.6	80.5	81.5	67.8	44.8	34.8	32.6	32.2	31.1	32.8	34.1	42.9	57.6
1947	53.5	72.6	90.1	75.7	52.2	37.9	34.1	40.0	35.1	33.8	40.4	56.1	76.7
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	69.9	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	70.6	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1941	71.6	99.6	111.6	87.1	68.8	53.7	54.5	58.8	56.6	56.9	61.0	66.5	86.2
1940	76.2	119.9	121.2	104.9	74.4	58.7	54.0	61.8	57.4	55.2	58.2	65.9	83.8
1939	79.9	131.3	120.9	102.2	71.7	57.4	51.6	58.1	58.1	58.4	65.7	78.5	106.9
1938	85.1	143.4	145.6	107.4	79.6	60.0	57.2	61.5	59.5	61.1	65.9	81.9	101.7
1937	89.5	121.7	116.7	106.0	82.5	65.7	66.2	68.8	69.1	72.6	82.4	93.7	127.3
1936	96.8	160.3	154.2	134.4	94.5	70.6	69.4	74.0	70.7	68.0	76.9	88.2	128.8
1935	92.9	131.5	134.2	111.2	83.4	68.0	64.6	66.9	70.4	75.5	83.0	99.4	126.7
1934	100.4	152.2	152.9	137.4	100.0	76.6	71.1	74.5	72.8	74.5	87.1	97.1	112.3
1933	100.1	153.7	153.1	136.6	96.4	74.1	72.6	74.7	72.3	71.6	84.2	96.8	118.8
1932	89.4	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1931	94.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1930	92.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1929	103.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1928	101.9	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1927	103.9	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1926	110.6	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1925	112.2	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1924	118.1	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1923	126.7	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1922	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1921	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1920	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA

See footnotes at end of table.

TABLE 16. - 1/ DEATH RATES FOR SELECTED CAUSES BY MONTH: JAPAN, 1920 - 1949 - Cont'd
PREMATURE BIRTH (159)

Year	Annual	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
1949	16.6	21.3	20.2	19.5	16.5	14.5	14.4	14.4	14.4	14.6	13.8	16.5	19.7
1948	17.1	23.6	23.0	19.4	15.0	14.0	14.2	13.5	13.9	14.3	15.8	18.5	20.4
1947	18.7	25.6	26.5	20.6	16.8	14.8	14.9	16.3	15.5	14.1	15.4	20.9	23.8
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	6.8	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	7.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1941	7.2	9.5	10.3	7.4	5.8	5.8	6.6	6.0	6.3	6.7	7.2	7.2	8.3
1940	7.3	11.0	10.0	8.1	6.7	6.3	5.5	6.0	6.3	6.0	6.4	7.3	8.2
1939	7.2	10.3	9.6	7.7	5.6	5.7	4.9	5.8	6.3	5.9	7.5	7.9	9.8
1938	7.6	12.1	11.5	8.6	6.8	5.8	5.8	5.8	5.6	6.1	6.3	7.8	9.2
1937	8.4	10.8	10.1	8.5	6.9	7.0	6.7	7.0	7.9	7.1	8.4	9.6	10.3
1936	9.0	12.9	13.0	10.5	7.8	6.5	7.2	7.2	6.8	7.0	8.4	9.5	11.4
1935	8.2	11.0	10.4	7.3	6.6	5.8	7.0	6.5	7.1	8.4	8.2	9.0	11.1
1934	8.3	11.5	11.3	9.4	7.8	5.7	5.9	6.5	6.6	7.1	8.1	8.9	10.8
1933	7.5	11.2	9.6	8.2	6.5	5.9	5.8	5.9	6.2	6.7	7.2	7.5	9.6
1932	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1931	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1930	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1929	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1928	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1927	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1926	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1925	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1924	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1923	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1922	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1921	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1920	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA

See footnotes at end of table.

TABLE 16. - 1/ DEATH RATES FOR SELECTED CAUSES BY MONTH: JAPAN, 1920 - 1949 - Cont'd
INJURY AT BIRTH (160)

Year	Annual	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
*1949	1.3	1.3	1.4	1.3	1.4	1.1	1.3	1.4	1.3	1.3	1.4	1.3	1.3
**1948	1.3	1.6	1.7	1.5	1.2	1.2	1.1	1.2	1.3	1.2	0.8	1.0	1.5
1947	1.1	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	0.6	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	0.6	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1941	0.6	0.6	1.0	0.7	0.7	0.5	0.4	0.5	0.7	0.6	0.5	0.6	0.7
1940	0.4	0.5	0.3	0.7	0.4	0.3	0.4	0.5	0.4	0.4	0.3	0.3	0.4
1939	0.4	0.3	0.6	0.3	0.5	0.4	0.4	0.5	0.4	0.3	0.5	0.4	0.7
1938	0.5	0.7	0.7	0.6	0.5	0.5	0.5	0.6	0.6	0.5	0.4	0.4	0.4
1937	0.6	0.6	0.7	0.7	0.6	0.7	0.6	0.6	0.7	0.5	0.6	0.5	0.6
1936	0.6	0.6	0.7	0.8	0.6	0.3	0.3	0.6	0.7	0.7	0.6	0.5	0.6
1935	0.6	0.7	0.7	0.8	0.6	0.5	0.5	0.7	0.6	0.6	0.5	0.7	0.6
1934	0.6	0.8	0.6	0.9	0.5	0.5	0.5	0.5	0.4	0.7	0.6	0.7	0.5
1933	0.5	0.5	0.7	0.7	0.4	0.3	0.4	0.7	0.6	0.5	0.4	0.5	0.7
1932	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1931	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1930	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1929	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1928	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1927	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1926	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1925	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1924	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1923	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1922	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1921	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1920	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA

See footnotes at end of table.

TABLE 16. - 1/ DEATH RATES FOR SELECTED CAUSES BY MONTH: JAPAN, 1920 - 1949 . Cont'd
19/ SENILITY (162)

Year	Annual	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
*1949	81.1	95.0	93.5	97.6	87.7	68.0	63.2	65.3	68.5	71.9	79.4	86.6	97.6
1948	79.9	119.6	113.4	105.1	85.9	63.1	61.1	60.7	63.4	63.4	68.6	73.9	81.4
1947	101.2	140.0	163.5	134.9	103.2	84.5	76.5	73.9	77.0	72.1	81.6	94.6	117.1
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	134.1	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	130.7	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1941	123.2	163.0	171.4	132.8	112.3	95.1	93.3	107.1	110.5	118.8	122.6	118.8	136.1
1940	124.9	163.5	169.5	143.5	116.7	97.9	91.4	104.5	123.6	120.6	116.6	116.4	136.1
1939	133.5	171.6	166.9	142.8	122.1	100.3	96.5	109.1	128.3	131.9	130.8	128.3	152.2
1938	138.7	185.9	194.7	149.8	117.8	97.8	96.6	120.5	140.9	151.0	138.5	135.0	139.9
1937	119.4	130.0	131.0	117.7	105.0	90.7	90.1	97.2	126.8	130.8	134.1	123.9	155.6
1936	131.0	196.6	207.8	160.2	118.0	95.7	93.7	102.2	135.3	130.5	127.0	120.5	118.8
1935	114.4	137.7	136.2	117.3	104.3	87.0	88.5	97.5	118.6	117.3	119.1	116.1	134.5
1934	127.9	180.0	177.4	158.6	118.1	97.7	91.8	96.0	112.6	121.2	130.9	129.3	123.7
1933	123.4	147.6	161.8	144.4	111.5	95.1	91.5	101.1	124.2	125.6	126.9	122.8	131.0
1932	116.8	130.2	138.5	136.3	114.5	93.5	86.0	98.8	116.0	120.7	124.4	117.2	125.4
1931	131.0	165.1	181.7	161.6	131.5	108.2	101.0	103.5	113.9	125.1	133.4	122.4	127.6
1930	118.8	142.4	138.1	120.4	105.4	101.2	96.0	104.7	121.0	127.7	121.5	118.5	130.0
1929	124.7	151.4	156.8	142.0	112.3	102.3	92.9	109.0	130.0	132.2	125.7	120.7	110.8
1928	123.2	140.3	143.0	140.6	118.1	90.0	86.5	98.3	119.9	135.9	144.3	127.7	134.3
1927	121.0	137.9	161.6	147.2	111.7	100.8	91.9	97.0	122.6	133.3	120.2	113.7	117.1
1926	111.1	131.1	124.8	118.1	103.3	88.0	83.5	87.0	107.7	117.2	126.7	119.2	127.7
1925	117.3	142.7	187.9	158.3	108.6	91.4	82.9	93.4	100.7	110.4	113.5	107.1	115.4
1924	125.6	149.5	152.8	154.6	114.4	100.2	91.3	100.1	134.6	129.4	131.2	125.4	124.5
1923	131.6	143.3	142.3	133.4	111.3	94.7	97.9	117.4	143.0	164.1	163.0	136.2	132.7
1922	132.3	171.9	177.2	161.9	120.1	97.5	89.8	103.2	130.1	145.6	134.7	124.2	134.4
1921	133.4	141.0	147.7	135.5	110.0	92.3	95.6	109.2	145.0	170.9	158.8	155.1	141.4
1920	131.3	163.6	172.0	149.8	112.6	99.8	91.9	104.0	137.9	152.4	133.4	118.2	134.9

See footnotes at end of table.

TABLE 16. - 1/ DEATH RATES FOR SELECTED CAUSES BY MONTH: JAPAN, 1920 - 1949 - Cont'd
SUICIDE (163-164)

Year	Annual	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
* 1949	17.3	12.4	13.6	16.0	19.9	22.8	18.5	20.0	18.4	17.3	16.3	16.1	15.8
1948	15.9	11.9	13.0	16.4	18.9	19.2	19.1	19.7	17.6	16.3	13.9	12.4	12.2
1947	15.7	10.5	10.4	14.8	18.8	21.6	19.0	21.3	17.7	15.7	13.8	12.9	11.5
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	11.8	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	12.7	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1941	13.3	10.8	11.7	13.7	15.4	15.8	15.2	15.5	14.7	11.9	12.4	11.7	10.4
1940	13.7	10.4	11.1	13.5	15.5	17.9	15.0	17.0	14.5	13.5	12.6	12.1	11.1
1939	15.1	12.3	13.2	15.4	18.0	19.6	16.8	17.5	16.1	14.6	13.4	12.7	10.9
1938	17.2	13.2	15.0	18.8	21.7	22.9	19.9	19.2	19.5	16.1	14.6	12.7	12.2
1937	20.1	17.2	18.7	21.0	25.9	25.4	22.7	24.6	21.7	18.8	16.5	15.5	13.3
1936	22.0	16.4	18.1	21.9	26.5	28.8	26.3	25.4	23.3	22.0	19.5	18.5	16.8
1935	20.5	16.5	18.6	20.5	24.3	25.7	22.6	23.4	21.0	18.8	16.7	18.5	16.6
1934	21.3	18.2	17.9	21.1	25.9	26.5	24.6	24.4	23.2	19.5	18.0	17.8	18.4
1933	22.0	18.2	19.3	21.7	27.4	29.1	25.7	27.0	22.9	20.7	17.5	18.1	15.9
1932	22.2	19.1	17.3	21.7	24.7	28.1	26.2	28.6	25.7	21.9	19.1	17.4	16.5
1931	21.9	16.6	17.8	22.7	25.3	26.8	27.0	24.5	24.8	20.1	19.1	18.6	19.6
1930	21.6	16.6	17.8	21.4	25.0	25.9	24.8	27.7	24.1	22.1	18.0	18.2	17.6
1929	20.1	15.6	15.6	20.0	23.6	24.5	22.8	24.8	21.5	18.9	18.5	17.0	17.6
1928	20.8	16.8	17.3	21.2	25.9	26.5	22.9	23.7	21.9	20.8	18.4	16.4	15.4
1927	20.8	15.7	16.3	19.1	25.0	25.2	24.6	26.4	22.5	20.0	17.7	17.5	17.3
1926	20.5	16.6	16.5	19.9	23.0	27.0	23.5	26.3	23.5	19.9	17.5	16.9	15.2
1925	20.5	14.7	16.2	19.0	24.3	27.2	26.2	26.0	21.0	20.1	18.5	17.1	15.0
1924	19.1	14.5	15.1	18.6	22.4	24.5	21.9	24.4	21.7	18.1	17.3	15.9	14.5
1923	19.7	14.8	15.9	21.0	22.7	23.6	24.5	25.1	22.1	19.7	16.1	15.0	13.7
1922	20.1	15.5	19.5	20.2	23.9	25.0	23.9	25.0	22.6	18.5	17.6	14.9	14.4
1921	20.0	16.2	16.3	18.4	23.2	24.8	21.9	24.0	24.0	19.3	18.8	16.6	16.4
1920	19.0	13.9	14.7	16.4	21.3	23.0	22.9	25.0	23.4	20.2	16.9	15.6	14.3

See footnotes at end of table.

TABLE 16. - 1/ DEATH RATES FOR SELECTED CAUSES BY MONTH: JAPAN, 1920 - 1949 - Cont'd
HOMICIDE (165-168)

Year	Annual	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
*1949	2.0	1.6	1.9	2.1	2.0	2.1	1.7	2.1	2.4	2.4	1.9	2.1	2.0
1948	2.0	1.8	1.9	2.3	2.8	2.3	1.8	2.4	2.2	1.7	1.7	1.7	1.9
1947	1.7	1.0	1.3	1.6	2.0	1.9	2.0	1.9	1.8	1.7	1.7	1.6	1.7
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	0.3	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	0.4	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1941	0.3	0.3	0.3	0.3	0.3	0.2	0.3	0.3	0.5	0.4	0.3	0.3	0.4
1940	0.4	0.2	0.4	0.3	0.4	0.4	0.4	0.5	0.5	0.5	0.3	0.3	0.3
1939	0.4	0.3	0.3	0.5	0.2	0.5	0.4	0.4	0.4	0.2	0.4	0.3	0.4
1938	0.6	0.5	0.6	0.9	0.7	1.3	0.8	0.5	0.3	0.5	0.5	0.4	0.5
1937	0.6	0.7	0.7	0.7	0.5	1.0	0.8	0.7	0.5	0.5	0.5	0.6	0.4
1936	0.7	0.4	0.8	0.9	1.0	0.8	0.5	0.6	0.8	0.7	0.6	0.5	0.5
1935	0.6	0.6	0.5	0.6	0.5	0.5	1.0	0.9	0.5	0.5	0.7	0.7	0.4
1934	0.6	0.6	0.4	0.6	0.7	0.8	0.8	0.7	0.7	0.7	0.6	0.5	0.4
1933	0.7	0.7	0.6	0.8	0.6	0.7	0.9	0.7	0.7	0.6	0.8	0.6	0.5
1932	0.7	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1931	0.8	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1930	0.6	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1929	0.6	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1928	0.7	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1927	0.7	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1926	0.8	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1925	0.7	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1924	0.7	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1923	0.6	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1922	0.8	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1921	0.6	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1920	0.7	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA

See footnotes at end of table.

TABLE 16. - 1/ DEATH RATES FOR SELECTED CAUSES BY MONTH: JAPAN, 1920 - 1949 - Cont'd
ACCIDENTAL DEATHS (1920-1949)

Year	Annual	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
* 1919	42.0	32.3	31.5	36.1	37.6	41.1	46.3	61.9	63.9	46.9	36.2	33.2	35.7
1918	48.6	39.7	35.6	37.0	38.0	40.0	100.5	65.6	61.6	56.5	39.1	33.1	36.3
1917	49.3	39.5	43.3	39.9	45.2	44.6	43.8	77.1	75.8	66.2	39.5	36.6	38.8
1916	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1915	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1914	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1913	45.1	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1912	42.1	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1911	39.3	35.1	32.8	35.5	37.1	36.8	41.4	51.1	58.2	38.9	38.9	31.1	33.4
1910	39.3	41.6	34.8	33.9	36.9	37.4	40.7	58.1	49.8	39.5	33.0	31.7	33.3
1909	40.9	41.6	37.6	36.6	38.5	39.2	40.4	63.0	52.1	39.6	35.6	32.6	33.5
1908	44.4	40.4	38.4	37.5	38.9	39.4	47.2	67.9	57.4	48.8	45.4	34.9	36.1
1907	42.4	33.1	36.7	36.8	39.3	37.7	43.7	60.0	65.5	41.8	37.4	37.5	40.2
1906	42.8	40.2	39.9	38.2	43.8	38.9	44.7	60.9	52.9	43.8	38.2	38.3	34.1
1905	41.7	34.0	34.2	36.1	38.6	42.1	45.2	61.3	54.1	46.0	39.5	33.6	34.5
1904	46.6	37.3	36.7	67.8	38.1	38.3	43.2	55.1	49.8	87.0	35.9	34.4	35.0
1903	44.4	34.4	36.1	81.7	36.6	39.4	44.4	64.4	54.7	39.9	36.8	31.7	31.4
1902	40.1	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1901	38.5	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1900	40.6	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1909	41.8	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1908	41.4	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1907	47.4	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1906	42.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1905	41.7	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1904	43.6	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1903	122.5	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1902	44.3	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1901	43.1	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1900	46.6	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA

See footnotes at end of table.

TABLE 16. - 1/ DEATH RATES FOR SELECTED CAUSES BY MONTH: JAPAN, 1920 - 1945 - Cont'd
20/ III*DEFINED AND UNKNOWN (200+ and 200e)

Year	Annual	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
* 1949	20.6	23.7	23.8	24.6	23.6	19.1	19.0	18.6	17.7	16.5	17.9	20.1	22.4
**1948	21.4	27.0	27.6	27.1	28.8	20.7	20.0	18.8	19.3	17.5	18.0	18.4	20.2
**1947	25.8	29.9	34.8	32.8	28.6	23.8	20.6	23.7	24.0	20.0	20.5	22.2	27.6
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	42.6	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	43.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1941	42.3	57.9	64.3	50.7	43.3	38.9	34.9	33.7	32.3	30.9	35.9	38.5	45.4
1940	45.8	66.6	67.7	57.9	50.6	43.2	38.8	36.9	35.2	34.4	34.9	37.3	44.5
1939	49.2	68.4	68.2	66.7	53.5	44.4	40.2	37.6	37.5	37.6	39.2	42.2	55.8
1938	50.9	76.7	81.9	64.0	53.2	43.1	38.9	38.0	39.0	41.8	39.1	44.9	52.7
1937	46.2	57.7	54.9	54.0	47.2	43.0	39.6	36.4	36.6	37.8	41.9	43.6	62.4
1936	53.9	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1935	48.5	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1934	54.7	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1933	56.3	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1932	61.7	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1931	68.2	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1930	63.8	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1929	71.6	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1928	72.5	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1927	69.6	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1926	64.3	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1925	68.3	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1924	73.4	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1923	78.8	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1922	126.5	165.7	168.6	165.2	119.3	103.0	97.9	104.4	111.8	117.2	116.4	117.7	133.1
1921	127.8	133.0	148.6	148.8	113.5	106.8	102.2	111.3	127.3	130.7	134.6	138.8	140.9
1920	132.0	177.8	189.2	192.3	132.6	108.8	100.8	109.0	115.9	120.9	119.6	114.6	137.7

See footnotes at end of table.

TABLE 16. - 1/ DEATH RATES FOR SELECTED CAUSES BY MONTH: JAPAN, 1920 - 1949 - Cont'd

FOOTNOTES:

- * Data for 1949 are provisional.
 ** The 1947 and 1948 monthly death rates for syphilis (30) and ill-defined and unknown causes (200 and c) and the 1948 monthly data for simple meningitis (81a), pleurisy (110) and injury at birth (160) are estimates based on preliminary figures. The annual data are final. Rates are the number of deaths per 100,000 population (per annum), estimated as of 1 October 1920 - 1945 and 1 July 1946 - 1949.
 1/ Estimated populations used to compute all rates 1920 - 1941 and annual rates 1942 - 1943 include civilian and military in Japan and Okinawa. All rates 1944 - 1949 and monthly rates 1942 - 1943 are based upon the estimated civilian population in Japan Proper only.
 2/ Typhoid fever, 1920-1943: includes typhoid pneumonia. 1944-1946, unknown whether typhoid pneumonia is included. 1947-1949, excludes typhoid pneumonia.
 3/ Scarlet fever, 1920-1943: includes puerperal scarlet fever. 1944-1946: unknown whether puerperal scarlet fever is included. 1947-1949: excludes puerperal scarlet fever.
 4/ Diphtheria, 1920-1922 and 1933-1943: excludes diphtheric endocarditis; includes croupous bronchitis and laryngeal croup. 1923-1932: excludes diphtheric endocarditis and croupous bronchitis; includes laryngeal croup. 1944-1946: unknown whether diphtheric endocarditis, croupous bronchitis, and laryngeal croup are included. 1947-1949: includes diphtheric endocarditis; excludes croupous bronchitis and laryngeal croup.
 5/ Tuberculosis (all forms). 1920-1943: includes chronic pleuro pneumonia; excludes spondylitis; includes all adenopathy. 1947-1949: excludes chronic pleuro pneumonia; includes spondylitis; excludes nontubercular adenopathy. 1944-1946: are estimates of the Ministry of Welfare.
 6/ Syphilis. 1920-1922: excludes general paralysis of the insane; excludes aneurysm of the aorta. 1923-1932: includes general paralysis of the insane, a small amount of non-syphilitic paralytic psychosis, aneurysm of the aorta and a small amount of non-syphilitic aneurysm. 1933-1943: includes general paralysis of the insane, aneurysm of the aorta and a small amount of non-syphilitic aneurysm. 1947-1949: includes general paralysis of the insane and aneurysm of the aorta.
 7/ Typhus fever and typhus-like diseases. 1920-1943, 1947-1949: includes typhus-like diseases. 1944-1946: excludes typhus-like diseases.
 8/ Simple meningitis. 1920-1943: includes acute cerebrospinal meningitis not due to meningococcus. 1947-1949: excludes acute cerebrospinal meningitis not due to meningococcus.
 9/ Intracranial lesions of vascular origin. 1920-1932: includes alcoholic cerebral apoplexy and congestion and a small amount of paralysis not due to intracranial lesions of vascular origin. 1933-1943: includes alcoholic cerebral apoplexy and congestion. 1947-1949: excludes alcoholic apoplexy and congestion.
 10/ Diseases of the heart. 1920-1943: includes diphtheric endocarditis. 1947-1949: excludes diphtheric endocarditis.
 11/ Bronchitis. 1920-1922 and 1933-1943: excludes croupous bronchitis. 1923-1932 and 1947-1949: includes croupous bronchitis.
 12/ Pneumonia (all forms). 1920-1943: excludes typhoid pneumonia. 1947-1949: includes typhoid pneumonia.
 13/ Ulcer of stomach or duodenum. 1920-1922: excludes ulcer of duodenum. 1923-1943: includes ulcer of duodenum.
 14/ Diarrhea, enteritis and ulceration of the intestines (all ages). 1920-1943: includes malnutrition 1 year of age and over. 1947-1949: excludes malnutrition 1 year of age and over.
 15/ Diarrhea, enteritis and ulceration of the intestines (under 2 years). 1920-1943: includes malnutrition 1 to 2 years of age. 1947-1949: excludes malnutrition 1 to 2 years of age.

TABLE 16. - 1/ DEATH RATES FOR SELECTED CAUSES BY MONTH: JAPAN, 1920 - 1949 - Cont'd

FOOTNOTES - Cont'd:

- 16/ Diarrhea, enteritis and ulceration of the intestines (2 years and over). 1920-1943: includes malnutrition, 2 years of age and over. 1947-1949: excludes malnutrition, 2 years of age and over.
- 17/ Nephritis. 1920-1943: includes alcoholic nephritis. 1947-1949: excludes alcoholic nephritis.
- 18/ Diseases of pregnancy, childbirth and the puerperium. 1920-1943: excludes puerperal scarlet fever. 1947-1949: includes puerperal scarlet fever.
- 19/ Senility. 1920-1922: excludes senility with mention of senile dementia, psychosis and paresis. 1923-1943 and 1947-1949: includes senility with mention of senile dementia.
- 20/ Ill-defined and unknown. 1920-1922: excludes malnutrition 1 year of age and over. Includes sudden death and found dead. 1923-1943: excludes malnutrition 1 year of age and over. Excludes sudden death and found dead. 1947-1949: includes malnutrition 1 year of age and over. Excludes sudden death and found dead.

"NA" indicates data are not available.

A dash (-) indicates that no deaths were reported.

A rate of 0.0 indicates that there were some deaths but that the rate was less than 0.1.

Sources:

Rates computed by Public Health and Welfare Section, GHQ, SCAP.
Sources of original death data are as follows:

- 1920-1941: Annual and Monthly data - Annual Reports, Cabinet Bureau of Statistics.
- 1942-1943: Annual data - Annual Reports, Cabinet Bureau of Statistics.
Monthly data - Unpublished Annual Reports, Ministry of Welfare.
- 1944-1945: Annual and Monthly data - Unpublished Annual Reports, Ministry of Welfare.
- 1946: Annual and Monthly data - Weekly Reports, Ministry of Welfare.
- 1947-1948: Annual and Monthly data - Final Annual Schedule Report, Ministry of Welfare.
- 1949: Annual and Monthly data - Monthly Schedule Reports, Ministry of Welfare.

TABLE 17. - 1/ DEATHS FROM TEN SELECTED CAUSES BY PREFECTURE: JAPAN, 1920-1949
2/ TUBERCULOSIS, ALL FORMS (13-22)

Year	All Japan	Aichi	Akita	Aomori	Chiba	Ehime	Fukui	Fukushima	Gifu	Gumma
1949	138,764	5,383	2,120	2,710	2,961	2,268	1,166	2,997	2,575	2,007
1948	143,909	5,547	2,280	2,777	3,069	2,471	1,206	3,110	2,972	2,326
1947	146,241	5,884	2,079	2,567	2,956	2,617	1,397	3,165	2,773	2,382
1946	198,900	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	203,000	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	176,800	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	171,473	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	160,398	6,860	1,904	2,517	2,720	2,536	1,531	2,510	2,641	2,040
1941	153,250	6,625	1,691	2,265	2,477	2,602	1,501	2,373	2,704	2,021
1940	152,019	6,657	1,687	2,276	2,513	2,615	1,611	2,517	2,773	2,275
1939	153,209	6,578	1,545	2,275	2,406	2,628	1,497	2,314	2,825	2,324
1938	147,685	6,618	1,524	1,972	2,137	2,554	1,566	2,317	2,917	2,192
1937	143,424	6,126	1,470	1,846	2,051	2,536	1,634	2,207	2,985	2,178
1936	143,855	6,387	1,466	2,002	2,128	2,528	1,730	2,258	2,893	2,191
1935	130,763	5,703	1,336	1,736	1,913	2,401	1,464	2,052	2,619	2,013
1934	130,262	5,571	1,341	1,823	2,028	2,367	1,576	2,240	2,679	2,032
1933	125,492	5,392	1,377	1,649	2,053	2,169	1,464	2,046	2,699	2,028
1932	118,023	5,040	1,237	1,586	1,936	2,060	1,545	1,952	2,533	1,943
1931	120,629	4,967	1,246	1,578	1,937	2,260	1,563	2,172	2,565	2,080
1930	118,345	4,751	1,164	1,416	2,104	2,175	1,404	2,092	2,511	2,163
1929	122,221	5,034	1,192	1,606	2,018	2,313	1,463	2,219	2,527	2,270
1928	118,377	5,018	1,083	1,527	2,032	2,194	1,354	2,153	2,467	2,154
1927	118,112	4,802	1,142	1,473	2,001	2,333	1,420	2,249	2,458	2,165
1926	111,728	4,376	1,085	1,468	1,962	2,245	1,536	2,130	2,263	2,027
1925	114,650	4,545	1,065	1,476	1,963	2,316	1,479	2,300	2,351	2,344
1924	112,938	4,472	1,093	1,438	2,095	2,233	1,403	2,327	2,372	2,241
1923	117,036	4,728	1,134	1,348	2,249	2,306	1,553	2,367	2,277	2,208
1922	124,328	4,685	1,181	1,561	2,459	2,356	1,774	2,548	2,308	2,234
1921	119,526	4,698	1,115	1,196	2,319	2,357	1,830	2,328	2,116	2,104
1920	123,988	5,182	1,143	1,219	2,372	2,361	1,736	2,367	2,352	2,090

TABLE 17. - 1/ DEATHS FROM TEN SELECTED CAUSES BY PREFECTURE, JAPAN, 1920-1949 Cont'd
2/ TUBERCULOSIS, ALL FORMS (13-22) (Cont'd)

Year	Hiroshima	Hokkaido	Hyogo	Ibaraki	Ishikawa	Iwate	Kagawa	Kagoshima	Kanagawa	Kochi	Kumamoto	Kyoto
1949	3,324	10,284	5,478	2,315	1,631	2,737	1,398	2,771	4,240	1,358	2,796	3,915
1948	3,679	9,522	5,798	2,336	1,715	2,755	1,342	2,933	4,398	1,239	2,819	3,951
1947	3,826	8,919	6,392	2,273	1,788	2,394	1,676	2,776	4,387	1,365	2,993	4,280
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	4,433	9,467	8,307	2,066	2,114	2,114	1,681	3,834	5,300	1,472	2,663	4,877
1941	4,179	8,704	7,885	1,908	2,286	1,870	1,525	3,385	5,253	1,402	2,521	4,745
1940	3,834	9,351	7,812	1,974	2,454	1,700	1,601	3,211	5,208	1,341	2,252	4,546
1939	4,082	9,609	8,125	1,862	2,371	1,638	1,635	3,280	4,944	1,420	2,342	4,770
1938	3,726	8,754	7,751	1,834	2,606	1,524	1,502	3,091	4,640	1,312	2,228	4,697
1937	3,510	8,378	7,764	1,722	2,660	1,521	1,427	3,215	4,200	1,355	2,496	4,529
1936	3,600	8,360	7,759	1,756	2,689	1,525	1,459	2,976	4,328	1,253	2,474	4,649
1935	3,321	7,396	6,925	1,660	2,403	1,337	1,337	2,830	3,814	1,142	2,247	4,407
1934	3,217	7,308	6,612	1,664	2,302	1,438	1,382	2,709	3,890	1,143	2,324	4,284
1933	3,001	7,068	6,406	1,623	2,120	1,304	1,110	2,528	3,816	1,122	2,300	4,299
1932	2,792	6,286	6,008	1,561	2,212	1,201	1,283	2,407	3,495	1,057	2,249	3,900
1931	2,900	6,379	5,940	1,741	2,179	1,144	1,343	2,459	3,545	1,008	2,462	3,991
1930	2,896	6,080	5,907	1,634	2,204	1,098	1,378	2,526	3,340	1,053	2,457	3,862
1929	2,971	6,407	6,041	1,765	2,303	1,163	1,291	2,441	3,681	1,052	2,521	3,803
1928	2,972	5,832	5,725	1,741	2,084	1,064	1,368	2,384	3,405	1,021	2,435	3,909
1927	3,153	5,527	5,844	1,759	2,118	1,122	1,431	2,444	3,367	987	2,455	3,723
1926	2,898	5,189	5,556	1,606	2,118	1,098	1,305	2,311	3,018	1,014	2,184	3,573
1925	3,113	5,627	5,715	1,770	2,004	1,053	1,349	2,293	3,122	1,011	2,336	3,462
1924	3,190	5,310	5,664	1,828	2,106	1,032	1,413	2,248	2,838	949	2,376	3,506
1923	3,313	5,427	5,531	1,911	2,313	1,002	1,452	2,172	3,218	1,001	2,495	3,454
1922	3,457	5,734	5,829	1,966	2,531	1,154	1,454	2,191	3,714	1,022	2,429	3,628
1921	3,347	4,956	5,702	1,836	2,284	1,010	1,465	2,280	3,640	1,035	2,398	4,005
1920	3,877	4,672	6,151	2,010	2,239	1,051	1,663	2,174	3,639	1,006	2,479	4,146

TABLE 17. - DEATHS FROM TEN SELECTED CAUSES BY PREFECTURE: JAPAN, 1920-49 Cont'd
2/ TUBERCULOSIS, ALL FORMS (13-22) (Cont'd)

Year	Mie	Miyagi	Miyazaki	Nagano	Nagasaki	Nara	Niigata	Oita	Okayama	Osaka	Saga	Saitama
1949	2,239	2,690	1,810	2,620	2,717	1,232	4,237	2,248	2,443	7,259	1,643	2,918
1948	2,220	2,475	1,785	3,043	2,717	1,282	4,519	2,360	2,746	7,323	1,768	2,952
1947	2,482	2,561	1,878	2,980	3,084	1,503	4,414	2,516	2,767	7,802	1,945	2,887
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	2,586	2,211	1,693	2,626	2,911	1,233	3,897	2,341	2,818	13,090	1,527	2,438
1941	2,605	2,038	1,509	2,508	2,828	1,224	3,672	2,265	2,718	12,495	1,546	2,340
1940	2,327	1,973	1,394	2,587	2,694	1,173	3,752	2,105	2,712	12,355	1,419	2,433
1939	2,293	1,889	1,492	2,709	2,678	1,168	3,754	2,079	2,668	12,749	1,316	2,389
1938	2,302	1,920	1,289	2,629	2,936	1,160	3,751	1,989	2,411	12,143	1,332	2,414
1937	2,393	1,713	1,370	2,614	2,826	1,260	3,688	2,061	2,278	11,441	1,225	2,451
1936	2,346	1,926	1,349	2,617	2,749	1,234	3,720	2,019	2,307	11,938	1,281	2,469
1935	2,260	1,701	1,225	2,490	2,496	1,288	3,538	1,933	2,141	10,861	1,143	2,351
1934	2,226	1,756	1,107	2,592	2,557	1,231	3,742	1,921	2,142	9,818	1,160	2,425
1933	2,069	1,573	1,004	2,607	2,382	1,151	3,615	1,755	2,096	9,811	1,122	2,404
1932	2,070	1,592	997	2,399	2,268	1,029	3,454	1,766	1,924	9,041	1,088	2,233
1931	2,091	1,615	1,025	2,502	2,408	986	3,509	1,733	1,958	9,153	1,084	2,392
1930	1,979	1,528	949	2,378	2,376	1,006	3,330	1,735	1,980	8,865	1,193	2,414
1929	2,081	1,606	1,005	2,559	2,397	1,013	3,530	1,964	2,094	9,150	1,180	2,436
1928	2,164	1,515	923	2,532	2,378	1,029	3,548	1,808	2,090	8,662	1,094	2,413
1927	2,123	1,532	948	2,519	2,298	1,006	3,413	1,851	2,108	8,493	1,218	2,409
1926	1,939	1,435	890	2,413	2,179	906	3,470	1,766	2,035	7,499	1,153	2,408
1925	2,168	1,468	909	2,562	2,013	982	3,617	1,714	2,054	7,527	1,183	2,531
1924	2,095	1,378	821	2,592	2,023	997	3,657	1,726	2,109	7,302	1,217	2,559
1923	2,264	1,571	882	2,675	2,182	963	3,820	1,704	2,194	7,367	1,206	2,796
1922	2,309	1,629	814	2,755	2,163	1,045	4,004	1,806	2,267	8,094	1,288	2,861
1921	2,311	1,604	814	2,671	2,293	982	3,923	1,746	2,304	7,725	1,327	2,615
1920	2,394	1,510	834	2,860	2,363	1,070	3,977	1,810	2,395	8,199	1,326	2,770

TABLE 17. - 1/ DEATHS FROM TEN SELECTED CAUSES BY PREFECTURES: JAPAN, 1920-1949 Cont'd
2/ TUBERCULOSIS, ALL FORMS (13-22) (Cont'd)

Area	Shiga	Shimane	Shizuoka	Tochigi	Tokushima	Tokyo	Tottori	Toyama	Wakayama	Yamagata	Yamaguchi	Yamanashi
1949	1,350	1,772	3,204	1,852	1,533	11,632	1,013	1,906	1,422	2,048	2,915	769
1948	1,442	1,911	3,471	2,020	1,520	11,870	1,113	1,875	1,528	2,334	3,167	850
1947	1,538	2,154	3,590	2,025	1,716	10,692	1,186	1,670	1,514	2,210	3,430	791
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	1,553	1,675	3,926	1,882	1,741	18,259	1,102	1,906	1,898	1,853	3,047	830
1941	1,543	1,730	3,812	1,793	1,777	17,379	1,091	1,899	1,594	1,648	2,942	781
1940	1,469	1,726	3,882	1,701	1,750	17,524	953	1,896	1,668	1,621	2,764	881
1939	1,470	1,510	4,103	1,811	1,778	17,802	1,018	1,896	1,742	1,554	2,898	828
1938	1,559	1,619	3,750	1,669	1,705	17,666	1,001	2,026	1,684	1,503	2,780	809
1937	1,582	1,555	3,790	1,742	1,719	16,095	923	1,991	1,734	1,460	2,782	789
1936	1,616	1,532	3,687	1,747	1,599	15,875	969	1,935	1,704	1,465	2,553	834
1935	1,488	1,513	3,351	1,483	1,530	13,930	869	1,635	1,682	1,364	2,338	792
1934	1,558	1,504	3,368	1,573	1,410	14,131	844	1,666	1,595	1,455	2,318	847
1933	1,501	1,382	3,157	1,485	1,413	14,116	772	1,541	1,454	1,459	2,201	798
1932	1,400	1,410	2,976	1,464	1,413	12,750	778	1,552	1,406	1,352	2,077	767
1931	1,407	1,399	3,047	1,479	1,501	13,404	813	1,504	1,487	1,437	2,186	813
1930	1,330	1,509	3,060	1,540	1,417	12,515	772	1,512	1,509	1,366	2,179	846
1929	1,472	1,529	3,089	1,598	1,536	12,735	893	1,525	1,430	1,312	2,225	847
1928	1,440	1,528	3,148	1,553	1,561	12,173	774	1,536	1,494	1,272	2,113	832
1927	1,438	1,518	3,076	1,537	1,488	12,153	774	1,613	1,532	1,286	2,113	806
1926	1,376	1,460	2,992	1,435	1,466	11,296	696	1,655	1,432	1,206	2,097	890
1925	1,496	1,422	3,053	1,580	1,585	11,732	749	1,521	1,234	1,319	2,115	822
1924	1,461	1,400	2,935	1,579	1,558	11,065	710	1,482	1,248	1,261	2,078	935
1923	1,623	1,538	3,110	1,590	1,509	11,962	790	1,550	1,300	1,384	2,167	844
1922	1,619	1,587	3,217	1,660	1,578	13,849	819	1,725	1,436	1,479	2,271	1,010
1921	1,637	1,608	3,125	1,623	1,547	12,860	713	1,478	1,473	1,392	2,233	975
1920	1,719	1,473	3,073	1,599	1,720	13,519	665	1,537	1,399	1,429	2,333	991

See footnotes at end of table

TABLE 17 - DEATHS FROM TEN SELECTED CAUSES BY HOPFETURE: JAPAN, 1920-1949 Cont'd
CANCER AND OTHER MALIGNANT TUMORS (45 - 55)

Year	All Japan	Aichi	Akita	Aomori	Chiba	Ehime	Fuku	Fukuoka	Fukushima	Gifu	Gunma
1949	58,890	2,368	755	664	1,802	1,110	554	2,642	1,394	1,047	1,180
1948	55,677	2,228	769	624	1,763	1,080	475	2,395	1,273	935	1,112
1947	55,047	1,982	651	567	1,678	1,145	454	2,141	1,342	983	1,086
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	52,678	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	52,693	2,054	640	559	1,388	997	452	2,241	1,182	845	979
1941	51,688	2,069	618	536	1,427	919	442	2,288	1,143	906	1,017
1940	50,676	1,897	638	494	1,368	862	459	2,144	1,086	868	995
1939	50,898	1,981	616	492	1,419	937	436	2,148	1,152	872	986
1938	50,218	1,917	644	493	1,362	956	442	2,048	1,112	824	986
1937	50,419	1,860	647	512	1,414	979	457	2,134	1,120	853	1,001
1936	48,989	1,810	614	485	1,377	936	400	1,933	1,037	857	915
1935	48,856	1,846	623	508	1,267	988	464	1,979	1,107	828	919
1934	47,732	1,771	621	516	1,347	883	428	1,892	1,012	851	878
1933	46,406	1,729	567	486	1,316	905	420	1,828	983	847	895
1932	45,030	1,666	510	475	1,284	807	389	1,777	1,031	769	870
1931	44,306	1,633	555	413	1,247	864	488	1,766	979	795	841
1930	44,687	1,645	539	436	1,221	851	424	1,785	1,031	822	831
1929	43,492	1,515	485	424	1,222	870	352	1,746	977	729	842
1928	44,267	1,584	492	435	1,255	914	391	1,728	1,035	755	801
1927	42,627	1,438	501	429	1,243	827	393	1,636	1,038	735	818
1926	42,333	1,448	514	438	1,205	845	409	1,676	1,017	730	794
1925	41,505	1,384	480	376	1,209	844	414	1,664	941	775	794
1924	41,058	1,334	508	347	1,238	828	417	1,539	974	706	819
1923	41,587	1,349	510	430	1,352	825	414	1,553	962	775	792
1922	40,578	1,325	567	420	1,217	780	990	1,536	960	726	766
1921	40,351	1,367	509	351	1,156	837	430	1,540	922	724	677
1920	40,102	1,267	512	345	1,217	799	397	1,624	967	713	714

TABLE 17 - DEATHS FROM TEN SELECTED CAUSES BY PREFECTURE: JAPAN, 1920-1949 Cont'd
CANCER AND OTHER MALIGNANT TUMORS (45-55) Cont'd

Year	Hiroshima	Hokkaido	Hyogo	Ibaraki	Ishikawa	Iwate	Kagawa	Kagoshima	Kanagawa	Kochi	Kumamoto	Kyoto
1949	1,557	2,445	2,270	1,570	805	635	769	997	1,586	610	1,223	1,420
1948	1,479	2,211	2,126	1,535	772	590	762	918	1,423	606	1,161	1,360
1947	1,380	2,119	1,964	1,520	713	549	737	956	1,314	582	1,104	1,401
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	1,424	1,729	2,136	1,381	675	557	676	987	1,388	549	1,005	1,514
1941	1,302	1,723	2,173	1,312	661	549	663	906	1,245	527	986	1,408
1940	1,310	1,648	2,093	1,308	686	534	648	897	1,245	469	990	1,412
1939	1,340	1,713	2,055	1,348	668	510	599	925	1,288	506	943	1,389
1938	1,277	1,700	2,080	1,355	609	543	629	900	1,254	513	965	1,383
1937	1,282	1,678	2,011	1,287	688	533	613	948	1,155	554	1,005	1,413
1936	1,278	1,641	2,028	1,265	619	438	564	881	1,226	532	976	1,323
1935	1,319	1,608	2,020	1,309	649	500	534	931	1,100	480	894	1,349
1934	1,284	1,499	1,923	1,312	686	484	618	863	1,168	468	944	1,287
1933	1,310	1,429	1,932	1,207	637	477	602	845	1,111	500	951	1,368
1932	1,149	1,372	1,809	1,275	613	480	564	890	991	507	897	1,234
1931	1,145	1,320	1,754	1,199	627	433	590	801	1,029	460	874	1,229
1930	1,172	1,391	1,774	1,339	618	420	532	847	962	443	870	1,245
1929	1,115	1,311	1,727	1,224	624	459	572	811	983	477	898	1,188
1928	1,181	1,315	1,772	1,177	648	461	542	783	994	446	941	1,220
1927	1,146	1,249	1,713	1,223	591	425	519	732	957	490	879	1,187
1926	1,125	1,179	1,693	1,173	578	359	542	726	936	487	893	1,161
1925	1,087	1,209	1,709	1,191	598	413	517	790	806	460	855	1,162
1924	1,138	1,120	1,648	1,234	612	400	478	752	832	423	925	1,159
1923	1,109	1,130	1,623	1,179	669	388	493	765	917	486	928	1,175
1922	1,118	1,167	1,507	1,143	606	388	528	724	886	478	893	1,104
1921	1,147	1,104	1,693	1,111	596	397	504	675	904	390	885	1,172
1920	1,129	1,059	1,545	1,164	578	398	512	676	866	450	893	1,130

TABLE 17 - 1/ DEATHS FROM TEN SELECTED CAUSES BY PREFECTURE: JAPAN, 1920-49 Cont'd.
CANCER AND OTHER MALIGNANT TUMORS (45-55) Cont'd.

Area	Mie	Miyagi	Miyazaki	Nagano	Nagasaki	Nara	Niigata	Oita	Okayama	Osaka	Saga	Saitama
1949	1,098	1,205	658	1,704	1,050	793	2,075	885	1,308	2,771	754	1,652
1948	1,056	1,123	613	1,630	957	790	2,043	860	1,354	2,458	797	1,533
1947	948	1,005	641	1,515	920	727	1,928	863	1,291	2,423	728	1,506
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	983	978	568	1,434	879	716	1,908	702	1,092	3,292	587	1,368
1941	1,001	979	528	1,446	833	742	1,829	657	1,140	3,185	569	1,274
1940	917	1,031	497	1,486	842	715	1,760	676	1,107	3,165	628	1,298
1939	1,027	974	512	1,422	818	753	1,813	732	1,138	3,200	589	1,299
1938	1,206	929	518	1,430	842	726	1,749	732	1,140	3,034	577	1,350
1937	985	923	542	1,462	844	742	1,774	724	1,162	3,099	633	1,350
1936	979	916	498	1,404	830	761	1,778	710	1,111	2,973	612	1,303
1935	994	1,001	485	1,461	832	759	1,755	710	1,097	3,044	551	1,294
1934	954	892	500	1,405	806	789	1,738	668	1,086	2,764	580	1,229
1933	1,009	881	458	1,356	742	772	1,690	692	1,011	2,783	606	1,154
1932	931	821	482	1,318	764	711	1,681	667	987	2,721	574	1,235
1931	931	801	482	1,299	730	770	1,645	651	998	2,623	615	1,169
1930	925	781	526	1,306	783	731	1,619	676	1,035	2,558	603	1,230
1929	980	777	477	1,380	769	708	1,572	564	1,005	2,539	553	1,151
1928	1,010	827	478	1,366	741	713	1,623	669	998	2,448	604	1,224
1927	945	803	439	1,293	755	742	1,610	605	975	2,413	542	1,192
1926	936	799	473	1,241	718	751	1,593	656	991	2,326	617	1,208
1925	928	745	429	1,229	687	732	1,596	626	889	2,230	592	1,113
1924	908	804	456	1,244	715	774	1,606	636	967	2,128	561	1,184
1923	942	779	435	1,251	735	782	1,553	624	976	2,054	568	1,165
1922	935	744	429	1,222	680	697	1,618	611	915	2,128	582	1,044
1921	907	771	387	1,164	776	740	1,582	569	920	2,130	566	1,128
1920	927	764	388	1,148	728	760	1,581	619	929	2,135	608	1,106

TABLE 17 - 1/ DEATHS FROM TEN SELECTED CAUSES BY PREFECTURE: JAPAN, 1920-49 Cont'd
CANCER AND OTHER MALIGNANT TUMORS (45-55) - Cont'd.

Year	Shiga	Shimane	Shizuoka	Tochigi	Tokushima	Tokyo	Tottori	Toyama	Wakayama	Yamagata	Yamaguchi	Yamanshi
1949	679	708	1,418	1,189	646	3,980	505	837	859	1,075	1,081	617
1948	639	694	1,359	1,133	570	3,584	498	805	806	979	1,012	584
1947	619	646	1,367	1,152	594	3,203	507	690	793	967	1,057	579
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	613	522	1,170	1,000	535	4,609	440	684	862	888	925	580
1941	612	542	1,219	992	561	4,430	449	644	842	836	872	566
1940	640	552	1,184	990	505	4,346	452	675	863	891	857	548
1939	624	603	1,177	960	501	4,234	373	679	823	879	906	539
1938	638	595	1,160	935	497	4,085	471	614	860	849	931	539
1937	591	558	1,099	1,030	501	4,036	428	680	803	852	930	527
1936	626	589	1,109	901	517	3,990	449	627	744	832	940	505
1935	619	615	1,139	910	495	3,785	412	587	767	860	909	493
1934	677	561	1,130	890	506	3,715	410	640	846	818	886	507
1933	670	545	1,056	878	482	3,523	364	606	782	801	896	504
1932	556	552	1,034	906	481	3,336	403	586	726	792	898	509
1931	601	534	1,012	820	431	3,263	398	601	781	811	826	452
1930	634	592	1,003	807	474	3,221	355	637	803	849	851	485
1929	561	599	985	891	461	3,067	416	628	757	803	844	460
1928	611	619	1,014	860	482	3,052	356	648	825	816	919	497
1927	623	580	964	794	484	2,822	384	588	799	784	857	435
1926	588	597	968	796	439	2,895	400	576	751	774	863	459
1925	623	569	956	859	477	2,735	425	597	737	777	861	415
1924	611	615	899	764	455	2,500	408	592	711	762	884	443
1923	603	614	987	772	486	2,447	430	632	764	790	925	429
1922	577	558	897	795	444	2,675	391	575	735	770	919	440
1921	639	628	903	778	432	2,531	392	557	724	726	956	455
1920	632	591	863	759	456	2,463	375	524	698	738	877	463

See footnotes at end of table

TABLE 17. - 1/ DEATHS FROM TEN SELECTED CAUSES BY PREFECTURE, JAPAN, 1920-1949 Cont'd
3/ INTRACRANIAL LESIONS OF VASCULAR ORIGIN (83)

Area	All Japan	Aichi	Akita	Aomori	Obiba	Ehime	Fukui	Fukuoka	Fukushima	Gifu	Gumma
1949	100,650	3,906	2,476	1,712	3,340	1,614	902	3,594	3,019	1,918	2,201
1948	94,329	3,650	2,506	1,773	3,285	1,454	800	3,413	2,956	1,722	2,087
1947	101,095	3,969	2,648	1,665	3,326	1,745	932	3,833	3,181	1,902	2,071
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	120,985	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	125,001	4,357	3,339	1,756	3,506	1,809	1,106	5,004	3,591	2,230	2,368
1941	124,786	4,466	3,245	1,952	3,822	1,822	1,080	5,069	3,647	2,241	2,404
1940	127,523	4,438	3,412	1,906	3,600	1,948	1,186	5,107	3,661	2,242	2,372
1939	130,496	4,296	3,328	1,912	3,508	2,115	1,164	5,536	3,682	2,269	2,406
1938	126,540	4,471	3,191	1,856	3,570	1,900	1,081	5,403	3,519	2,200	2,337
1937	118,456	3,985	2,962	1,640	3,237	1,890	1,106	5,216	3,375	2,150	2,089
1936	117,847	3,984	2,897	1,559	3,400	1,845	1,076	5,052	3,249	2,194	2,113
1935	114,268	3,794	2,809	1,518	3,278	1,811	989	4,859	3,244	2,049	2,055
1934	114,192	3,890	2,893	1,468	3,282	1,798	1,050	4,687	3,370	2,099	2,028
1933	110,441	3,591	2,818	1,395	3,318	1,848	1,021	4,387	3,195	1,955	1,992
1932	107,176	3,550	2,801	1,301	3,243	1,852	963	4,285	3,176	1,899	1,928
1931	107,150	3,564	2,677	1,411	3,304	1,790	967	4,113	3,255	1,829	2,032
1930	104,747	3,252	2,663	1,323	3,194	1,769	903	4,250	3,189	1,811	1,948
1929	108,202	3,518	2,976	1,403	3,373	1,776	906	4,281	3,233	1,936	1,993
1928	102,722	3,432	2,834	1,240	3,298	1,728	851	4,001	3,023	1,797	1,896
1927	101,502	3,197	2,744	1,197	3,136	1,692	810	3,961	3,128	1,808	1,888
1926	98,486	3,145	2,759	1,211	3,185	1,702	862	3,747	3,051	1,785	1,805
1925	96,085	3,210	2,636	1,181	3,069	1,571	809	3,593	3,088	1,797	1,825
1924	102,584	3,012	2,888	1,309	3,168	1,768	894	3,872	3,239	1,841	1,980
1923	94,406	3,050	2,548	1,071	3,244	1,547	816	3,619	3,047	1,722	1,729
1922	91,244	2,740	2,577	1,222	3,021	1,476	762	3,588	3,077	1,603	1,683
1921	90,312	3,003	2,405	1,075	2,964	1,557	819	3,494	3,044	1,611	1,602
1920	87,984	2,923	2,374	1,057	2,789	1,520	721	3,667	2,961	1,630	1,517

TABLE 17. - 1/ DEATHS FROM TEN SELECTED CAUSES BY PREFECTURE: JAPAN, 1920-1949 Cont'd
3/ INTRACRANIAL LESIONS OF VASCULAR ORIGIN (83) - Cont'd

Year	Hiroshima	Hokkaido	Hyogo	Ibataki	Ishikawa	Iwate	Kagawa	Kagoshima	Kanagawa	Kochi	Kumamoto	Kyoto
1949	2,415	3,670	3,112	3,348	1,235	2,407	1,116	2,324	2,514	1,284	2,258	1,732
1948	2,192	3,230	2,952	3,193	1,077	2,373	1,004	2,144	2,300	1,161	2,074	1,554
1947	2,494	3,175	3,314	3,244	1,230	2,424	1,138	2,265	2,289	1,311	2,275	1,899
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	3,178	3,893	4,473	3,546	1,350	2,661	1,290	2,827	3,277	1,420	2,773	2,446
1941	3,144	3,956	4,333	3,702	1,496	2,768	1,254	2,654	3,269	1,316	2,588	2,329
1940	3,286	4,084	4,394	3,590	1,542	2,712	1,334	2,753	3,183	1,327	2,676	2,505
1939	3,510	4,306	4,624	3,652	1,483	2,644	1,411	2,661	3,295	1,432	2,707	2,648
1938	3,462	3,765	4,482	3,465	1,559	2,623	1,360	2,611	3,088	1,387	2,807	2,526
1937	3,070	3,680	4,398	3,230	1,540	2,521	1,250	2,488	2,912	1,361	2,663	2,387
1936	3,146	3,275	4,142	3,275	1,536	2,442	1,260	2,530	2,814	1,390	2,499	2,371
1935	2,915	3,286	4,107	3,286	1,417	2,347	1,250	2,365	2,754	1,431	2,562	2,370
1934	2,898	3,091	3,950	3,462	1,490	2,417	1,173	2,489	2,746	1,353	2,413	2,414
1933	2,893	3,040	3,996	3,364	1,378	2,237	1,226	2,259	2,534	1,311	2,398	2,344
1932	2,977	2,724	3,660	3,287	1,363	2,243	1,164	2,307	2,488	1,252	2,268	2,240
1931	2,813	2,861	3,612	3,519	1,320	2,241	1,179	2,254	2,464	1,213	2,304	2,536
1930	2,978	2,818	3,579	3,287	1,273	2,295	1,108	2,474	2,353	1,274	2,301	2,169
1929	3,056	2,715	3,728	3,446	1,329	2,236	1,267	2,276	2,473	1,386	2,292	2,357
1928	2,835	2,529	3,547	3,342	1,246	2,292	1,072	2,148	2,336	1,252	2,119	2,203
1927	2,871	2,475	3,624	3,253	1,386	2,294	1,230	2,207	2,285	1,288	2,102	2,232
1926	2,923	2,400	3,559	3,184	1,328	2,241	1,059	2,063	2,200	1,298	2,117	2,073
1925	2,603	2,374	3,254	3,255	1,230	2,339	944	2,098	2,202	1,201	2,116	2,089
1924	3,144	2,310	3,767	3,335	1,529	2,451	1,331	2,209	1,927	1,378	2,100	2,198
1923	2,607	2,157	3,235	3,206	1,255	2,275	1,003	2,097	2,131	1,229	2,009	1,970
1922	2,546	1,972	3,105	3,024	1,362	2,312	1,266	2,106	1,946	1,165	1,878	1,837
1921	2,505	1,888	3,017	2,983	1,231	2,194	1,100	2,091	1,962	1,183	2,002	1,965
1920	2,459	1,922	3,054	3,050	1,131	2,103	1,007	2,125	1,886	1,130	1,893	1,857

TABLE 17. - 1/ ILLATES FROM TEN SELECTED CAUSES BY HEMISPHERE: JAPAN, 1920-1949 Cont'd
3/ INTRACRANIAL LESIONS OF VASCULAR ORIGIN (83) - Cont'd

Year	Hio	Miyagi	Miyazaki	Nagano	Nagasaki	Nara	Miigata	Oita	Okayama	Osaka	Saga	Saitama
1949	1,768	2,284	1,279	3,223	1,660	960	3,985	1,697	2,077	3,096	1,217	3,004
1948	1,650	2,448	1,102	3,080	1,503	978	3,947	1,578	2,002	2,791	1,074	2,898
1947	1,868	2,166	1,236	3,216	1,692	1,073	4,321	1,808	2,288	3,232	1,206	2,822
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	2,156	2,561	1,472	3,998	1,849	1,248	4,849	2,051	2,618	5,552	1,420	3,030
1941	2,448	2,491	1,494	3,875	1,866	1,230	4,471	2,022	2,550	5,288	1,322	3,240
1940	2,079	2,570	1,475	4,054	1,946	1,251	4,744	1,985	2,634	5,737	1,361	3,259
1939	2,439	2,472	1,535	4,131	2,440	1,373	4,826	2,446	2,789	6,173	1,334	3,217
1938	2,085	2,403	1,466	3,906	2,055	1,273	4,746	2,084	2,667	5,681	1,361	3,241
1937	1,986	2,361	1,343	3,051	1,912	1,186	4,411	1,917	2,647	5,323	1,379	2,931
1936	1,974	2,293	1,398	3,777	1,837	1,188	4,304	1,956	2,572	5,456	1,320	2,926
1935	1,921	2,191	1,357	3,740	1,807	1,224	4,179	1,931	2,538	5,226	1,250	2,917
1934	1,937	2,255	1,217	3,826	1,629	1,210	4,481	1,960	2,445	5,108	1,290	3,092
1933	1,615	2,167	1,293	3,694	1,586	1,141	4,258	1,849	2,373	4,992	1,299	2,895
1932	1,883	2,120	1,298	3,686	1,535	1,164	3,959	1,914	2,343	4,595	1,209	2,802
1931	1,234	2,190	1,234	3,718	1,527	1,174	3,929	1,694	2,418	4,552	1,266	2,925
1930	1,721	2,252	1,258	3,479	1,534	1,086	3,739	1,783	2,371	4,544	1,214	2,624
1929	1,758	2,234	1,134	3,763	1,465	1,153	4,080	1,856	2,467	4,546	1,283	2,815
1928	1,940	2,209	1,073	3,540	1,510	1,033	3,763	1,696	2,344	4,349	1,202	2,719
1927	1,721	2,165	1,148	3,446	1,488	1,006	3,871	1,761	2,487	4,443	1,195	2,565
1926	1,672	2,059	1,077	3,385	1,359	973	3,934	1,627	2,370	4,086	1,194	2,509
1925	1,703	2,211	1,116	3,377	1,334	988	3,767	1,600	2,217	3,772	1,130	2,504
1924	1,638	2,293	1,155	3,716	1,391	1,023	4,440	1,664	2,657	3,892	1,230	2,461
1923	1,624	2,193	1,074	3,374	1,371	941	3,620	1,531	2,270	3,516	1,173	2,333
1922	1,423	2,171	1,100	3,267	1,272	933	3,527	1,494	2,292	3,285	1,226	2,166
1921	1,571	2,442	1,110	3,195	1,235	982	3,507	1,669	2,223	3,369	1,145	2,130
1920	1,512	2,049	1,003	3,078	1,136	930	3,567	1,565	2,157	3,217	1,163	2,038

TABLE 17. - 1/ DEATHS FROM TEN SELECTED CAUSES BY PREFECTURE, JAPAN, 1920-1949, Cont'd
3/ INTRACRANIAL LESIONS OF VASCULAR ORIGIN (83) (Cont'd)

Year	Shiga	Shimane	Shizuoka	Tochigi	Tokushima	Tokyo	Tottori	Toyama	Wakayama	Yamagata	Yamaguchi	Yamanehi
1949	1,083	1,376	2,883	2,271	962	5,241	813	1,307	1,173	2,382	1,825	1,027
1948	1,029	1,175	2,626	2,191	926	4,688	773	1,271	1,072	2,296	1,737	974
1947	1,225	1,332	2,302	2,084	1,156	4,309	811	1,298	1,174	2,470	2,155	1,021
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	1,299	1,477	3,352	2,603	1,154	9,390	958	1,666	1,684	2,661	2,575	1,148
1941	1,338	1,700	3,408	2,613	1,194	9,351	992	1,522	1,564	2,850	2,515	1,205
1940	1,372	1,685	3,416	2,637	1,199	9,711	994	1,715	1,640	2,954	2,607	1,230
1939	1,415	1,782	3,325	2,518	1,289	10,010	1,162	1,606	1,711	2,804	2,781	1,229
1938	1,349	1,791	3,359	2,593	1,210	9,549	1,044	1,643	1,612	2,784	2,863	1,202
1937	1,320	1,612	3,107	2,266	1,163	8,483	956	1,531	1,535	2,538	2,607	1,061
1936	1,401	1,675	3,150	2,340	1,229	8,547	938	1,553	1,490	2,565	2,574	1,146
1935	1,285	1,514	2,978	2,215	1,258	8,221	959	1,452	1,562	2,481	2,418	1,158
1934	1,430	1,632	2,983	2,300	1,157	7,920	937	1,498	1,513	2,528	2,327	1,056
1933	1,230	1,604	2,892	2,298	1,151	7,776	923	1,415	1,460	2,489	2,291	1,119
1932	1,277	1,624	2,827	2,197	1,083	7,396	892	1,407	1,316	2,419	2,265	1,054
1931	1,295	1,457	2,827	2,238	1,119	7,515	862	1,469	1,346	2,456	2,167	977
1930	1,279	1,555	2,655	2,151	1,027	7,138	824	1,288	1,399	2,373	2,240	1,002
1929	1,352	1,598	2,768	2,270	1,174	7,211	895	1,415	1,353	2,366	2,268	1,022
1928	1,308	1,428	2,708	2,199	1,031	6,860	877	1,294	1,343	2,322	2,124	979
1927	1,215	1,406	2,497	2,119	1,154	6,559	896	1,260	1,280	2,304	2,134	934
1926	1,235	1,360	2,547	2,094	1,008	6,072	763	1,221	1,209	2,171	2,070	914
1925	1,175	1,319	2,438	2,030	974	5,928	702	1,167	1,160	2,145	2,064	840
1924	1,298	1,544	2,545	2,068	1,167	5,687	877	1,561	1,330	2,171	2,440	946
1923	1,245	1,518	2,434	1,889	947	5,541	780	1,301	1,177	1,973	2,204	810
1922	1,123	1,263	2,311	1,823	1,005	5,281	711	1,360	1,094	1,990	2,098	811
1921	1,201	1,166	2,324	1,817	1,010	5,027	707	1,332	1,201	1,776	2,009	770
1920	1,146	1,140	2,281	1,880	901	4,875	646	1,266	1,098	1,838	1,960	742

See footnotes at end of table.

TABLE 17. - 1/ DEATHS FROM TEN SELECTED CAUSES BY PREFECTURE: JAPAN, 1920-1949 Cont'd
 1/ DISEASES OF THE HEART (90 - 95)

Year	All Japan	Aichi	Akita	Iomori	Chiba	Shizu	Fukui	Fukuoka	Fukushima	Gifu	Gunma
1949	52,825	2,198	911	662	1,642	969	520	1,847	1,337	1,051	1,236
1948	49,046	1,993	872	646	1,566	955	430	1,632	1,250	980	1,130
1947	48,575	2,051	831	564	1,455	908	418	1,678	1,272	1,137	1,089
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	45,428	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	43,255	1,828	668	507	1,265	713	453	1,470	1,017	843	976
1941	42,280	1,768	704	556	1,253	675	475	1,467	983	818	923
1940	45,275	1,883	796	540	1,367	773	526	1,610	1,003	904	894
1939	47,166	1,910	713	588	1,401	760	530	1,721	1,123	866	896
1938	47,192	1,955	670	517	1,492	821	543	1,753	1,096	963	943
1937	42,569	1,797	655	459	1,323	687	541	1,533	908	850	837
1936	42,650	1,735	612	613	1,184	764	469	1,536	962	916	835
1935	39,628	1,581	609	523	1,194	716	491	1,317	934	837	706
1934	42,283	1,666	668	543	1,217	782	580	1,375	1,029	981	834
1933	39,897	1,508	666	438	1,222	714	554	1,287	925	850	724
1932	38,784	1,475	636	457	1,196	660	519	1,329	941	851	711
1931	41,697	1,738	636	489	1,210	788	562	1,397	956	906	718
1930	40,956	1,433	688	489	1,234	698	566	1,406	986	982	742
1929	41,342	1,550	861	580	1,248	691	543	1,250	997	889	720
1928	39,711	1,619	680	492	1,236	729	493	1,219	920	936	784
1927	38,786	1,474	711	449	1,113	741	512	1,230	874	851	737
1926	37,796	1,446	725	543	1,181	687	485	1,147	845	773	664
1925	39,731	1,548	761	507	1,181	738	475	1,198	908	858	738
1924	40,485	1,525	737	540	1,238	755	456	1,218	1,025	872	747
1923	41,989	1,630	1,030	609	1,311	797	529	1,197	1,014	944	736
1922	37,145	1,327	787	560	1,071	674	511	1,102	960	845	740
1921	37,279	1,504	702	480	1,146	738	454	1,129	872	896	684
1920	35,401	1,294	669	469	1,031	653	409	1,067	867	767	624

TABLE 17. - 1/ DEATHS FROM TEN SELECTED CAUSES BY PREFECTURE: JAPAN, 1920-49 Cont'd
1/ DISEASES OF THE HEART (90 - 95) - Cont'd.

Year	Hiroshima	Hokkaido	Hyogo	Ibaraki	Ishikawa	Iwate	Kagawa	Kagoshima	Kanagawa	Kochi	Kumamoto	Kyoto
1919	1,322	2,250	1,860	1,629	711	885	624	1,122	1,263	456	1,239	1,054
1918	1,370	1,996	1,660	1,500	612	882	586	1,034	1,159	408	1,096	1,053
1917	1,392	1,773	1,813	1,410	666	740	593	1,022	1,099	431	1,152	1,087
1916	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1915	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1914	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1913	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1912	1,251	1,412	1,594	1,185	526	745	435	912	1,181	381	847	999
1911	1,145	1,551	1,543	1,135	520	742	408	859	1,132	331	845	926
1910	1,322	1,738	1,654	1,179	645	791	434	923	1,141	390	919	1,101
1909	1,290	1,895	1,792	1,202	653	848	559	891	1,137	388	890	1,061
1908	1,358	1,692	1,851	1,169	690	722	540	958	1,132	389	1,027	1,043
1907	1,226	1,575	1,630	1,043	663	694	492	831	1,000	334	892	992
1906	1,219	1,583	1,633	1,012	683	695	490	884	1,020	396	939	920
1905	1,129	1,405	1,485	944	675	602	488	807	999	402	856	867
1904	1,177	1,421	1,627	1,009	735	648	561	856	1,052	399	872	1,060
1903	1,182	1,437	1,563	1,026	635	654	547	774	1,003	377	832	890
1902	1,133	1,341	1,620	1,029	666	559	449	787	943	342	785	909
1901	1,217	1,414	1,592	1,070	656	635	544	792	1,013	369	924	1,007
1900	1,245	1,357	1,601	956	695	650	507	868	958	414	828	988
1909	1,259	1,536	1,633	1,037	761	611	599	771	990	374	865	1,016
1908	1,204	1,234	1,583	1,040	717	601	534	806	926	430	773	942
1907	1,220	1,235	1,538	982	698	615	526	726	904	444	772	955
1906	1,231	1,228	1,465	929	697	601	501	866	873	447	816	943
1905	1,236	1,190	1,684	1,038	744	655	512	743	933	448	743	1,061
1904	1,454	1,178	1,584	1,063	789	611	546	855	895	485	817	1,067
1903	1,415	1,262	1,639	1,002	885	627	523	894	964	446	782	1,113
1902	1,259	1,427	1,427	860	790	525	468	801	852	460	691	1,026
1901	1,274	1,020	1,410	905	711	528	467	824	884	431	812	972
1900	1,209	1,010	1,377	892	642	475	429	804	750	423	716	971

TABLE 17. - 1/ DEATHS FROM TEN SELECTED CAUSES BY PREFECTURE, JAPAN, 1920-1949 Cont'd
1/ DISEASES OF THE HEART (90 - 95) - Cont'd

Year	Mie	Miyagi	Miyazaki	Nagano	Nagasaki	Nara	Niigata	Oita	Okayama	Osaka	Saga	Saitama
1949	1,038	1,086	769	1,604	995	520	1,651	910	1,077	2,146	616	1,607
1948	992	941	581	1,438	951	506	1,601	811	1,002	1,978	632	1,400
1947	962	932	620	1,415	1,022	553	1,563	794	963	2,071	623	1,332
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	865	889	468	1,252	842	445	1,428	675	749	2,389	456	1,084
1941	815	850	456	1,180	805	418	1,429	620	773	2,127	452	1,041
1940	854	811	458	1,298	856	475	1,461	628	819	2,381	490	1,113
1939	845	859	573	1,265	926	484	1,539	690	975	2,511	508	1,137
1938	987	798	562	1,299	935	433	1,349	701	885	2,493	494	1,119
1937	842	749	486	1,145	827	440	1,302	699	839	2,301	412	993
1936	821	824	487	1,215	865	429	1,412	678	847	2,242	470	979
1935	764	738	459	1,076	718	435	1,287	615	814	2,101	447	943
1934	885	751	430	1,187	725	487	1,444	661	859	2,248	446	993
1933	765	709	443	1,123	608	405	1,223	643	757	2,209	359	999
1932	751	699	436	1,049	668	438	1,204	635	798	2,014	427	942
1931	929	688	520	1,142	709	454	1,357	670	831	2,164	410	978
1930	809	743	460	1,148	714	443	1,331	664	814	2,207	466	892
1929	822	706	484	1,134	706	426	1,487	652	839	1,999	435	925
1928	905	685	457	1,110	667	402	1,310	590	805	2,018	421	852
1927	823	730	403	1,081	643	384	1,292	662	829	1,941	358	898
1926	788	631	413	1,040	603	398	1,377	569	764	1,835	386	878
1925	827	684	409	1,178	616	441	1,451	530	826	1,976	421	917
1924	839	716	425	1,220	641	442	1,493	580	869	2,010	427	951
1923	952	761	424	1,195	695	351	1,507	606	963	2,015	436	990
1922	769	683	375	1,101	618	338	1,328	597	852	1,889	426	782
1921	821	669	421	1,118	617	382	1,218	633	856	1,865	408	846
1920	720	668	390	1,077	653	343	1,262	622	822	1,815	396	831

TABLE 17. - $\frac{1}{2}$ DEATHS FROM TEN SELECTED CAUSES BY PREFECTURE, JAPAN, 1920-49 Cont'd
 $\frac{1}{2}$ DISEASES OF THE HEART (90 - 95) - Cont'd

Year	Shiga	Shimane	Shizuoka	Tochigi	Tokushima	Tokyo	Tottori	Toyama	Wakayama	Yamagata	Yamaguchi	Yamanashi
1949	657	743	1,477	1,085	682	3,131	381	700	597	995	973	617
1948	649	719	1,437	1,053	691	2,631	368	682	567	968	910	628
1947	722	675	1,472	1,022	789	2,275	374	636	606	949	1,048	576
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	583	552	1,182	815	568	3,251	305	532	530	829	816	492
1941	552	553	1,187	864	565	3,416	318	512	471	854	775	488
1940	581	623	1,234	836	623	3,465	313	627	549	880	838	479
1939	622	623	1,308	895	658	3,637	361	606	599	860	965	501
1938	668	628	1,382	858	618	3,644	390	715	594	832	988	496
1937	549	606	1,218	785	573	3,233	339	656	502	733	928	452
1936	623	580	1,200	689	637	3,005	360	609	495	727	864	492
1935	543	538	1,179	685	554	2,740	310	590	510	715	831	449
1934	650	589	1,115	705	633	2,818	344	651	533	723	851	482
1933	564	629	1,194	746	609	2,775	303	576	516	682	828	424
1932	558	641	1,140	666	584	2,514	311	569	498	628	862	404
1931	587	658	1,162	721	710	2,763	301	635	533	741	916	445
1930	607	704	1,166	718	594	2,663	316	629	527	728	955	429
1929	568	673	1,180	686	610	2,763	340	677	508	749	940	424
1928	566	621	1,122	672	651	2,603	320	585	515	676	816	444
1927	514	677	1,101	601	626	2,491	361	656	500	642	793	389
1926	542	668	1,098	600	547	2,340	300	624	462	724	764	378
1925	550	704	1,122	714	617	2,352	373	613	468	758	766	419
1924	509	682	1,146	736	614	2,258	354	639	487	784	813	393
1923	564	784	1,161	729	638	2,388	357	678	470	784	821	371
1922	503	544	983	637	616	2,186	321	607	440	654	771	382
1921	481	579	1,071	619	630	2,111	323	603	453	642	780	350
1920	478	591	966	641	571	2,037	262	546	411	650	701	390

See footnotes at end of table.

TABLE 17. - 1/ DEATHS FROM TEN SELECTED CAUSES BY PREFECTURE: JAPAN, 1920-1949 Cont'd
2/ PNEUMONIA, ALL FORMS (107-109)

Year	All Japan	Aichi	Akita	Aomori	Chiba	Ehime	Fukui	Fukuoka	Fukushima	Gifu	Gumma
1949	54,636	2,258	832	1,149	1,568	1,082	490	2,508	1,792	981	1,342
1948	52,979	1,804	849	1,180	1,309	973	445	2,003	1,392	901	1,234
1947	501,601	3,996	1,337	1,604	2,481	1,257	883	4,269	2,577	1,914	2,078
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	116,494	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	105,110	4,247	1,522	2,076	2,503	1,503	908	4,725	2,488	1,529	1,904
1941	103,060	5,087	1,569	1,874	2,417	1,593	1,214	4,804	2,262	1,953	2,076
1940	109,620	5,071	1,694	1,976	2,507	1,821	1,150	4,661	2,587	1,858	2,071
1939	130,246	4,862	1,970	2,288	2,939	2,167	1,280	6,755	3,038	1,858	2,554
1938	117,162	5,466	1,511	1,852	2,782	1,885	1,303	6,289	2,826	2,059	2,091
1937	107,162	5,002	1,460	2,007	2,274	1,787	1,353	5,465	2,597	1,817	2,167
1936	111,093	4,718	1,715	2,043	2,172	2,088	1,301	6,461	2,657	1,988	1,833
1935	104,061	3,764	1,665	1,962	2,487	1,605	1,261	4,803	2,462	1,752	1,915
1934	122,980	6,643	1,459	2,028	2,648	2,226	1,349	5,797	2,583	2,438	2,559
1933	105,163	4,347	1,535	1,874	2,210	1,973	1,204	5,653	2,337	1,583	1,726
1932	111,644	4,990	1,742	1,842	2,503	1,976	1,352	5,274	2,711	1,752	1,922
1931	128,352	5,299	1,521	2,198	2,818	2,172	1,477	6,888	3,392	2,418	2,553
1930	100,071	4,656	1,503	1,964	2,035	1,928	1,174	5,163	2,453	1,775	1,754
1929	119,860	4,900	1,732	1,709	2,418	2,193	1,243	6,892	2,794	1,786	2,112
1928	124,706	5,714	1,552	2,053	2,921	2,011	1,402	5,900	3,272	2,676	2,790
1927	118,915	5,623	1,640	1,926	2,621	2,017	1,324	5,591	3,054	2,196	2,741
1926	107,035	4,498	1,613	1,658	2,199	2,223	1,099	6,178	2,698	1,682	1,740
1925	127,547	5,319	1,680	2,063	3,076	2,449	1,379	5,980	3,071	2,209	2,590
1924	122,331	5,198	1,744	1,829	2,796	2,628	1,292	5,908	2,929	2,104	2,367
1923	118,396	5,035	1,473	1,734	3,106	1,954	1,325	5,865	3,137	2,218	2,512
1922	111,699	4,369	1,586	1,965	2,651	1,766	1,308	5,212	2,881	1,949	2,622
1921	112,594	4,484	1,567	1,657	2,611	1,776	1,015	6,850	2,643	1,661	2,448
1920	173,244	7,268	2,223	2,851	4,324	2,941	1,780	8,904	4,937	3,070	3,371

TABLE 17. - DEATHS FROM TEN SELECTED CAUSES BY PREFECTURE, JAPAN, 1920-49 Cont'd
5/ PNEUMONIA, ALL FORMS (107-109) - Cont'd

Year	Hiroshima	Hokkaido	Hyogo	Ibaraki	Ishikawa	Iwate	Kagawa	Kagoshima	Kanagawa	Kochi	Kumamoto	Kyoto
1949	1,154	3,276	1,798	1,308	747	1,367	590	1,265	1,647	482	1,259	990
1948	1,245	3,191	1,791	1,188	785	1,393	675	1,557	1,441	599	1,299	926
1947	2,563	5,450	3,555	2,255	1,372	2,025	1,198	2,853	2,865	1,860	2,719	2,029
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	2,633	5,172	4,147	2,399	1,421	2,156	927	2,685	3,297	804	2,145	1,731
1941	2,443	4,601	3,828	2,581	1,375	1,905	999	2,254	2,865	1,142	2,287	1,965
1940	2,720	5,086	4,164	2,694	1,622	2,325	1,175	2,678	2,771	927	2,350	2,427
1939	3,495	5,159	5,693	3,074	1,919	1,989	1,302	3,143	3,537	1,127	2,577	2,501
1938	3,027	4,785	4,284	2,906	1,593	1,911	1,157	3,003	3,075	1,302	2,509	2,112
1937	2,686	4,715	4,221	2,245	1,593	1,908	1,014	2,482	2,515	1,096	2,506	2,041
1936	3,432	4,165	4,204	2,521	1,781	2,405	1,454	2,757	2,227	1,238	2,490	2,675
1935	2,813	4,382	3,927	2,603	1,893	1,866	1,030	2,575	2,595	900	2,216	1,782
1934	3,174	5,021	4,347	2,928	1,906	1,947	1,668	3,090	2,899	1,162	2,898	2,401
1933	2,670	4,831	3,866	2,458	1,587	1,934	1,264	2,650	2,457	1,195	2,380	2,221
1932	3,638	4,371	4,325	2,419	1,852	1,964	1,422	2,816	2,580	1,294	2,609	2,262
1931	3,399	4,662	4,874	3,108	1,874	1,994	1,545	3,263	2,593	1,187	3,144	2,735
1930	2,741	4,897	3,633	2,254	1,430	1,709	1,282	2,686	2,029	993	2,322	2,063
1929	2,896	4,616	4,955	2,492	1,978	1,719	1,365	2,879	3,032	1,440	2,931	2,459
1928	3,437	5,209	4,411	3,377	1,860	2,096	1,513	3,059	3,201	1,234	2,721	2,392
1927	3,152	5,496	4,262	2,669	1,803	1,882	1,378	2,765	2,606	1,105	2,610	2,230
1926	3,196	4,388	4,206	2,079	1,820	1,695	1,476	3,038	2,121	1,074	2,882	2,091
1925	3,492	5,357	4,699	3,483	1,896	1,875	1,295	3,243	2,952	1,415	2,655	2,398
1924	4,166	5,177	5,032	2,760	1,877	2,035	1,506	3,286	2,767	1,157	3,111	2,107
1923	3,672	5,445	4,065	3,331	1,727	1,607	1,431	2,912	2,527	1,201	2,318	2,429
1922	4,472	4,917	4,195	2,917	1,885	1,812	1,490	3,244	2,562	1,118	2,874	1,679
1921	3,069	4,096	2,669	1,464	1,464	1,413	1,495	3,210	2,793	930	2,965	2,616
1920	3,136	7,786	8,246	3,778	2,258	2,001	1,770	3,719	3,566	1,780	3,445	3,290

TABLE 17. - DEATHS FROM TEN SELECTED CAUSES BY PREFECTURE: JAPAN, 1920-49 Cont'd
5/ PNEUMONIA, ALL FORMS (107-109) - Cont'd.

Year	Mie	Miyagi	Miyazaki	Nagano	Nagasaki	Nara	Niigata	Oita	Okayama	Osaka	Saga	Saitama
1949	888	1,250	678	1,272	1,265	489	1,880	936	876	2,731	562	1,570
1948	913	1,252	724	1,202	1,176	402	1,856	852	1,054	2,063	551	1,320
1947	1,688	2,196	1,412	2,400	2,214	961	2,805	1,808	2,080	5,161	1,217	2,675
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	1,580	2,108	1,278	1,972	2,356	718	3,058	1,703	1,457	6,762	1,170	2,544
1942	1,710	1,928	1,105	2,114	2,137	746	3,019	1,663	1,467	6,526	1,063	2,631
1941	1,679	2,115	1,078	2,182	2,545	797	3,493	1,710	1,659	7,423	1,142	2,833
1940												
1939	2,163	2,245	1,789	2,584	2,726	970	3,585	2,119	2,277	10,310	1,472	3,246
1938	2,429	1,960	1,470	2,510	2,990	805	3,092	1,928	1,801	7,296	1,378	2,658
1937	1,900	1,957	1,280	2,073	2,464	918	3,109	2,162	1,629	7,392	1,319	2,555
1936	2,232	2,278	1,452	2,558	2,340	924	3,068	2,023	2,088	6,410	1,424	2,288
1935	1,916	2,206	1,414	2,291	2,044	953	3,348	1,626	1,835	6,658	1,030	2,669
1934	2,459	2,055	1,371	2,496	2,384	1,191	3,526	2,035	2,278	6,656	1,452	3,100
1933	1,867	2,230	1,408	1,868	2,347	830	2,972	1,862	1,828	6,240	1,287	2,535
1932	1,926	1,974	1,541	2,247	2,778	1,014	3,460	2,197	2,002	6,152	1,412	2,702
1931	2,647	2,175	1,426	2,740	2,536	974	3,868	2,256	1,993	7,271	1,492	3,409
1930	2,009	2,168	1,097	2,167	2,396	718	3,262	1,688	1,721	4,810	1,442	2,090
1929	2,188	1,945	1,815	2,338	2,644	1,075	3,469	2,553	2,018	7,886	1,677	2,915
1928	2,557	2,476	1,306	2,986	2,651	913	4,059	2,070	1,989	5,906	1,499	3,451
1927	2,225	2,354	1,129	2,625	2,162	1,211	3,620	1,878	2,055	7,443	1,385	3,172
1926	1,858	2,050	1,377	2,065	2,375	786	3,287	2,252	2,161	5,432	1,612	2,306
1925	3,035	2,393	1,387	3,157	2,342	1,204	3,783	1,884	1,887	7,635	1,556	3,258
1924	2,237	2,500	1,334	2,892	2,788	1,063	4,090	2,125	2,259	6,288	1,602	2,941
1923	2,006	2,354	1,141	2,601	2,129	1,015	3,553	2,031	2,062	6,697	1,645	3,205
1922	2,116	2,375	1,279	2,794	2,142	857	4,026	1,750	1,844	5,443	1,576	2,958
1921	1,970	2,109	1,193	2,300	2,732	947	3,560	2,039	1,895	6,523	1,738	2,943
1920	3,008	3,622	1,422	3,761	3,686	1,673	5,676	2,528	2,586	11,610	2,034	4,681

TABLE 17 - 1/ DEATHS FROM TEN SELECTED CAUSES BY PREFECTURE, JAPAN, 1920-49 Cont'd
5/ PNEUMONIA, ALL FORMS (107-109) - Cont'd

Year	Shiga	Shimane	Shizuoka	Tochigi	Tokushima	Tokyo	Tottori	Toyama	Wakayama	Yamagata	Yamaguchi	Yamanashi
1949	522	577	1,517	1,118	620	3,657	307	814	562	1,068	1,056	536
1948	511	674	1,381	879	749	3,105	355	771	581	973	1,008	447
1947	1,099	1,192	2,653	1,922	1,312	6,477	619	1,217	1,089	1,683	1,800	1,041
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	790	1,100	2,437	1,888	1,056	10,413	574	1,715	992	1,454	1,994	969
1941	862	1,053	2,390	1,747	1,091	9,363	660	1,579	845	1,579	1,696	874
1940	943	1,123	2,517	1,860	1,267	9,055	640	1,901	1,050	1,618	1,790	855
1939	1,066	1,680	2,766	2,402	1,658	10,548	733	1,996	1,378	1,879	2,398	1,035
1938	1,101	1,498	2,850	1,880	1,447	10,094	584	1,799	1,022	1,612	2,161	1,089
1937	962	1,106	2,417	1,750	1,411	8,198	633	1,701	962	1,725	1,769	879
1936	1,044	1,369	2,585	1,800	1,527	7,052	637	1,679	1,298	1,739	2,208	805
1935	866	1,216	2,714	1,896	1,289	7,502	609	1,775	1,170	1,945	1,956	1,095
1934	1,249	1,437	3,212	2,209	1,819	9,686	791	2,164	1,303	2,008	1,927	1,070
1933	933	1,178	2,812	2,047	1,517	8,246	614	1,708	948	1,740	1,841	920
1932	995	1,482	2,694	1,910	1,677	7,452	581	1,812	1,347	1,908	1,874	928
1931	1,473	1,662	3,065	2,566	1,906	10,067	788	2,000	1,392	1,957	2,258	1,136
1930	829	1,250	2,649	1,667	1,318	6,281	586	1,638	934	1,995	2,002	910
1929	1,159	1,514	2,681	1,963	1,619	8,803	784	2,103	1,344	1,900	2,161	975
1928	1,047	1,345	2,983	2,434	1,722	9,338	579	2,034	1,119	1,976	2,098	1,317
1927	1,084	1,377	2,697	2,376	1,649	8,485	666	1,942	1,186	2,015	2,024	1,074
1926	1,021	1,400	2,362	1,667	1,610	6,535	688	2,209	1,065	1,977	2,378	908
1925	1,236	1,348	3,673	2,649	1,812	9,483	807	1,820	1,490	2,137	2,058	1,217
1924	1,090	1,659	2,984	2,103	1,822	6,827	759	2,036	1,250	2,012	2,606	1,348
1923	1,341	1,363	2,555	2,406	1,857	8,376	688	2,261	1,032	2,041	1,956	1,057
1922	1,043	1,293	2,746	2,204	1,702	6,397	646	1,864	1,068	2,340	1,915	1,273
1921	915	1,104	2,735	2,210	1,585	7,775	552	1,696	1,180	1,947	2,286	1,009
1920	1,869	1,658	4,191	3,119	2,769	11,049	1,102	2,514	1,492	3,276	3,064	1,940

See footnotes at end of table.

TABLE 17. - 1/ DEATHS FROM TEN SELECTED CAUSES BY PREFECTURE, JAPAN, 1920-1949, Cont'd
2/ DIARRHEA, ENTERITIS AND ULCERATION OF THE INTESTINES (ALL AGES) (119-120)

Year	All Japan	Aichi	Akita	Aomori	Chiba	Ehime	Fukui	Fukuoka	Fukushima	Gifu	Gumma
1949	72,186	2,376	1,655	1,798	1,421	1,203	1,087	2,897	1,757	1,585	1,246
1948	83,264	2,477	2,344	2,513	1,772	1,250	1,456	2,598	2,127	1,717	1,645
1947	101,454	3,341	2,387	2,312	2,194	1,884	1,537	3,810	2,262	2,085	1,817
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	105,811	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	95,618	3,130	2,087	2,523	1,985	1,391	1,432	3,240	2,577	1,948	1,947
1941	95,353	4,151	2,081	2,397	2,159	1,393	1,633	2,834	2,530	2,194	2,061
1940	106,889	4,718	2,366	2,637	2,443	1,534	1,887	3,299	3,151	2,401	2,776
1939	111,232	5,017	2,367	2,829	2,324	1,713	1,798	3,411	3,043	2,424	2,629
1938	115,455	5,048	2,135	2,867	2,568	1,569	2,074	3,826	3,198	2,789	3,010
1937	118,460	5,181	2,776	2,950	2,411	1,678	2,180	3,847	3,508	2,594	2,842
1936	119,343	4,893	2,511	3,501	2,494	1,557	2,451	4,179	3,666	2,759	2,654
1935	110,599	4,350	2,408	2,723	2,357	1,577	1,817	3,623	3,121	2,539	2,313
1934	126,357	5,586	2,844	3,506	2,442	1,668	2,654	3,622	3,610	3,124	2,728
1933	129,202	4,860	2,824	2,800	2,954	2,102	2,299	4,217	3,540	2,771	2,698
1932	135,761	5,161	3,521	4,499	2,674	2,072	2,447	4,447	4,001	2,977	2,827
1931	138,409	5,135	3,195	3,325	3,025	2,001	2,401	5,236	3,319	3,212	3,012
1930	140,946	5,436	3,218	4,007	3,122	2,111	2,359	5,072	3,913	3,202	2,926
1929	154,993	6,277	3,422	3,879	3,674	2,227	2,679	5,793	4,573	3,490	3,693
1928	144,962	6,371	3,570	4,063	3,378	2,103	2,446	5,190	3,946	3,540	3,340
1927	140,171	5,093	3,207	3,613	3,158	2,416	2,692	4,793	4,237	3,317	3,835
1926	139,478	5,504	3,321	3,402	3,160	2,225	2,380	4,523	3,586	3,153	3,274
1925	140,335	5,899	3,193	3,506	2,916	2,020	2,369	4,372	3,507	3,383	3,759
1924	150,134	6,299	3,406	4,052	3,393	2,378	2,438	5,051	3,943	3,483	3,652
1923	162,660	6,704	3,925	3,928	4,114	2,357	2,763	4,997	4,222	3,619	4,174
1922	151,990	5,566	3,444	3,979	3,255	2,380	2,768	4,571	3,819	3,228	3,604
1921	153,126	6,869	3,159	3,491	3,422	2,631	2,703	4,779	3,753	3,629	3,475
1920	140,444	6,286	2,542	3,280	3,143	2,355	2,517	4,663	3,382	3,263	3,482

TABLE 17. - 1/ DEATHS FROM TEN SELECTED CAUSES BY PREFECTURE: JAPAN, 1920-1949, Cont'd
 5/ DIARRHEA, ENTERITIS AND ULCERATION OF THE INTESTINES (ALL AGES) (119-120) Cont'd

Year	Hiroshima	Hokkaido	Hyogo	Ibaraki	Ishikawa	Iwate	Kagawa	Kagoshima	Kanagawa	Kochi	Kumamoto	Kyoto
1949	1,665	4,933	2,942	1,680	1,167	1,592	704	1,815	1,094	639	1,684	1,282
1948	1,911	5,327	3,261	1,782	1,627	2,098	791	2,091	1,410	792	1,823	1,519
1947	2,663	5,605	4,722	1,967	1,620	1,736	905	2,492	1,644	993	2,195	2,358
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	2,177	5,321	3,664	2,190	1,464	2,391	882	3,320	1,901	792	1,802	1,817
1941	2,010	5,327	3,864	2,327	1,640	2,274	773	2,543	1,876	725	1,714	1,679
1940	2,289	5,690	3,874	2,947	1,622	2,652	837	3,271	1,979	750	1,760	1,598
1939	2,896	6,655	4,225	2,686	1,592	2,670	1,119	2,917	1,937	802	1,896	1,827
1938	2,690	6,148	4,605	2,996	1,767	2,963	1,111	2,883	2,008	878	2,184	2,015
1937	2,350	6,763	4,464	2,759	1,826	3,322	977	3,183	2,179	850	2,133	1,838
1936	2,789	6,664	4,605	2,783	1,873	2,601	1,060	3,019	2,049	861	2,086	2,300
1935	2,351	6,352	4,339	2,632	1,712	2,456	949	3,012	2,198	1,017	2,063	2,078
1934	2,784	6,194	4,796	2,853	2,407	2,937	1,227	3,208	2,441	985	2,241	2,312
1933	2,991	7,062	5,147	2,739	2,028	2,604	1,408	3,054	2,610	1,181	2,228	2,334
1932	2,933	7,666	5,248	2,929	2,190	3,691	1,292	3,184	2,639	1,228	2,469	2,631
1931	2,864	6,917	5,028	3,216	2,104	2,541	1,283	3,675	2,629	1,147	2,719	2,894
1930	2,842	8,092	5,422	3,255	2,044	3,465	1,213	3,539	2,757	1,355	2,828	2,701
1929	3,243	7,713	5,676	4,049	2,304	3,929	1,428	3,597	3,277	1,406	2,907	2,630
1928	3,069	7,289	5,462	3,753	2,062	3,461	1,221	3,294	2,931	1,421	2,630	2,396
1927	3,601	7,625	5,639	3,799	2,326	2,903	1,328	3,055	3,345	1,623	2,533	2,819
1926	3,375	5,514	5,323	3,482	2,017	2,758	1,211	3,576	2,926	1,546	2,255	2,680
1925	3,316	7,030	5,577	3,599	2,084	2,574	1,205	3,170	3,291	1,293	2,558	2,893
1924	3,784	6,525	5,599	3,958	2,043	3,247	1,664	3,179	3,093	1,480	2,583	2,908
1923	3,692	6,669	6,401	4,061	2,407	3,018	1,536	3,640	4,388	1,796	2,824	3,290
1922	3,893	6,375	5,929	3,929	2,185	3,248	1,470	3,376	3,295	1,493	2,897	3,094
1921	3,496	6,040	5,965	3,703	2,098	3,007	1,462	2,926	3,742	1,565	2,953	3,490
1920	3,345	6,237	5,871	3,668	1,863	2,719	1,395	2,318	3,213	1,452	2,511	2,876

TABLE 17. - 1/ DEATHS FROM TEN SELECTED CAUSES BY PREFECTURE: JAPAN, 1920-1949 Cont'd
6/ DIARRHEA, ENTERITIS AND ULCERATION OF THE INTESTINES (ALL AGES) Cont'd

Year	Mie	Miyagi	Miyazaki	Nagano	Nagasaki	Nara	Niigata	Oita	Okayama	Osaka	Saga	Saitama
1949	1,450	1,247	1,010	1,403	1,418	812	2,803	1,324	1,474	2,934	934	1,978
1948	1,612	1,647	1,098	1,882	1,564	1,012	3,426	1,416	1,798	2,951	1,070	2,245
1947	2,304	1,513	1,238	2,354	1,982	1,220	3,735	1,926	2,577	4,018	1,400	2,794
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	1,932	1,750	1,240	2,272	1,766	903	4,110	1,476	1,780	4,283	890	2,667
1941	2,037	1,561	1,066	2,435	1,515	1,001	3,884	1,367	1,723	4,349	936	3,278
1940	1,969	2,106	1,261	2,735	1,597	952	4,970	1,507	1,960	4,095	929	4,109
1939	2,058	2,282	1,519	2,847	1,765	1,075	5,169	1,497	2,279	4,397	989	3,762
1938	2,434	2,293	1,342	2,757	1,826	1,156	5,186	1,496	2,409	4,095	950	4,091
1937	2,149	2,609	1,446	2,942	2,170	1,131	5,373	1,825	2,252	4,436	1,105	3,730
1936	2,326	2,331	1,247	2,857	2,173	1,074	5,618	1,575	2,219	4,063	1,402	3,558
1935	2,079	2,244	1,344	2,540	1,853	1,102	4,336	1,555	2,068	4,686	1,143	3,213
1934	2,731	2,473	1,279	3,098	1,776	1,353	6,081	1,604	2,128	5,235	1,212	3,776
1933	2,635	2,568	1,452	3,109	1,882	1,371	5,611	1,678	2,754	5,299	1,352	3,842
1932	2,539	3,176	1,536	3,181	2,154	1,389	5,547	1,855	2,344	4,786	1,640	4,168
1931	2,898	3,074	1,648	3,107	2,345	1,393	6,198	1,956	2,405	5,508	1,756	4,566
1930	2,724	3,222	1,415	3,838	2,149	1,246	5,670	1,798	2,378	5,519	1,742	4,110
1929	3,010	3,752	1,520	3,820	2,340	1,486	6,979	1,955	2,858	5,644	1,746	4,917
1928	3,117	3,216	1,229	3,479	2,050	1,282	6,337	1,788	2,603	5,262	1,405	4,795
1927	2,847	3,212	1,561	3,684	2,094	1,333	5,997	1,922	2,862	6,067	1,610	4,612
1926	2,728	2,900	1,237	3,677	1,567	1,253	6,385	1,700	2,531	6,269	1,554	4,073
1925	2,944	2,660	1,149	3,573	1,609	1,420	6,181	1,609	2,233	6,181	1,389	4,492
1924	2,959	3,049	1,308	3,965	1,853	1,374	6,741	1,693	2,761	7,058	1,587	4,691
1923	3,286	3,097	1,523	3,919	1,918	1,456	6,975	1,688	2,967	7,114	1,596	5,167
1922	2,785	2,774	1,400	4,041	1,871	1,467	7,147	1,464	2,696	8,052	1,490	4,319
1921	3,529	2,691	1,244	3,757	1,898	1,611	6,929	1,825	2,859	6,905	1,626	4,807
1920	3,174	2,414	1,108	3,965	1,674	1,573	5,675	1,361	2,705	6,164	1,432	4,339

TABLE 17. - 1/ DEATHS FROM TEN SELECTED CAUSES BY PREFECTURE: JAPAN, 1920-1949 Cont'd
6/ DIARRHEA, ENTERITIS AND ULCERATION OF THE INTESTINES (ALL AGES) (119-120) CONT'D

Year	Shiga	Shimane	Shizuoka	Tochigi	Tokushima	Tokyo	Tottori	Toyama	Wakayama	Yamagata	Yamaguchi	Yamanashi
1949	1,204	939	1,890	1,626	853	2,244	608	1,757	988	1,728	1,224	736
1948	1,268	994	2,057	1,673	945	2,648	673	1,855	951	2,013	1,279	856
1947	1,935	1,147	2,494	2,005	1,267	3,196	866	2,413	1,284	2,109	1,974	1,074
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	1,338	1,071	2,291	2,118	880	3,776	710	2,229	1,231	2,396	1,498	990
1941	1,427	1,005	2,488	2,329	853	3,235	621	2,016	1,097	2,379	1,261	1,238
1940	1,472	1,095	2,419	2,723	909	3,585	654	2,501	1,220	2,854	1,325	1,281
1939	1,177	1,409	2,505	2,858	984	3,565	819	2,428	1,279	3,054	1,461	1,263
1938	1,642	1,161	2,780	3,103	1,023	3,792	775	2,634	1,372	2,973	1,562	1,331
1937	1,609	1,129	2,982	2,819	1,227	4,016	846	2,804	1,238	3,179	1,421	1,411
1936	1,573	1,208	3,453	2,945	889	5,176	873	2,377	1,284	3,450	1,517	1,390
1935	1,488	1,147	2,846	2,860	1,095	4,231	892	2,349	1,714	2,964	1,538	1,385
1934	1,813	1,327	3,073	3,379	1,167	4,762	946	2,910	1,493	3,707	1,548	1,317
1933	1,754	1,402	3,465	3,003	1,254	6,117	951	2,454	1,732	3,781	1,609	1,676
1932	1,759	1,674	3,398	2,920	1,309	5,109	1,173	2,606	1,629	3,731	1,808	1,457
1931	1,956	1,467	3,406	3,445	1,378	5,810	1,053	2,848	1,598	3,600	1,891	1,485
1930	1,873	1,637	3,968	3,152	1,195	5,778	1,251	2,438	1,715	3,843	1,951	1,449
1929	1,908	1,900	4,088	3,673	1,470	7,409	1,223	2,822	1,672	4,229	2,118	1,688
1928	1,890	1,549	3,873	2,416	1,338	6,858	1,088	2,510	1,718	3,947	1,899	1,438
1927	2,067	1,514	3,825	3,712	1,520	6,714	1,110	2,515	1,703	3,979	2,137	1,627
1926	1,996	1,518	3,938	3,141	1,437	6,937	1,129	2,582	1,745	4,167	1,937	1,526
1925	2,078	1,485	4,084	3,247	1,206	6,883	1,005	2,497	1,742	3,456	1,821	1,983
1924	1,970	1,559	3,822	3,317	1,612	7,527	1,110	2,509	1,857	3,713	2,135	1,896
1923	2,538	1,702	4,926	4,077	1,719	9,055	1,276	2,821	2,053	4,126	2,267	1,869
1922	2,075	1,347	3,671	3,479	1,745	8,989	1,127	2,555	2,179	3,935	1,920	1,981
1921	2,643	1,346	4,238	3,441	1,692	8,479	1,137	2,298	2,043	3,789	1,934	2,079
1920	2,432	1,364	3,842	3,262	1,632	7,671	1,197	1,950	1,850	3,394	1,921	1,944

See footnotes at end of table.

TABLE 17. - 1/ DEATHS FROM TEN SELECTED CAUSES BY PREFECTURE: JAPAN, 1920-49
2/ NEPHRITIS, ALL FORMS (130-132)

Year	All Japan	Aichi	Akita	Aomori	Chiba	Ehime	Fukui	Fukuoka	Fukushima	Gifu	Gumma
1949	33,859	1,010	674	383	1,076	562	276	1,478	737	521	685
1948	36,561	1,187	753	470	1,182	568	303	1,539	840	560	718
1947	45,017	1,462	803	474	1,329	717	318	2,223	984	718	852
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	49,708	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	50,598	1,688	980	656	1,211	637	403	2,383	1,048	756	904
1941	59,445	1,570	984	630	1,247	619	418	2,165	977	798	841
1940	54,142	1,757	1,240	739	1,281	752	495	2,327	1,020	784	892
1939	58,314	1,703	1,079	754	1,334	799	558	2,726	1,067	864	934
1938	61,535	1,897	962	726	1,606	829	662	3,207	1,182	892	978
1937	55,183	1,711	951	644	1,392	791	586	2,955	1,065	756	935
1936	57,239	1,757	995	952	1,411	788	560	2,831	1,056	875	925
1935	55,510	1,737	969	620	1,355	864	578	2,645	1,017	852	970
1934	59,059	1,816	980	708	1,435	866	687	2,514	1,190	884	1,006
1933	59,538	1,831	1,007	611	1,592	856	580	2,647	1,112	911	1,001
1932	60,899	1,836	1,093	715	1,613	900	607	2,807	1,198	910	1,067
1931	63,790	1,850	1,099	657	1,688	900	709	3,023	1,173	1,005	1,070
1930	62,910	1,766	1,159	802	1,676	889	718	2,965	1,237	926	1,016
1929	64,930	1,899	1,626	1,024	1,775	874	595	2,762	1,221	944	1,073
1928	62,421	1,904	1,150	753	1,910	883	584	2,737	1,242	955	1,007
1927	59,554	1,684	1,298	716	1,523	874	572	2,486	1,174	835	1,003
1926	59,110	1,714	1,458	792	1,566	894	544	2,545	1,106	867	990
1925	59,353	1,657	1,319	705	1,690	885	558	2,556	1,125	847	998
1924	61,657	1,644	1,223	673	1,710	966	543	2,544	1,242	871	1,119
1923	62,046	1,707	1,872	891	1,788	890	577	2,460	1,211	904	1,061
1922	60,449	1,596	1,446	940	1,609	902	565	2,378	1,296	852	1,019
1921	59,320	1,830	1,103	691	1,618	896	595	2,480	1,266	901	979
1920	54,941	1,526	944	594	1,434	817	558	2,585	1,202	779	903

TABLE 17. - DEATHS FROM TEN SELECTED CAUSES BY PREFECTURE: JAPAN, 1920-1949, Cont'd
7 NEPHRITIS, ALL FORMS (130-132) - Cont'd

Year	Hiroshima	Hokkaido	Hyogo	Ibaraki	Ishikawa	Iwate	Kagawa	Kagoshima	Kanagawa	Kochi	Kumamoto	Kyoto
1949	811	1,115	1,158	1,075	369	505	419	1,046	772	426	871	563
1948	922	1,147	1,335	1,146	432	572	439	1,016	875	412	1,071	648
1947	1,176	1,215	1,728	1,313	525	671	562	1,367	1,099	530	1,315	888
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	1,225	1,527	2,045	1,339	509	684	568	1,666	1,441	500	1,214	1,141
1941	1,118	1,659	1,940	1,306	521	735	492	1,588	1,386	492	1,136	1,113
1940	1,270	1,796	2,134	1,423	549	874	550	1,698	1,415	502	1,252	1,229
1939	1,417	2,046	2,397	1,453	630	821	638	1,682	1,609	571	1,451	1,289
1938	1,394	1,823	2,451	1,501	679	826	679	1,751	1,772	597	1,514	1,426
1937	1,226	1,816	2,248	1,385	662	777	569	1,527	1,552	545	1,325	1,230
1936	1,964	1,947	2,317	1,410	638	782	614	1,597	1,567	594	1,463	1,401
1935	1,341	1,741	2,178	1,326	616	800	595	1,587	1,530	639	1,397	1,265
1934	1,393	1,853	2,239	1,510	759	861	614	1,600	1,605	596	1,420	1,363
1933	1,387	1,933	2,343	1,528	660	850	617	1,554	1,764	619	1,451	1,398
1932	1,464	1,877	2,195	1,565	702	857	559	1,614	1,791	640	1,462	1,434
1931	1,567	2,049	2,325	1,585	692	887	650	1,603	1,637	644	1,559	1,515
1930	1,595	1,860	2,189	1,592	686	977	638	1,647	1,672	684	1,452	1,464
1929	1,591	2,318	2,118	1,619	711	1,009	633	1,598	1,798	710	1,460	1,457
1928	1,499	1,697	2,265	1,573	675	883	557	1,476	1,714	756	1,451	1,548
1927	1,487	1,949	2,137	1,481	688	874	645	1,472	1,563	643	1,391	1,520
1926	1,519	1,762	2,023	1,471	636	882	597	1,480	1,558	603	1,449	1,367
1925	1,604	1,709	2,249	1,650	609	993	624	1,434	1,675	541	1,291	1,509
1924	1,625	1,870	2,438	1,628	635	1,059	634	1,468	1,638	628	1,280	1,602
1923	1,584	1,811	2,224	1,481	755	1,133	618	1,466	1,556	627	1,320	1,559
1922	1,493	1,657	2,220	1,550	718	1,144	599	1,551	1,681	682	1,410	1,494
1921	1,470	1,706	2,175	1,403	663	961	573	1,520	1,637	715	1,350	1,544
1920	1,321	1,616	1,998	1,365	558	922	597	1,490	1,446	649	1,252	1,571

TABLE 17. - 1/ DEATHS FROM TEN SELECTED CAUSES BY PREFECTURE: JAPAN, 1920-49 Cont'd
2/ MALE, ALL FORMS (130-132) - (Cont'd)

Year	Mie	Miyagi	Miyazaki	Nagano	Nagasaki	Nara	Niigata	Oita	Okayama	Osaka	Saga	Saitama
1949	617	619	622	947	697	394	1,155	661	659	1,582	579	1,016
1948	628	624	583	1,090	728	400	1,324	734	705	1,847	630	1,066
1947	795	764	722	1,172	924	527	1,528	828	857	2,241	736	1,394
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	780	977	729	1,301	1,000	464	1,584	840	693	3,253	616	1,427
1941	810	904	680	1,289	1,035	498	1,586	772	727	2,962	676	1,339
1940	898	902	749	1,283	1,072	491	1,793	905	809	3,461	736	1,546
1939	917	999	828	1,473	1,170	589	1,870	912	890	3,759	732	1,502
1938	1,047	1,060	829	1,411	1,234	648	1,966	962	958	3,893	836	1,592
1937	891	1,092	732	1,326	1,162	573	1,752	901	803	3,690	856	1,389
1936	1,000	1,007	779	1,358	1,132	574	1,813	911	844	3,751	799	1,474
1935	1,001	1,043	774	1,446	1,162	595	1,741	884	859	3,512	730	1,410
1934	1,073	1,052	898	1,625	1,106	624	2,247	903	853	3,671	704	1,712
1933	1,047	1,101	772	1,576	1,133	600	1,981	939	938	3,761	860	1,741
1932	1,064	1,101	891	1,615	1,148	641	1,948	938	844	3,847	887	1,719
1931	1,269	1,227	895	1,689	1,174	693	2,184	888	926	3,889	851	1,976
1930	1,081	1,417	882	1,872	1,137	602	2,187	948	898	3,807	884	1,797
1929	1,155	1,382	832	1,843	1,103	624	2,421	944	982	3,705	782	1,831
1928	1,246	1,293	833	1,681	1,105	617	2,072	889	979	3,735	852	1,838
1927	1,027	1,222	800	1,597	1,071	628	2,124	885	920	3,437	821	1,679
1926	954	1,267	731	1,661	1,008	570	2,307	847	877	3,337	780	1,630
1925	1,142	1,356	661	1,718	955	602	2,092	768	941	3,447	755	1,749
1924	1,056	1,392	752	1,793	963	577	2,413	898	998	3,646	788	1,821
1923	1,109	1,519	759	1,787	1,026	617	2,271	772	927	3,293	802	1,626
1922	959	1,439	704	1,906	975	557	2,258	806	953	3,361	894	1,585
1921	1,155	1,284	730	1,706	1,022	697	2,036	821	913	3,291	890	1,664
1920	1,088	1,245	653	1,671	928	633	2,004	775	830	3,083	814	1,399

TABLE 17. - 1/ DEATHS FROM TEN SELECTED CAUSES BY PREFECTURE: JAPAN, 1920-1949, Cont'd
2/ NEPHRITIS, ALL FORMS (130-132) - Cont'd

Year	Shiga	Shimane	Shizuoka	Tochigi	Tokushima	Tokyo	Tottori	Toyama	Wakayama	Yamagata	Yamaguchi	Yamaguchi
1949	425	434	878	676	397	2,068	374	510	267	667	679	404
1948	469	450	956	710	428	2,085	354	507	312	779	778	459
1947	579	509	1,245	837	688	2,324	465	647	435	956	1,082	557
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	566	572	1,237	939	536	4,415	522	677	415	931	1,079	520
1942	587	643	1,233	791	522	4,379	510	682	457	1,025	1,093	470
1941	640	737	1,345	882	592	4,798	552	749	502	1,163	1,090	468
1940												
1939	646	803	1,515	938	688	5,306	644	835	554	1,132	1,225	565
1938	751	855	1,640	969	668	5,706	637	888	571	1,121	1,345	594
1937	695	729	1,467	881	637	4,996	548	807	508	1,034	1,158	568
1936	681	772	1,419	855	645	4,845	533	798	536	1,028	1,298	543
1935	664	734	1,409	805	701	4,774	548	778	571	1,107	1,170	530
1934	797	794	1,443	917	731	4,983	537	981	566	1,255	1,184	564
1933	780	862	1,476	1,012	679	5,240	547	821	513	1,171	1,198	600
1932	744	833	1,618	978	732	5,363	562	894	585	1,241	1,213	587
1931	742	953	1,538	1,059	830	5,745	599	934	668	1,216	1,368	650
1930	741	995	1,599	992	775	5,470	575	876	580	1,251	1,304	640
1929	739	878	1,572	1,028	839	6,021	567	923	557	1,470	1,293	664
1928	729	874	1,684	931	834	5,759	556	842	612	1,335	1,249	627
1927	682	844	1,593	964	784	5,433	575	905	542	1,283	1,348	617
1926	704	813	1,438	947	721	5,712	519	899	581	1,191	1,199	654
1925	775	833	1,488	940	783	5,326	512	795	604	1,165	1,212	565
1924	775	845	1,585	1,057	819	5,329	549	864	573	1,169	1,345	670
1923	817	956	1,620	1,004	765	5,528	576	923	604	1,188	1,410	652
1922	707	773	1,488	997	725	5,843	493	742	568	1,077	1,212	645
1921	813	809	1,580	953	723	5,565	492	792	577	955	1,134	712
1920	743	724	1,303	891	709	5,167	365	747	553	1,036	1,020	623

See footnotes at end of table.

TABLE 17. - 1/ DEATHS FROM TEN SELECTED CAUSES BY PREFECTURE, JAPAN, 1920-1949 Cont'd
CONGENITAL DEBILITY (158)

Year	All Japan	Aichi	Akita	Aomori	Chiba	Ehime	Fukui	Fukuoka	Fukushima	Gifu	Gumma
1949	37,825	1,654	1,340	1,114	901	674	495	1,537	943	922	607
1948	38,204	1,716	1,077	1,125	992	719	553	1,596	1,055	947	691
1947	41,751	1,617	1,218	1,063	1,097	721	535	1,669	1,133	984	716
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	51,941	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	52,074	2,242	1,410	1,222	1,158	853	685	2,025	1,339	1,213	829
1941	52,348	2,266	1,330	1,178	1,202	834	728	2,027	1,322	1,196	879
1940	54,925	2,397	1,588	1,380	1,292	858	802	2,034	1,400	1,349	917
1939	57,102	2,419	1,576	1,339	1,362	913	673	2,244	1,572	1,386	874
1938	60,556	2,504	1,685	1,525	1,367	869	853	2,501	1,659	1,447	939
1937	63,367	2,550	1,739	1,454	1,549	1,039	894	2,595	1,700	1,510	956
1936	67,967	2,861	1,842	1,770	1,615	1,111	992	2,734	1,784	1,701	967
1935	64,355	2,611	1,838	1,590	1,564	1,038	853	2,451	1,780	1,560	964
1934	68,599	2,861	1,956	1,630	1,525	1,175	1,119	2,571	2,024	1,821	1,060
1933	67,480	2,617	1,823	1,642	1,603	1,149	972	2,701	1,836	1,679	946
1932	59,379	2,398	1,580	1,417	1,501	1,000	841	2,153	1,712	1,502	861
1931	61,440	2,527	1,704	1,485	1,601	1,036	931	2,271	1,769	1,460	919
1930	59,276	2,309	1,667	1,524	1,581	1,030	825	2,179	1,661	1,468	947
1929	65,400	2,541	1,821	1,439	1,587	1,116	935	2,345	1,862	1,608	1,111
1928	63,899	2,610	1,755	1,431	1,698	986	886	2,319	1,759	1,533	1,126
1927	64,123	2,568	1,806	1,336	1,637	1,011	964	2,244	1,894	1,553	1,121
1926	67,219	2,784	1,878	1,524	1,835	1,056	985	2,317	1,897	1,653	1,133
1925	67,014	2,740	1,931	1,549	1,783	1,019	948	2,315	1,934	1,661	1,212
1924	69,539	2,808	2,006	1,540	1,914	1,097	911	2,407	2,062	1,632	1,246
1923	73,740	2,897	2,256	1,734	2,131	1,054	1,060	2,524	2,279	1,849	1,350
1922	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1921	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1920	NA	NA	NA	NA	na	NA	NA	NA	NA	NA	NA

TABLE 17. - $\frac{1}{2}$ DEATHS FROM TEN SELECTED CAUSES BY PREFECTURE: JAPAN, 1920-1949, Cont'd
CONGENITAL DEBILITY (158) - Cont'd

Year	Hiroshima	Hokkaido	Hyogo	Ibaraki	Ishikawa	Iwate	Kagawa	Kagoshima	Kanagawa	Kochi	Kumamoto	Kyoto
1949	806	1,616	1,161	1,120	488	937	566	707	546	451	807	660
1948	870	1,666	1,350	1,157	568	909	609	688	619	450	752	682
1947	1,031	1,644	1,429	1,327	602	963	784	723	724	NA	822	759
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	1,275	1,953	2,125	1,449	770	1,405	744	767	1,073	544	869	1,164
1941	1,292	1,953	2,041	1,553	708	1,372	701	697	1,056	510	904	1,136
1940	1,263	2,119	1,993	1,755	816	1,541	775	734	1,143	507	896	1,190
1939	1,341	2,273	2,280	1,724	737	1,428	832	778	1,168	551	925	1,234
1938	1,532	2,539	2,377	1,882	840	1,515	861	822	1,195	650	986	1,427
1937	1,587	2,298	2,378	2,045	927	1,592	906	787	1,192	769	982	1,469
1936	1,695	2,421	2,777	2,093	1,031	1,653	957	856	1,184	724	1,093	1,568
1935	1,617	2,395	2,501	1,998	869	1,507	839	798	1,240	777	1,028	1,427
1934	1,717	2,486	2,715	2,124	995	1,652	956	855	1,187	791	1,143	1,549
1933	1,733	2,600	2,760	2,046	955	1,710	994	787	1,248	880	1,105	1,506
1932	1,473	2,116	2,460	1,901	803	1,509	879	706	1,085	811	980	1,284
1931	1,497	2,377	2,448	1,996	915	1,539	898	706	1,141	806	1,090	1,304
1930	1,485	2,235	2,371	1,933	754	1,485	847	768	1,165	797	924	1,279
1929	1,606	2,465	2,797	2,120	921	1,620	956	653	1,291	1,008	1,012	1,388
1928	1,606	2,337	2,656	2,138	910	1,697	916	614	1,261	973	963	1,299
1927	1,552	2,337	2,649	2,071	1,002	1,619	1,004	722	1,302	894	968	1,346
1926	1,738	2,452	2,898	2,262	974	1,655	1,064	671	1,246	1,037	953	1,495
1925	1,727	2,568	2,650	2,316	955	1,638	981	624	1,522	1,022	865	1,501
1924	1,790	2,714	2,937	2,515	1,036	1,853	1,021	611	1,401	965	980	1,492
1923	1,888	2,671	3,087	2,558	1,071	1,861	1,048	663	1,404	1,128	994	1,520
1922	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1921	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1920	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA

TABLE 17. - 1/ DEATHS FROM TEN SELECTED CAUSES BY PREFECTURE, JAPAN, 1920-1949 Cont'd
CONGENITAL DEBILITY (158) - Cont'd

Year	Mie	Miyagi	Miyazaki	Nagano	Nagasaki	Nara	Niigata	Oita	Okayama	Osaka	Saga	Saitama
1949	698	798	585	642	889	408	1,045	707	890	1,385	669	1,137
1948	754	883	480	732	760	486	1,077	689	1,000	1,416	672	1,107
1947	876	842	516	819	844	528	1,128	882	1,150	1,584	801	1,190
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	998	1,078	569	910	1,108	582	1,202	954	1,214	2,767	749	1,295
1941	1,054	1,009	552	1,011	996	607	1,261	939	1,216	2,841	762	1,502
1940	1,091	1,241	578	1,059	1,028	630	1,333	895	1,288	2,581	822	1,586
1939	1,116	1,151	632	1,065	1,116	686	1,354	998	1,389	3,012	865	1,609
1938	1,325	1,291	639	1,157	1,161	750	1,317	1,083	1,482	2,894	920	1,627
1937	1,344	1,378	679	1,133	1,203	783	1,440	1,246	1,646	3,107	987	1,724
1936	1,354	1,443	769	1,278	1,334	785	1,693	1,329	1,822	3,090	1,126	1,812
1935	1,329	1,566	658	1,308	1,162	757	1,523	1,216	1,613	3,125	1,042	1,656
1934	1,468	1,553	654	1,365	1,263	820	1,614	1,249	1,772	3,192	1,109	1,840
1933	1,538	1,483	714	1,290	1,207	821	1,547	1,246	1,874	3,045	1,132	1,888
1932	1,209	1,294	565	1,162	1,021	740	1,235	1,082	1,639	2,845	970	1,654
1931	1,165	1,291	535	1,196	959	730	1,438	1,037	1,687	2,896	927	1,800
1930	1,175	1,260	511	1,220	975	735	1,320	973	1,564	2,862	946	1,626
1929	1,263	1,442	583	1,321	1,039	841	1,476	1,057	1,857	3,063	1,037	1,941
1928	1,289	1,283	479	1,255	1,002	786	1,426	1,000	1,799	2,943	1,025	1,929
1927	1,260	1,391	503	1,335	1,011	899	1,463	986	1,785	3,056	1,007	1,928
1926	1,412	1,453	545	1,457	1,001	844	1,516	1,080	1,976	3,023	1,053	2,021
1925	1,412	1,491	505	1,489	967	826	1,494	1,033	1,798	3,091	949	2,083
1924	1,366	1,577	541	1,435	936	927	1,670	1,127	2,039	3,253	1,007	2,130
1923	1,538	1,707	501	1,478	1,010	880	1,643	1,121	2,119	3,378	1,098	2,502
1922	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1921	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1920	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA

TABLE 17. - 1/ DEATHS FROM TEN SELECTED CAUSES BY PREFECTURE: JAPAN, 1920-1949, Cont'd
CONGENITAL DEBILITY (158) - Cont'd

Year	Shiga	Shimane	Shizuoka	Tochigi	Tohama	Tokyo	Tottori	Toyama	Wakayama	Yamagata	Yamanashi
1949	489	460	1,026	605	542	1,376	293	596	347	905	265
1948	466	486	998	589	595	1,425	306	549	363	872	276
1947	555	606	1,067	714	620	1,538	348	683	465	937	288
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	685	690	1,394	764	682	3,502	337	755	502	1,165	356
1941	631	737	1,444	781	730	3,537	415	697	501	1,176	377
1940	698	716	1,544	919	746	3,419	405	786	554	1,217	392
1939	663	854	1,673	905	801	3,447	373	693	612	1,290	399
1938	706	889	1,820	900	863	3,480	447	768	668	1,343	417
1937	788	920	1,831	956	894	3,627	449	778	706	1,486	437
1936	856	956	2,068	1,059	929	3,810	511	814	775	1,471	417
1935	772	929	2,010	950	878	3,629	443	778	728	1,559	502
1934	906	980	2,122	1,006	961	3,672	525	860	786	1,629	454
1933	819	1,002	1,946	988	935	3,799	480	863	827	1,534	454
1932	640	873	1,818	838	865	3,529	469	629	717	1,454	415
1931	697	859	1,709	861	905	3,785	475	710	661	1,532	399
1930	686	812	1,645	857	832	3,546	464	646	680	1,482	446
1929	675	875	1,856	987	890	4,162	498	670	714	1,631	440
1928	690	822	1,882	872	879	4,157	472	716	739	1,764	450
1927	702	875	1,734	958	777	3,854	489	720	701	1,760	417
1926	805	855	1,930	967	889	3,798	487	752	659	1,861	472
1925	797	899	1,854	979	844	3,920	563	717	727	1,864	508
1924	795	852	1,991	987	899	3,806	561	735	700	1,959	479
1923	897	976	2,012	1,044	985	3,996	566	712	813	2,072	518
1922	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1921	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1920	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA

See footnotes at end of table.

TABLE 17. - 1/ DEATHS FROM TEN SELECTED CAUSES BY PREFECTURE: JAPAN, 1920-1949 Cont'd
2/ SEXILITY (162)

Year	All Japan	Aichi	Akita	Amori	Chiba	Ehime	Fukui	Fukuoka	Fukushima	Gifu	Gumma
1920	66,699	2,685	1,096	965	2,188	1,776	895	2,348	1,599	1,399	1,140
1921	64,674	2,421	1,053	1,165	2,302	1,658	856	2,096	1,565	1,306	1,110
1922	76,052	3,121	1,201	1,004	2,590	2,027	1,004	2,909	2,039	1,672	1,270
1923	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1924	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1925	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1926	98,694	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1927	95,436	3,586	1,374	1,133	3,038	2,127	1,183	3,243	2,278	1,923	1,604
1928	89,333	3,562	1,297	1,057	2,989	1,900	1,167	2,910	2,140	1,987	1,509
1929	89,023	3,724	1,469	1,253	2,714	1,999	1,202	2,766	2,205	1,900	1,604
1930	94,496	3,653	1,439	1,334	2,854	2,193	1,216	3,058	2,190	1,859	1,616
1931	97,921	4,140	1,282	1,214	3,356	2,264	1,266	3,265	2,380	2,319	1,688
1932	84,624	3,448	1,098	1,009	2,705	2,040	1,164	2,824	1,965	1,891	1,385
1933	90,990	3,736	1,157	1,282	2,840	2,106	1,286	2,901	2,166	2,137	1,571
1934	78,421	3,386	1,108	892	2,582	1,969	1,117	2,592	1,791	1,779	1,338
1935	86,529	3,742	1,103	986	2,780	2,114	1,470	2,577	2,058	2,051	1,482
1936	82,452	3,320	1,078	953	2,691	2,101	1,227	2,553	1,893	1,857	1,325
1937	76,736	3,291	1,038	889	2,374	1,959	1,156	2,433	1,748	1,715	1,349
1938	84,774	3,769	1,113	986	2,734	2,103	1,295	2,784	2,042	1,926	1,483
1939	75,755	3,059	1,025	833	2,407	1,926	1,173	2,665	1,786	1,643	1,318
1940	78,373	3,274	1,246	1,048	2,496	2,014	1,247	2,668	1,856	1,694	1,373
1941	76,522	3,552	1,083	898	2,685	1,905	1,104	2,584	1,746	1,801	1,259
1942	73,835	3,168	1,165	867	2,472	1,892	1,142	2,413	1,829	1,627	1,308
1943	66,799	2,899	1,193	887	2,208	1,693	926	2,244	1,577	1,371	1,130
1944	69,223	3,152	1,163	743	2,327	1,728	1,017	2,262	1,621	1,628	1,211
1945	73,155	2,982	1,181	752	2,392	1,891	968	2,476	1,747	1,598	1,284
1946	75,763	3,533	1,598	894	2,887	1,861	1,035	2,384	1,634	1,692	1,288
1947	75,206	3,017	1,482	1,069	2,628	1,886	1,028	2,388	1,892	1,560	1,291
1948	74,927	3,940	1,281	841	2,763	1,958	1,016	2,474	1,648	1,607	1,291
1949	72,645	3,442	1,178	705	2,649	1,892	1,051	2,594	1,618	1,597	1,185

TABLE 17. - 1/ DEATHS FROM TEN SELECTED CAUSES BY PREFECTURE: JAPAN, 1920-1949 Cont'd
3/ SENILITY (162) - Cont'd

Year	Hiroshima	Hokkaido	Hyogo	Ibaraki	Ishikawa	Iwate	Kagawa	Kagoshima	Kanagawa	Kochi	Kumamoto	Kyoto
1949	2,024	2,269	2,533	1,871	1,076	1,104	1,056	1,664	1,398	1,179	1,693	1,418
1948	1,817	2,167	2,605	1,901	921	1,057	886	1,397	1,325	1,176	1,484	1,426
1947	2,378	2,501	3,509	2,224	999	1,186	1,260	1,802	1,490	1,441	1,869	1,859
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	2,806	2,346	4,374	2,580	1,295	1,640	356	2,687	2,031	1,606	2,299	2,239
1941	2,526	2,337	3,778	2,442	1,276	1,503	1,220	2,227	1,948	1,590	1,941	2,017
1940	2,712	2,293	3,804	2,439	1,377	1,551	1,291	2,262	1,804	1,494	1,932	2,081
1939	3,051	2,985	4,180	2,504	1,336	1,611	1,542	2,311	1,898	1,571	2,044	2,216
1938	3,102	2,574	4,433	2,642	1,464	1,511	1,463	2,097	1,869	1,628	2,091	2,313
1937	2,565	2,407	3,793	2,099	1,379	1,427	1,303	1,899	1,639	1,474	1,925	1,895
1936	2,784	2,660	4,104	2,357	1,516	1,545	1,264	2,134	1,651	1,417	2,160	2,026
1935	2,452	2,209	3,617	2,041	1,245	1,271	1,188	1,882	1,371	1,319	1,941	1,726
1934	2,446	2,138	4,017	2,379	1,696	1,388	1,317	1,949	1,599	1,402	1,897	1,879
1933	2,521	2,313	4,001	2,168	1,309	1,514	1,261	1,810	1,560	1,453	1,835	1,853
1932	2,269	1,851	3,651	2,004	1,363	1,434	1,084	1,788	1,351	1,482	1,728	1,793
1931	2,479	2,125	3,973	2,101	1,497	1,466	1,270	1,891	1,533	1,451	1,959	2,027
1930	2,273	1,808	3,617	1,794	1,294	1,352	1,110	1,814	1,319	1,256	1,849	1,843
1929	2,326	2,074	3,535	1,899	1,525	1,350	1,147	1,704	1,346	1,331	1,820	1,863
1928	2,115	1,766	3,334	1,879	1,357	1,356	957	1,639	1,432	1,466	1,748	1,779
1927	2,025	1,815	3,361	1,999	1,522	1,360	1,122	1,598	1,231	1,273	1,579	1,693
1926	1,916	1,653	3,021	1,601	1,343	1,188	955	1,483	1,231	1,258	1,493	1,553
1925	2,070	1,670	3,044	1,817	1,292	1,300	947	1,499	1,159	1,374	1,411	1,579
1924	2,282	1,615	3,458	1,871	1,435	1,272	1,202	1,535	1,278	1,471	1,575	1,597
1923	2,216	1,476	3,490	1,861	1,591	1,216	1,108	1,464	1,629	1,368	1,455	1,609
1922	2,280	1,473	3,400	1,896	1,610	1,413	1,284	1,560	1,295	1,523	1,557	1,583
1921	2,159	1,342	3,324	1,729	1,299	1,207	1,136	1,562	1,404	1,426	1,669	1,781
1920	2,074	1,340	3,436	1,869	1,367	1,109	1,185	1,395	1,340	1,466	1,477	1,710

TABLE 17. - 1/ DEATHS FROM TEN SELECTED CAUSES BY PREFECTURE: JAPAN, 1920-1949 Cont'd
8/ SENILITY (162) - Cont'd

Year	Mie	Miyagi	Miyazaki	Nagano	Nagasaki	Nara	Niigata	Oita	Okayama	Oosaka	Saga	Saitama
1949	1,262	1,070	928	1,731	1,709	713	1,729	1,493	1,971	2,033	853	1,770
1948	1,340	1,113	750	1,711	1,509	692	1,711	1,354	1,953	2,213	847	1,738
1947	1,686	1,165	1,045	2,095	1,984	843	2,025	1,612	2,324	2,827	1,141	1,827
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	2,008	1,485	1,267	2,588	2,356	945	2,572	1,897	2,759	4,074	1,134	1,993
1941	1,941	1,392	1,142	2,533	2,088	906	2,499	1,684	2,621	3,463	1,006	2,219
1940	1,945	1,365	1,060	2,435	2,018	860	2,660	1,702	2,552	3,399	962	2,145
1939	1,794	1,441	1,188	2,678	2,291	919	2,611	1,894	2,886	3,615	954	2,091
1938	2,191	1,412	1,103	2,749	2,346	949	2,646	1,925	3,107	3,333	1,083	2,283
1937	1,734	1,148	1,029	2,394	2,079	743	2,425	1,675	2,579	3,007	965	1,827
1936	1,934	1,323	1,041	2,554	2,388	973	2,599	1,874	2,895	3,028	1,258	1,996
1935	1,676	1,072	1,022	2,236	1,990	703	2,029	1,641	2,346	2,644	1,007	1,561
1934	1,886	1,178	989	2,564	2,009	779	2,814	1,690	2,582	2,892	1,023	1,899
1933	1,713	1,156	952	2,308	2,058	796	2,373	1,765	2,790	2,754	988	1,728
1932	1,617	1,036	969	2,204	1,976	748	2,218	1,717	2,255	2,482	985	1,718
1931	1,855	1,161	944	2,371	2,065	825	2,454	1,851	2,500	2,845	1,069	1,932
1930	1,518	1,071	878	2,149	2,025	792	2,008	1,663	2,265	2,520	1,095	1,592
1929	1,520	1,055	849	2,110	2,015	772	2,471	1,718	2,374	2,510	1,050	1,787
1928	1,689	1,052	903	2,113	2,017	731	2,238	1,681	2,260	2,452	1,032	1,720
1927	1,473	1,016	871	2,095	1,921	725	2,261	1,614	2,294	2,404	895	1,598
1926	1,282	892	894	1,818	1,787	650	2,088	1,599	2,014	2,191	936	1,380
1925	1,537	879	764	2,074	1,644	702	2,241	1,485	2,094	2,243	820	1,503
1924	1,316	906	707	2,013	1,829	755	2,245	1,755	2,446	2,470	910	1,578
1923	1,594	949	754	2,025	1,860	690	2,303	1,622	2,380	2,468	960	1,565
1922	1,406	1,029	872	2,066	1,886	637	2,339	1,825	2,266	2,555	969	1,534
1921	1,640	932	899	1,975	1,954	695	2,063	1,888	2,501	2,539	982	1,599
1920	1,594	870	791	1,894	1,810	750	2,083	1,854	2,285	2,677	962	1,437

TABLE 17. - 1/ DEATHS FROM TEN SELECTED CAUSES BY PREFECTURE, JAPAN, 1920-1949 Cont'd
8/ SENILITY (162) - Cont'd

Year	Shiga	Shimane	Shizuoka	Tochigi	Tokushima	Tokyo	Tottori	Toyama	Wakayama	Yamagata	Yamanashi	
1949	870	968	1,859	1,216	1,036	1,599	730	664	972	1,017	1,591	599
1948	862	848	1,732	1,122	978	2,352	721	632	924	1,005	1,432	603
1947	1,091	1,314	2,359	1,216	1,301	2,779	929	766	1,264	1,128	2,083	794
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	1,287	1,634	2,880	1,355	1,551	4,937	1,078	951	1,574	1,381	2,172	870
1941	1,230	1,622	2,810	1,418	1,436	4,312	1,089	911	1,362	1,389	2,010	857
1940	1,281	1,727	2,684	1,373	1,424	4,043	995	1,014	1,325	1,484	1,865	829
1939	1,232	2,023	2,848	1,453	1,715	4,065	1,144	939	1,448	1,544	2,294	928
1938	1,486	1,973	3,084	1,545	1,699	4,338	1,072	996	1,322	1,443	2,428	938
1937	1,083	1,551	2,671	1,372	1,580	3,653	984	943	1,198	1,372	2,014	760
1936	1,274	1,777	2,996	1,350	1,666	3,679	980	926	1,340	1,333	2,243	893
1935	1,053	1,524	2,515	1,155	1,523	3,035	951	771	1,288	1,099	1,910	774
1934	1,278	1,795	2,855	1,397	1,561	3,121	1,038	1,046	1,346	1,326	2,153	898
1933	1,165	1,662	2,731	1,316	1,489	3,173	943	795	1,274	1,246	1,930	865
1932	1,035	1,597	2,568	1,123	1,383	2,721	966	828	1,107	1,188	1,861	744
1931	1,223	1,862	2,630	1,228	1,579	2,923	1,002	977	1,345	1,309	2,015	742
1930	1,072	1,934	2,309	1,062	1,380	2,461	922	757	1,196	1,089	2,010	703
1929	975	1,945	2,510	1,065	1,311	2,705	963	781	1,099	1,296	2,005	671
1928	1,002	1,781	2,618	1,081	1,367	2,807	848	732	1,239	1,196	1,845	643
1927	938	1,705	2,273	1,037	1,453	2,437	948	746	990	1,150	1,966	704
1926	795	1,609	2,035	1,044	1,211	2,124	727	669	930	1,077	1,699	625
1925	937	1,618	2,273	1,009	1,270	2,271	802	676	1,044	1,088	1,619	636
1924	933	1,763	2,175	1,035	1,457	2,412	917	673	1,117	999	1,961	681
1923	1,059	1,955	2,666	1,063	1,323	2,792	971	711	993	1,090	2,033	688
1922	963	1,725	2,233	1,051	1,449	2,620	894	761	1,140	1,147	1,986	708
1921	1,082	1,709	2,514	1,021	1,368	2,503	958	618	1,102	956	1,927	705
1920	952	1,657	2,327	981	1,322	2,470	891	625	1,098	1,037	1,918	711

See footnotes at end of table

TABLE 17. - 1/ DEATHS FROM TEN SELECTED CAUSES BY PREFECTURE; JAPAN, 1920-1949, Cont'd
ACCIDENTAL DEATHS (1949-1955)

Year	All Japan	Aichi	Akita	Aomori	Chiba	Ehime	Fukui	Fukuoka	Fukushima	Gifu	Gunma
1949	34,498	1,168	587	490	719	769	428	1,945	933	610	630
1948	38,960	1,131	610	600	803	567	3,812	2,005	957	586	676
1947	38,471	1,227	629	541	812	691	389	2,168	999	651	1,117
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	33,476	1,111	495	430	632	529	257	1,806	762	512	446
1942	30,959	983	441	387	608	500	252	1,737	703	506	447
1941	28,573	895	501	397	586	446	330	1,635	726	473	482
1940	28,147	958	525	416	595	439	310	1,907	767	520	467
1939	31,483	975	530	417	710	481	256	1,844	842	540	562
1938	29,951	1,003	529	425	539	502	304	1,738	791	509	593
1937	29,928	925	803	478	627	528	333	1,853	845	540	540
1936	28,721	942	490	423	557	559	291	1,753	877	504	766
1935	31,680	937	460	389	502	503	332	1,535	698	530	459
1934	29,817	845	483	433	554	543	283	1,458	741	516	391
1933	26,532	877	442	395	483	510	305	1,358	709	524	487
1932	25,055	845	405	386	514	527	262	1,359	713	448	404
1931	26,054	835	414	387	477	470	275	1,621	717	435	419
1930	26,412	862	477	342	478	517	260	1,480	774	511	415
1929	25,824	816	432	379	494	515	280	1,492	686	455	394
1928	25,102	830	432	381	528	490	325	1,433	881	484	402
1927	25,414	890	431	371	519	528	269	1,396	699	406	463
1926	24,779	862	395	344	469	514	289	1,378	734	443	407
1925	25,557	844	466	369	502	459	350	1,361	828	481	448
1924	71,166	851	480	350	1,785	480	290	1,567	759	501	439
1923	25,338	760	465	401	549	514	307	1,315	913	445	411
1922	24,294	759	410	299	529	448	304	1,262	705	454	466
1921	25,944	801	492	389	535	544	265	1,409	805	459	455

TABLE 17. - 1/ DEATHS FROM TEN SELECTED CAUSES BY PREFECTURE: JAPAN, 1920-1949 Cont'd
ACCIDENTAL DEATHS (169-195) - Cont'd

Year	Hiroshima	Hokkaido	Hyogo	Ibarak	Ishikawa	Iwate	Kagawa	Kagoehima	Kanagawa	Kochi	Kumamoto	Kyoto
1949	938	2,232	1,361	719	398	732	441	861	1,084	416	761	573
1948	1,012	2,211	1,358	699	437	1,119	479	700	1,084	418	676	615
1947	1,255	2,134	1,659	778	460	763	466	644	1,154	517	756	683
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	967	2,077	1,440	593	295	532	451	628	969	362	514	531
1941	857	2,173	1,288	657	272	486	338	566	970	357	442	500
1940	812	2,006	1,227	494	320	544	367	552	799	313	482	429
1939	790	2,146	1,320	551	290	652	336	551	791	365	485	461
1938	913	2,131	1,837	672	328	636	388	907	929	375	561	457
1937	899	2,070	1,327	587	325	672	395	616	866	375	493	538
1936	849	1,905	1,336	601	280	557	368	610	796	399	492	553
1935	789	1,847	1,340	533	276	557	463	473	743	374	520	552
1934	845	3,521	1,491	514	336	467	350	492	675	440	512	817
1933	751	1,795	1,234	557	333	2,900	343	469	741	345	581	515
1932	740	1,769	1,223	497	272	1,800	355	466	697	324	485	522
1931	683	1,664	1,094	489	312	1,493	376	437	635	340	465	507
1930	694	1,674	1,063	495	310	1,469	372	537	629	290	524	510
1929	735	1,774	1,134	544	326	451	397	510	650	324	536	518
1928	709	1,806	1,152	496	306	454	354	461	627	325	576	449
1927	692	1,813	1,156	495	302	431	366	492	538	287	888	3,383
1926	899	1,801	1,090	464	266	439	349	499	578	301	543	469
1925	735	1,689	1,484	493	271	438	341	398	581	294	490	440
1924	700	1,862	1,103	477	294	487	380	443	599	287	473	429
1923	895	1,867	1,250	541	283	515	317	445	16,403	282	528	494
1922	732	1,894	1,112	526	337	472	376	424	557	304	486	468
1921	764	1,736	1,130	461	289	464	323	456	520	273	457	471
1920	808	2,231	1,045	620	306	588	363	424	662	439	405	417

TABLE 17. - 1/ DEATHS FROM TEN SELECTED CAUSES BY PREFECTURE, JAPAN, 1920-49 Cont'd
ACCIDENTAL DEATHS (1920-195) Cont'd

Year	Mie	Miyagi	Miyazaki	Nagano	Nagasaki	Nara	Niigata	Oita	Okayama	Osaka	Saga	Saitama
1949	523	683	466	790	822	253	1,036	548	695	1,277	532	669
1948	519	715	443	735	821	262	1,026	548	985	1,342	569	728
1947	592	709	463	806	886	328	1,115	566	860	1,634	NA	1,103
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	497	532	402	657	767	217	956	447	635	1,415	403	572
1941	438	450	396	644	669	215	801	438	576	1,364	387	461
1940	428	418	325	628	671	215	912	422	561	1,526	325	558
1939	434	473	429	664	640	188	1,042	496	539	1,404	317	526
1938	437	479	372	704	636	194	1,006	423	427	1,524	336	604
1937	437	520	351	756	557	209	972	459	640	1,476	307	544
1936	440	468	389	721	610	201	974	477	622	1,422	364	548
1935	459	478	352	704	529	239	944	446	576	1,415	307	483
1934	431	432	335	729	588	227	926	392	662	9,128	305	459
1933	411	672	399	688	562	200	915	454	572	1,321	288	493
1932	441	448	295	719	462	240	899	389	588	1,352	212	493
1931	448	438	341	637	512	182	811	495	495	1,205	279	462
1930	438	433	353	666	594	193	889	447	565	1,114	354	478
1929	423	438	281	734	618	228	865	399	553	1,131	326	485
1928	420	474	303	758	509	189	854	486	557	1,115	368	476
1927	405	450	309	741	573	188	986	393	551	1,083	355	476
1926	400	419	262	681	558	207	991	468	554	1,084	345	451
1925	422	423	280	696	601	224	835	382	532	923	327	447
1924	361	405	295	703	602	205	851	431	557	1,001	329	472
1923	396	422	288	798	550	204	921	382	577	1,040	349	711
1922	385	454	253	667	554	199	1,070	394	526	1,034	326	466
1921	465	405	291	702	483	231	794	460	566	968	317	471
1920	394	419	277	669	409	187	904	338	587	875	322	450

TABLE 17 - 1/ DEATHS FROM TEN SELECTED CAUSES BY PREFECTURE: JAPAN, 1920-49 Cont'd
ACCIDENTAL DEATHS (169-195) Cont'd

Year	Shiga	Shimane	Shizuoka	Tochigi	Tokushima	Tokyo	Tottori	Toyama	Wakayama	Yamagata	Yamaguchi	Yamanashi
1949	263	414	885	566	409	1,900	236	570	338	582	888	358
1948	295	461	890	543	390	2,091	221	540	367	594	916	349
1947	395	429	940	936	448	2,001	288	553	477	633	952	331
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	251	349	797	454	394	1,877	226	390	344	490	1,274	274
1941	230	336	764	418	323	1,776	207	361	355	464	726	264
1940	258	312	768	416	341	1,820	186	436	332	533	654	286
1939	272	335	759	495	366	1,896	197	437	269	465	656	228
1938	266	378	840	513	410	1,896	225	459	349	506	688	280
1937	271	344	790	489	386	1,913	232	477	320	499	671	255
1936	260	374	812	508	372	1,733	200	474	320	547	631	243
1935	257	390	725	467	399	1,758	188	443	324	443	615	271
1934	315	360	701	454	418	1,708	296	400	363	451	570	225
1933	241	305	769	454	378	1,737	219	407	305	464	586	226
1932	248	324	693	438	331	1,814	193	370	316	509	536	242
1931	236	336	610	380	325	1,442	173	361	292	446	605	234
1930	248	329	862	444	360	1,518	223	384	270	447	579	249
1929	247	372	706	464	389	1,600	187	373	324	470	554	230
1928	223	345	624	447	366	1,526	184	435	259	476	511	261
1927	236	317	570	429	358	1,457	189	439	288	478	578	230
1926	226	324	633	396	399	1,305	211	345	296	483	614	252
1925	213	286	639	390	366	1,416	184	340	267	429	517	211
1924	230	324	603	440	393	1,529	203	393	259	497	613	219
1923	219	332	1,015	423	385	2,896	212	368	303	476	571	266
1922	240	349	660	440	342	1,114	195	375	288	444	551	242
1921	266	338	662	403	355	1,086	194	339	331	486	501	260
1920	231	351	635	529	407	1,176	172	349	271	520	516	306

See footnotes on next page.

TABLE 17. - DEATHS FROM TEN SELECTED CAUSES BY PREFECTURE, JAPAN, 1920-1949 Cont'd

FOOTNOTES:-

- 1/ Data for 1949 are provisional.
Data refer to deaths of Japanese nationals in Japan for the period 1920-1942 and 1947-1949. Prefectural death data are not available for 1943 and the all Japan total includes death in Okinawa and Karafuto. No death data for these causes are available for 1944-1946 except for the tuberculosis (all forms) estimates made by the Ministry of Welfare.
All data for the period 1920-1945 include military deaths in the country.
- 2/ Tuberculosis (all forms). 1920-1943; includes chronic pleuro pneumonia and all adenopathy; excludes spondylitis.
1944-1946; are estimates of the Ministry of Welfare. 1947-1949; excludes chronic pleuro pneumonia and nontubercular adenopathy; includes spondylitis.
- 3/ Intracranial lesion of vascular origin. 1920-1932; includes alcoholic cerebral apoplexy, congestion and a small amount of paralysis not due to intracranial lesions of vascular origin. 1933-1943; includes alcoholic cerebral apoplexy and congestion. 1947-1949; excludes alcoholic apoplexy and congestion.
- 4/ Diseases of the heart. 1920-1943; include diphtheric endocarditis. 1947-1949; excludes diphtheric endocarditis.
- 5/ Pneumonia (all forms). 1920-1943; excludes typhoid pneumonia. 1947-1949; includes typhoid pneumonia.
- 6/ Diarrhea, enteritis and ulceration of intestines (all ages). 1920-1943; includes malnutrition 1 year of age and over. 1947-1949; excludes malnutrition 1 year of age and over.
- 7/ Nephritis (all forms). 1920-1943; includes alcoholic nephritis. 1947-1949; excludes alcoholic nephritis.
- 8/ Senility. 1920-1922; excludes senility with mention of senile dementia, psychosis and paresis. 1923-1943 and 1947-1949; includes senility with mention of senile dementia. "NA" indicates data are not available.

SOURCES:-

1920-1943; Final Vital Statistics Annual Reports, Cabinet Bureau of Statistics
1947-1948; Final Vital Statistics Annual Reports, Ministry of Welfare
1949 : Monthly Vital Statistics Schedule Reports, Ministry of Welfare.

TABLE 18. - 1/DEATH RATES FOR TEN SELECTED CAUSES BY PREFECTURE: JAPAN 1920-1949
(Rates per 100,000 population)
2/TUBERCULOSIS, ALL FORMS (13-22)

Year	All Japan	Aichi	Akita	Aomori	Chiba	Ehime	Fukui	Fukuoka	Fukushima	Gifu	Gumma
1949	168.8	162.8	161.2	217.1	135.0	149.4	155.2	203.8	144.3	164.8	121.7
1948	179.4	171.9	177.7	228.0	143.4	166.8	164.5	216.6	153.5	195.0	144.6
1947	187.5	188.7	165.6	217.8	140.1	180.2	192.6	243.2	159.1	185.9	161.7
1946	264.2	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	280.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	240.2	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	230.8	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	219.1	210.9	183.4	232.3	171.9	225.2	240.9	237.1	155.5	209.5	156.3
1941	211.1	207.4	162.7	227.9	156.9	222.2	236.0	240.6	147.1	215.1	155.8
1940	212.5	213.2	162.6	230.7	160.4	225.0	253.7	231.1	157.0	222.3	177.6
1939	216.1	217.3	146.4	224.5	155.9	226.6	228.4	252.2	144.1	228.3	184.4
1938	209.2	220.5	144.7	195.9	136.0	219.8	239.1	228.2	144.5	235.8	174.1
1937	203.7	209.0	139.4	184.6	130.4	216.7	249.0	220.8	137.4	240.5	172.7
1936	206.7	218.6	140.0	203.3	136.3	216.3	235.3	211.2	141.5	234.3	174.9
1935	190.4	199.2	128.7	179.5	123.7	206.1	226.4	196.1	129.7	213.7	162.0
1934	192.4	201.2	130.2	195.1	133.7	207.6	248.1	198.7	143.0	220.2	163.7
1933	187.9	196.2	124.9	178.9	135.5	185.1	231.8	178.1	131.7	223.4	186.1
1932	179.4	188.9	122.6	175.0	129.1	177.3	246.4	172.3	126.9	211.4	160.0
1931	186.2	189.9	124.9	176.7	130.5	196.3	219.0	186.7	142.7	216.0	173.4
1930	185.3	185.1	117.8	160.9	143.1	190.4	227.1	190.8	138.7	213.1	182.4
1929	194.3	198.6	122.1	183.0	137.8	201.5	243.2	206.6	146.6	211.5	191.6
1928	190.6	201.9	111.8	178.4	140.0	192.9	224.8	200.7	143.8	208.7	184.0
1927	193.3	197.7	119.3	175.1	136.6	207.7	273.4	202.2	152.3	211.0	187.8
1926	185.6	184.1	114.5	177.3	136.4	202.1	256.0	189.9	146.0	196.8	178.3
1925	193.7	195.9	113.7	181.6	140.3	211.2	247.4	195.5	160.0	207.6	209.5
1924	193.7	202.4	117.8	182.5	153.1	208.0	233.1	190.2	162.5	214.1	201.8
1923	203.2	217.4	123.5	173.3	167.8	216.4	259.4	193.2	168.0	207.8	202.4
1922	218.9	217.5	127.7	202.2	183.9	223.0	296.8	206.4	183.0	212.6	207.3
1921	213.1	221.9	123.2	156.7	173.4	224.0	305.6	204.7	168.9	196.2	197.5
1920	223.8	248.0	127.2	161.1	177.5	244.7	289.7	219.1	173.7	219.7	198.6

TABLE 18. - 1/DEATH RATES FOR TEN SELECTED CAUSES BY PREFECTURE: JAPAN 1920-1949 - Cont'd
(Rates per 100,000 population)

2/TUBERCULOSIS, ALL FORMS (13-22) - Cont'd

Year	Hiroshima	Hokkaido	Hyogo	Ibaraki	Ishikawa	Iwate	Karawa	Kazoshima	Kanagawa	Kochi	Kumamoto	Kyoto
1949	158.5	249.6	169.3	110.5	169.0	206.4	146.1	153.1	178.5	153.0	149.5	214.1
1948	179.8	236.9	183.7	114.3	182.1	212.9	165.1	166.1	189.8	143.1	157.8	221.5
1947	190.5	231.8	209.3	113.0	193.0	189.8	182.9	159.2	198.0	161.1	169.7	246.4
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	235.9	285.9	250.5	127.2	202.5	195.4	235.5	245.7	229.3	210.8	198.5	282.9
1942	223.8	265.3	242.1	118.1	205.4	171.2	212.0	215.4	234.3	200.0	186.8	275.8
1941	207.9	269.7	245.9	123.6	328.4	157.3	222.3	204.8	241.2	191.7	166.9	266.4
1940												
1939	220.2	300.3	264.4	119.0	311.0	151.9	219.1	206.7	251.2	203.1	169.1	267.9
1938	202.2	275.2	254.1	117.2	340.9	141.9	200.9	194.4	238.2	186.7	160.6	265.5
1937	191.9	264.8	256.4	109.7	345.3	141.9	189.6	200.9	218.2	190.5	178.8	259.6
1936	197.1	268.0	260.5	112.5	349.2	143.9	194.2	186.3	229.6	175.6	177.6	269.4
1935	184.0	241.0	236.9	107.2	312.7	127.8	191.9	177.8	207.3	159.7	162.0	289.9
1934	183.4	238.4	235.9	107.3	301.8	138.7	181.9	166.6	218.1	153.7	165.8	256.3
1933	172.5	235.2	231.6	105.6	278.2	127.6	187.1	157.0	218.9	152.0	165.3	261.6
1932	162.0	213.9	220.5	104.0	291.0	119.3	171.9	151.2	205.4	144.6	163.1	241.9
1931	169.9	222.0	221.3	115.9	287.4	115.5	181.7	156.3	213.6	139.2	180.3	252.3
1930	171.1	216.2	223.2	109.9	291.2	112.5	188.0	162.3	206.2	146.6	181.5	248.7
1929	175.2	242.8	231.9	120.0	303.1	121.7	178.0	159.3	244.1	148.7	185.3	250.2
1928	176.9	223.5	222.5	119.4	274.4	112.7	189.8	156.7	228.8	145.0	180.7	261.6
1927	190.1	215.0	220.8	122.0	284.0	120.7	200.5	160.5	230.0	141.4	184.6	254.9
1926	176.7	204.5	222.6	112.6	280.7	119.9	184.4	155.1	209.3	146.2	189.0	248.7
1925	192.4	225.2	232.8	125.6	266.9	116.9	192.6	155.8	220.4	147.1	180.2	246.2
1924	202.2	188.3	228.0	129.1	277.7	117.4	210.8	150.8	217.9	139.7	189.1	253.4
1923	211.6	200.8	227.6	137.3	307.7	115.3	215.1	147.6	234.5	147.2	199.8	254.9
1922	222.2	222.0	244.4	142.8	338.0	134.2	216.4	150.9	274.1	151.1	196.5	272.6
1921	216.0	207.8	243.3	134.6	302.5	118.4	217.0	159.0	271.7	153.6	193.7	305.8
1920	251.4	198.0	257.2	148.8	299.6	124.3	245.3	153.6	275.0	149.9	201.0	322.1

TABLE 18. - 1/DEATH RATES FOR TEN SELECTED CAUSES BY PREFECTURE: JAPAN 1920-1949 - Cont'd

2/TUBERCULOSIS, ALL FORMS (13-22) Cont'd

Year	Wie	Miyagi	Miyazaki	Nagano	Nagasaki	Nara	Mifurata	Oita	Okayama	Osaka	Saga	Saitama
1949	150.6	164.4	167.8	122.9	169.4	154.4	169.8	176.1	144.5	201.5	172.2	133.6
1948	153.0	155.1	165.7	146.4	173.6	164.8	185.6	179.6	166.4	208.3	189.9	138.5
1947	175.4	183.7	183.3	144.9	201.6	193.0	182.8	212.3	171.1	234.3	212.2	137.6
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	217.5	177.9	202.6	156.2	210.3	201.4	189.6	244.6	214.2	264.0	218.9	150.8
1941	218.9	161.7	180.5	148.5	206.1	199.3	178.9	235.5	206.1	257.1	221.8	145.7
1940	166.8	157.4	168.2	153.3	199.4	191.7	184.3	219.4	206.9	261.4	205.1	153.4
1939	196.8	147.6	174.3	161.3	203.0	186.4	187.3	210.5	198.6	267.0	196.7	154.0
1938	197.0	150.8	151.3	155.8	196.3	185.2	187.0	201.4	179.5	258.5	197.9	155.8
1937	203.2	135.0	161.6	153.2	199.2	200.5	183.1	207.9	169.0	249.0	179.8	157.9
1936	199.3	153.7	161.2	152.9	210.0	207.1	185.4	204.6	172.0	268.2	187.2	169.1
1935	192.4	137.8	143.6	145.3	192.5	207.6	177.3	197.2	160.7	248.1	166.6	153.8
1934	185.7	143.6	135.5	144.8	198.2	202.8	186.9	197.7	162.0	250.0	165.2	160.3
1933	173.9	130.7	125.0	147.0	186.5	190.3	181.9	181.7	159.5	256.0	161.0	160.2
1932	175.6	134.6	126.4	136.8	179.7	171.0	175.4	184.1	147.6	242.2	160.5	150.2
1931	179.1	139.0	132.4	144.2	183.1	164.7	179.9	182.1	151.4	251.8	156.4	162.5
1930	171.0	133.7	124.8	138.5	192.6	168.7	172.2	183.4	154.2	251.0	172.5	165.4
1929	161.0	143.0	137.5	150.6	200.2	167.4	183.1	202.6	165.4	262.3	169.6	165.8
1928	189.5	137.1	127.7	151.6	199.6	171.2	185.5	188.7	165.6	255.8	156.8	166.0
1927	187.9	143.1	135.2	151.2	194.4	169.0	180.5	186.2	174.3	259.3	175.7	168.1
1926	173.1	134.7	126.8	146.3	185.6	158.5	185.3	189.9	163.1	236.6	167.2	170.1
1925	195.7	140.6	131.5	157.3	172.9	168.2	195.5	187.3	165.9	246.0	172.7	181.5
1924	192.9	139.5	117.7	158.4	171.2	173.8	201.0	198.1	169.9	244.7	180.6	186.3
1923	210.2	160.9	128.7	165.9	196.6	169.1	212.4	196.6	175.0	256.2	179.6	207.6
1922	215.1	167.8	120.9	172.8	196.9	184.3	226.1	209.2	184.9	209.2	191.8	214.0
1921	215.7	165.9	123.0	169.1	194.6	175.2	220.0	202.5	186.5	248.2	197.2	196.8
1920	223.9	157.0	128.1	183.0	208.0	189.5	223.9	222.0	196.7	316.8	196.8	209.9

TABLE 18. - 1/DEATH RATES FOR TEN SELECTED CAUSES BY PREFECTURE: JAPAN 1920-1949 - Cont'd

2/TUBERCULOSIS, ALL FORMS (13-22) - Cont'd

Year	Shiga	Shimane	Shizuoka	Tochigi	Tokushima	Tokyo	Tottori	Toyama	Wakayama	Yamagata	Yamaguchi	Yamanashi
1949	150.9	191.4	190.9	116.0	172.1	209.5	166.7	196.3	141.6	148.4	189.0	92.0
1948	165.2	211.6	144.2	129.7	174.9	219.1	187.7	187.9	155.9	173.4	210.4	104.3
1947	191.1	241.2	169.8	132.2	201.0	214.1	202.1	170.8	168.3	165.7	232.2	98.1
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	225.5	229.8	194.4	158.2	247.0	235.2	232.3	231.3	222.2	169.0	230.2	126.2
1941	222.6	276.4	180.6	150.2	250.6	231.7	228.3	224.5	197.8	149.4	225.3	118.7
1940	211.7	236.2	195.1	142.9	246.9	241.6	199.5	233.7	195.5	146.8	216.5	134.7
1939	206.3	204.4	204.3	149.4	245.6	255.8	211.2	232.5	199.7	138.4	234.8	128.2
1938	218.5	218.8	187.6	137.9	234.9	257.3	203.7	233.4	193.1	133.8	230.3	125.0
1937	220.6	208.0	197.3	143.6	235.1	230.6	188.7	247.5	193.2	129.4	230.1	121.2
1936	226.0	204.8	197.4	145.0	218.8	241.8	197.7	241.2	195.8	130.4	212.6	128.4
1935	209.2	202.5	172.7	124.1	209.9	218.7	177.2	204.7	194.7	122.1	196.4	122.5
1934	217.6	190.4	177.2	132.8	190.8	229.7	167.6	207.3	184.1	129.5	198.1	129.1
1933	211.3	182.9	169.2	126.4	194.5	236.5	154.3	197.0	169.4	131.0	189.3	122.7
1932	198.8	184.7	160.9	125.8	192.1	220.7	160.7	196.0	165.6	122.7	180.1	119.1
1931	201.7	188.3	167.2	138.4	207.9	233.3	165.0	191.6	177.2	131.8	191.1	127.6
1930	192.3	204.1	170.2	134.9	197.8	231.4	157.8	194.1	181.6	126.5	191.9	134.1
1929	217.2	208.1	172.8	140.6	215.7	245.2	169.6	196.4	173.2	120.9	192.6	136.5
1928	213.2	208.3	178.6	137.7	220.4	242.3	188.8	199.1	182.7	120.4	188.3	134.9
1927	214.3	208.1	177.7	137.9	212.0	251.0	160.5	111.2	183.7	119.7	190.8	130.9
1926	206.2	200.8	175.7	130.9	210.5	249.1	141.7	207.0	179.4	116.6	188.9	136.8
1925	225.8	196.8	182.7	144.9	229.8	251.6	158.6	207.0	165.4	126.4	192.2	136.8
1924	222.8	199.2	177.3	142.9	229.9	278.7	157.5	200.3	161.0	126.8	195.8	153.6
1923	249.7	219.5	192.2	146.7	223.6	296.8	171.8	211.8	163.2	149.5	208.8	140.9
1922	257.3	224.7	201.8	155.1	234.6	333.7	177.9	236.9	188.5	161.1	216.5	170.2
1921	251.7	226.2	197.7	157.3	230.4	347.9	156.1	203.4	194.7	142.9	213.6	165.6
1920	264.0	206.1	193.2	152.8	256.6	363.4	146.3	212.2	186.4	147.5	224.1	169.9

See footnotes at end of table.

TABLE 18. - 1/DEATH RATES FOR TEN SELECTED CAUSES BY PREFECTURE: JAPAN 1920-1949 - Cont'd
CANCER AND OTHER MALIGNANT TUMORS (45-55)

Year	All Japan	Aichi	Akita	Aomori	Chiba	Ehime	Fukui	Fukuoka	Fukushima	Gifu	Gumma
1949	71.6	71.6	57.4	53.2	82.2	73.1	73.7	77.8	64.2	67.0	71.6
1948	69.4	69.1	59.9	51.2	82.4	72.9	64.8	72.3	62.8	61.4	69.1
1947	68.0	63.5	51.8	48.1	79.5	78.9	62.6	67.5	67.4	64.6	69.1
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	70.9	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	72.0	63.1	61.7	56.0	87.7	85.2	71.1	70.0	73.2	67.0	75.0
1941	71.2	64.8	59.4	53.9	90.4	78.5	69.5	72.9	70.8	72.1	78.4
1940	70.8	60.7	61.5	50.1	87.3	74.2	72.3	70.3	67.7	69.6	77.7
1939	71.8	65.5	58.4	48.6	90.2	80.8	66.5	74.8	71.7	70.5	78.2
1938	71.1	63.9	61.1	49.0	86.7	82.3	67.5	71.7	69.3	66.6	78.3
1937	71.6	62.5	61.3	51.2	89.9	83.7	69.7	75.1	69.7	68.7	79.4
1936	70.4	61.9	59.6	49.3	88.2	80.1	61.3	71.2	65.0	69.4	73.0
1935	71.2	64.5	60.0	52.5	81.9	84.6	71.8	71.8	70.0	67.5	74.0
1934	70.5	64.0	60.3	55.2	88.1	74.8	67.4	69.8	64.6	70.0	70.7
1933	69.8	63.6	55.6	52.7	86.8	77.2	66.5	68.6	63.3	70.1	72.8
1932	68.4	62.4	50.5	52.4	85.6	69.5	62.0	67.9	67.0	64.2	71.7
1931	68.4	62.4	50.6	46.2	84.0	76.8	76.4	68.7	64.3	66.3	70.1
1930	70.0	64.1	54.6	49.6	83.1	74.5	68.6	70.6	68.4	69.8	70.1
1929	69.1	60.8	49.7	48.9	83.4	75.8	58.5	72.2	64.6	61.0	71.1
1928	71.3	63.7	50.8	50.8	86.5	80.4	64.9	72.2	69.1	63.9	68.4
1927	69.8	59.2	52.3	51.0	86.7	73.6	68.4	68.8	70.3	63.1	70.9
1926	70.3	60.9	54.2	52.9	85.0	76.1	68.2	71.8	69.3	63.5	69.8
1925	70.1	59.7	51.3	46.2	86.4	77.0	69.2	72.3	65.5	69.4	71.0
1924	70.4	60.4	54.7	44.0	90.4	77.1	69.3	63.3	68.0	63.7	73.7
1923	72.2	62.0	55.6	55.3	100.9	77.4	69.2	65.5	68.3	70.7	72.6
1922	71.4	61.5	56.3	54.6	91.0	73.8	65.2	66.6	69.0	63.9	67.4
1921	71.9	64.5	56.2	46.0	86.4	79.6	71.8	68.6	66.9	67.1	63.6
1920	72.4	60.6	57.0	45.6	91.1	76.3	66.3	74.2	71.0	66.6	67.8

TABLE 16. - 1/DEATH RATES FOR TEN SELECTED CANCERS BY PREFECTURE: JAPAN 1920-1949 - Cont'd
CANCER AND OTHER MALIGNANT TUMORS (45-55) - Cont'd

Year	Fukushima	Iokkaido	Hyogo	Ibaraki	Ishikawa	Iwate	Karawa	Yagoshima	Kanarawa	Kochi	Kumamoto	Kyoto
1949	74.3	59.3	70.2	74.9	83.4	47.9	80.3	55.1	66.8	68.7	66.8	77.6
1948	72.3	58.0	67.4	75.1	82.0	45.6	81.6	52.0	65.0	70.0	65.0	76.2
1947	68.7	55.1	64.3	75.6	77.0	43.5	80.4	54.8	59.3	68.7	62.6	80.7
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	75.8	52.2	64.4	85.1	90.2	50.8	94.7	63.3	60.1	78.6	74.9	87.8
1941	69.7	52.5	66.7	81.2	88.3	50.3	92.2	57.7	60.0	75.2	73.1	81.8
1940	71.1	51.1	65.9	81.9	91.8	49.4	90.0	57.2	57.7	67.0	73.4	82.8
1939	72.3	53.5	66.9	86.1	87.6	47.3	80.3	58.3	65.4	72.4	68.1	78.0
1938	69.1	53.5	68.2	86.6	79.7	50.6	84.1	56.6	64.4	73.0	69.6	78.2
1937	69.5	53.0	66.4	82.0	83.3	49.7	81.5	58.2	60.0	77.9	72.0	80.4
1936	70.0	52.6	67.8	81.1	80.4	41.3	87.4	55.2	65.1	74.5	70.1	76.7
1935	73.1	52.4	69.1	84.5	84.5	47.8	79.3	56.5	59.8	67.1	64.5	79.2
1934	73.2	48.9	68.6	84.6	89.9	46.7	81.3	53.1	65.5	62.9	67.3	77.0
1933	75.3	47.6	69.8	78.5	83.6	46.7	79.9	52.5	63.7	67.8	68.3	83.5
1932	66.7	46.7	66.4	83.9	80.6	47.7	75.6	52.1	58.2	69.3	65.1	76.5
1931	67.1	45.9	65.3	79.8	82.7	43.7	79.8	50.9	62.0	63.5	64.0	77.7
1930	69.3	49.5	67.0	90.0	81.7	43.0	72.6	54.4	60.6	61.7	64.3	80.2
1929	65.8	49.7	66.3	83.2	82.1	48.0	78.9	52.9	65.2	67.4	65.0	78.2
1928	70.3	50.4	68.9	80.7	85.3	48.8	75.2	51.5	66.8	63.4	69.8	81.6
1927	69.1	45.6	67.6	84.8	78.1	45.7	72.7	50.0	65.4	70.2	66.1	81.1
1926	68.6	48.5	67.8	82.2	76.6	39.2	76.6	49.4	64.9	70.2	67.9	80.8
1925	67.2	48.4	69.6	84.5	79.6	45.8	73.8	53.7	56.9	66.9	66.0	82.6
1924	72.1	39.7	66.3	87.2	80.7	45.5	71.3	50.5	63.9	61.8	73.6	83.8
1923	70.8	41.8	66.8	84.7	89.0	44.7	73.0	52.0	66.8	71.5	74.3	86.7
1922	71.9	45.2	63.2	83.0	80.9	45.1	78.6	49.9	65.4	70.7	71.9	82.9
1921	74.0	44.7	69.7	81.5	79.6	46.5	74.6	47.1	67.5	57.9	71.5	89.4
1920	73.2	44.9	67.1	86.2	77.3	47.1	75.5	47.8	66.9	67.1	72.4	87.8

TABLE 18. - 1/DEATH RATES FOR TEN SELECTED CAUSES BY PREFECTURE: JAPAN 1920-1949 - Cont'd
CANCER AND OTHER MALIGNANT TUMORS (45-55) - Cont'd

Year	Kie	Miyagi	Miyazaki	Nagano	Nagasaki	Nara	Niigata	Oita	Okayama	Osaka	Saga	Saitama
1949	73.8	73.7	61.0	80.0	65.4	99.4	83.1-	63.3	77.3	76.9	79.0	75.6
1948	72.8	70.4	58.3	78.4	61.2	101.5	84.1	69.1	82.1	75.6	86.6	71.9
1947	67.0	64.2	62.6	73.6	60.1	93.3	79.8	70.0	79.8	72.8	80.5	71.8
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	82.7	77.6	68.0	85.3	63.5	117.0	92.9	73.4	83.0	66.4	84.1	84.6
1941	84.1	77.7	63.1	85.6	60.7	120.8	89.1	68.3	86.5	65.5	81.6	79.3
1940	77.6	82.2	60.0	88.1	62.3	116.8	86.4	70.4	84.4	67.0	90.8	81.8
1939	88.2	76.1	59.8	84.7	62.0	120.2	90.5	74.1	84.7	67.0	88.0	83.7
1938	87.8	73.0	60.8	84.7	63.9	115.9	87.2	74.1	84.9	64.6	85.7	87.1
1937	83.7	72.8	63.9	85.7	64.0	118.1	88.1	73.0	86.2	67.4	92.9	87.0
1936	83.2	73.1	59.5	82.0	63.4	121.8	88.6	71.9	82.8	66.8	89.4	84.5
1935	84.6	81.1	58.8	85.2	64.2	122.3	87.9	72.4	82.3	70.8	80.3	84.6
1934	79.6	72.9	61.2	78.5	62.5	130.0	86.8	68.8	82.1	70.4	83.1	81.2
1933	84.8	73.2	57.0	76.5	58.1	127.6	85.0	71.6	76.9	72.6	86.9	76.9
1932	79.0	69.4	61.1	76.9	60.5	123.1	85.4	69.5	75.7	72.9	82.6	83.1
1931	79.8	68.9	62.3	74.9	58.5	128.6	84.4	68.4	77.2	72.2	88.7	79.4
1930	79.9	68.3	69.2	76.1	62.3	122.6	83.7	71.5	80.6	72.3	87.2	84.3
1929	85.2	69.2	65.3	81.2	64.2	117.0	81.6	58.2	79.4	72.8	79.0	78.3
1928	88.5	74.8	66.2	81.0	62.2	118.6	84.9	68.8	79.1	72.3	86.6	84.2
1927	85.4	74.0	61.7	77.6	63.9	124.7	85.1	64.1	77.8	73.7	78.2	83.2
1926	83.6	74.1	67.4	75.2	61.1	127.3	84.5	70.5	79.4	73.4	89.4	85.3
1925	83.8	71.4	62.1	75.4	59.0	125.4	86.3	68.4	71.8	72.9	86.4	79.8
1924	83.6	81.4	65.4	76.0	60.5	135.0	88.3	73.0	77.9	71.3	83.3	86.5
1923	89.3	79.8	63.5	77.6	62.8	137.3	86.4	75.0	79.2	71.4	94.6	86.5
1922	87.1	76.7	63.7	76.7	58.8	122.9	92.7	70.8	74.6	76.7	86.7	78.1
1921	84.6	79.8	58.5	73.7	67.6	130.7	87.6	66.0	75.3	79.5	84.1	84.9
1920	82.7	79.4	59.8	73.5	64.1	134.6	89.0	72.6	76.3	82.5	90.2	83.8

TABLE 18. - 1/DEATH RATES FOR TEN SELECTED CAUSES BY PREPARENCE: JAPAN 1920-1949 - Cont'd
CANCER AND OTHER MALIGNANT TUMORS (45-56) Cont'd

Year	Shiga	Shiwan	Shizuoka	Fuchii	Tokushima	Tokyo	Fotbori	Toyama	Wakayama	Yamagata	Yamaguchi	Yamanashi
1949	75.9	76.5	57.5	74.5	72.5	71.7	83.1	81.8	85.5	77.9	70.1	73.8
1948	73.2	76.9	56.5	72.7	66.6	66.2	83.6	80.7	82.7	72.7	67.2	71.7
1947	72.2	72.3	59.0	72.2	69.6	64.1	80.4	70.6	82.7	72.5	71.5	71.8
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	89.0	71.6	57.9	84.6	75.9	59.9	92.8	82.0	100.9	81.0	69.9	88.2
1941	88.3	74.1	60.6	87.1	79.1	59.1	94.0	78.6	98.3	77.6	66.8	86.0
1940	92.2	75.5	59.5	87.2	71.2	59.9	94.6	83.2	101.2	80.7	67.1	83.8
1939	87.6	81.6	58.6	79.2	69.2	60.8	77.4	85.0	94.3	78.3	75.0	83.4
1938	88.4	80.3	58.0	77.2	68.5	59.5	97.3	76.8	98.6	75.6	77.1	83.1
1937	82.4	74.6	55.2	84.9	68.5	59.8	87.5	84.5	91.8	75.5	76.9	81.0
1936	97.6	78.7	56.4	74.8	70.8	60.8	91.6	78.1	85.5	74.1	78.3	77.8
1935	87.0	82.3	58.7	76.1	67.9	59.4	84.0	73.5	88.8	77.0	76.4	76.2
1934	84.6	74.4	59.4	75.2	68.5	60.4	81.4	79.6	97.7	72.8	75.7	77.3
1933	94.3	72.5	56.3	71.7	65.7	59.0	72.7	75.9	91.1	71.9	77.1	77.5
1932	79.0	73.9	55.9	77.9	66.1	57.7	81.2	74.0	85.5	71.9	77.9	79.1
1931	86.2	71.9	58.5	71.2	59.7	58.4	80.8	76.6	93.0	74.4	72.2	71.0
1930	91.7	80.1	55.8	77.7	66.2	59.6	72.6	81.8	96.7	78.6	74.9	76.9
1929	82.8	80.7	55.1	78.4	64.7	59.0	84.7	80.9	91.7	74.0	73.4	74.2
1928	90.5	84.4	57.5	78.3	68.1	60.7	73.0	83.6	100.9	76.0	80.8	80.6
1927	89.9	79.5	57.7	71.2	69.0	58.3	79.6	77.0	99.0	74.1	76.3	71.2
1926	88.1	82.1	56.8	72.1	63.0	62.0	83.7	76.1	94.1	74.2	77.7	75.7
1925	94.1	78.8	57.2	79.8	69.1	61.0	90.0	79.7	93.6	75.6	78.7	69.1
1924	97.2	87.5	54.5	69.2	67.1	63.0	88.2	80.0	91.7	76.6	83.3	72.8
1923	97.4	87.2	61.0	71.2	72.0	60.7	97.5	86.4	99.5	80.2	87.9	71.6
1922	88.6	79.0	56.3	74.3	66.0	68.3	88.4	79.0	96.5	78.7	87.6	74.1
1921	92.1	82.4	57.4	77.6	64.3	66.5	96.9	76.7	95.7	74.5	89.5	77.3
1920	97.1	82.7	55.7	72.5	68.0	66.6	92.5	72.3	93.0	76.2	84.2	79.4

See footnotes at end of table.

TABLE 18. - 1/DEATH RATES FOR TEN SELECTED CAUSES BY PREFECTURE: JAPAN 1920-1949 - Cont'd
3/INTRACRANIAL LESIONS OF VASCULAR ORIGIN (83)

Year	All Japan	Michi	Akita	Aomori	Chiba	Ehime	Fukui	Fukuoka	Fukushima	Sifu	Yamaguchi
1949	122.4	118.2	188.2	177.1	152.3	106.3	120.0	105.9	145.4	122.8	133.5
1948	117.6	113.1	184.3	144.6	152.6	98.2	103.1	102.0	145.9	113.0	129.7
1947	129.6	127.3	210.9	141.3	157.6	120.2	108.5	120.8	159.9	127.5	121.9
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	162.8	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	170.8	133.9	321.7	180.0	221.6	154.5	174.0	156.4	222.5	176.9	181.4
1941	171.9	139.8	312.1	196.4	229.8	155.6	169.8	161.5	226.0	178.3	185.4
1940	176.3	142.1	328.8	193.2	229.8	167.6	186.8	167.4	228.4	179.7	185.2
1939	184.1	141.9	315.4	188.7	222.9	185.0	177.6	192.7	229.3	183.3	190.9
1938	178.2	148.9	302.9	184.4	227.3	163.5	165.1	189.2	221.3	177.8	185.6
1937	168.3	134.0	280.8	164.0	226.9	136.4	162.6	183.6	210.1	173.2	165.6
1936	168.3	136.4	276.6	158.4	217.9	157.9	165.0	180.4	203.6	177.7	168.6
1935	166.4	132.5	270.7	157.0	212.0	155.5	152.9	176.3	205.1	167.2	165.4
1934	168.7	140.5	280.9	157.1	214.7	152.3	165.3	172.9	215.1	172.5	163.4
1933	165.3	132.0	276.2	151.4	218.4	157.7	161.7	164.5	205.7	161.8	162.1
1932	168.9	133.1	277.6	143.4	216.2	159.4	163.6	162.9	206.8	165.2	164.8
1931	165.4	136.2	268.3	156.0	222.6	180.3	158.4	159.9	213.8	164.0	164.0
1930	164.0	126.7	269.6	150.4	217.3	154.9	146.1	168.2	211.5	153.7	164.2
1929	172.0	138.8	304.8	151.7	230.3	154.7	150.6	177.1	213.6	162.0	168.2
1928	165.5	138.1	292.6	144.9	227.2	151.9	141.3	167.1	201.9	152.1	162.0
1927	166.1	131.6	286.6	142.3	218.6	150.6	134.8	167.7	211.8	155.2	163.8
1926	163.6	132.3	288.9	146.3	224.6	153.2	143.7	160.5	209.1	155.2	158.8
1925	162.4	130.4	281.5	145.3	219.3	143.3	135.3	156.1	214.8	158.7	163.1
1924	176.0	136.3	311.2	166.2	231.5	164.7	148.5	159.2	226.2	166.1	178.3
1923	163.9	140.2	277.5	137.7	242.0	145.2	136.3	152.6	216.2	157.2	158.5
1922	160.6	127.2	283.0	158.8	225.9	139.6	127.5	155.6	221.0	147.6	154.3
1921	161.0	141.8	265.7	140.8	223.1	148.0	136.8	152.9	220.8	149.4	150.4
1920	158.8	139.9	264.2	139.7	208.7	145.2	120.3	166.5	217.3	152.3	144.1

TABLE 18. - 1/DEATH RATES FOR TEN SELECTED CAUSES BY PREFECTURE: JAPAN 1920-1949 - Cont'd

3/INTRACRANIAL LESIONS OF VASCULAR ORIGIN (83) - Cont'd

Year	Hiroshima	Fokkaido	Hyogo	Itaraki	Isikawa	Iwate	Karawa	Karoshima	Kanagawa	Kochi	Kumamoto	Eyobo
1949	115.2	89.1	96.2	159.2	122.0	181.5	116.6	128.4	105.9	144.6	123.4	94.7
1948	107.1	80.3	92.5	156.2	114.3	183.4	107.5	121.4	99.3	134.1	116.1	87.1
1947	124.1	82.5	108.5	161.3	132.8	192.2	124.2	129.9	103.5	154.7	129.0	109.5
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	169.1	117.3	134.9	218.4	190.5	242.6	180.7	181.2	141.8	203.4	208.7	141.9
1941	168.4	120.6	133.1	229.1	199.0	257.4	174.3	166.9	145.8	187.7	191.7	135.4
1940	172.6	126.5	136.3	224.7	206.4	251.0	195.2	175.6	147.4	189.7	198.3	146.8
1939	189.3	134.6	150.5	233.3	194.5	242.4	189.1	167.7	167.4	204.8	185.4	148.7
1938	187.3	112.4	146.9	221.5	203.0	244.2	181.9	164.2	152.5	197.3	202.3	142.8
1937	166.4	116.3	143.3	206.9	199.9	233.2	166.1	155.5	151.3	191.4	190.8	135.7
1936	172.2	111.1	139.1	200.9	199.5	230.4	167.7	158.4	149.3	194.8	179.4	126.9
1935	161.5	107.4	140.5	212.8	184.4	224.4	167.0	158.6	149.7	222.1	184.7	139.2
1934	165.2	100.8	140.9	223.1	195.4	237.1	154.4	153.1	154.0	182.0	175.1	144.4
1933	162.3	101.2	144.5	218.9	190.2	218.8	152.6	145.3	145.3	177.7	172.3	142.6
1932	172.7	92.7	134.3	216.3	179.3	222.9	155.9	144.9	146.2	171.2	164.5	138.9
1931	164.8	99.5	134.6	234.2	174.1	228.2	159.5	143.3	148.5	187.5	167.7	147.7
1930	172.0	100.2	138.2	221.0	169.2	231.2	151.2	158.9	145.3	177.4	167.9	139.7
1929	180.2	102.9	143.1	234.3	174.9	234.0	174.7	148.9	164.0	196.0	169.5	155.1
1928	169.6	96.9	137.8	229.1	164.0	242.8	148.8	141.2	156.9	175.8	182.3	147.4
1927	173.1	96.3	143.1	225.7	183.2	240.9	172.4	146.7	156.1	184.5	156.1	152.4
1926	178.3	94.6	142.6	223.2	182.8	244.6	149.6	138.5	152.6	187.1	181.1	144.3
1925	160.9	95.0	132.6	231.0	183.8	250.6	134.8	142.5	155.4	174.7	163.3	142.5
1924	199.3	81.9	151.7	235.6	201.6	278.9	198.6	148.2	148.0	201.4	167.1	158.9
1923	166.5	79.8	133.1	230.3	166.9	261.9	148.6	142.5	155.3	180.7	160.9	145.4
1922	163.7	76.3	130.2	219.7	191.9	268.8	148.4	143.1	147.6	172.3	151.2	138.0
1921	161.7	76.5	128.7	218.7	164.5	257.2	142.9	145.8	146.5	175.5	161.7	132.1
1920	159.5	81.5	132.7	225.9	151.3	248.7	148.6	150.1	142.5	188.4	157.5	144.5

TABLE 18. 1/DEATH RATES FOR TEN SELECTED CAUSES BY PREFECTURE: JAPAN 1920-1949 - Cont'd

3/INTRACRANIAL LESIONS OF VASCULAR ORIGIN (83) - Cont'd

Year	Mie	Miyagi	Miyazaki	Narano	Narasaki	Nara	Miyata	Oita	Okayama	Osaka	Saga	Saitama
1949	120.2	139.6	118.6	151.2	103.5	120.3	159.7	132.9	122.8	84.3	127.5	137.5
1948	113.7	134.6	104.6	148.1	96.0	104.8	162.1	126.7	121.3	79.4	115.4	133.1
1947	132.0	136.4	120.7	156.3	110.6	137.8	178.9	146.7	141.5	97.0	131.6	134.5
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	181.3	203.3	176.1	237.9	133.6	203.9	236.0	214.3	199.0	112.0	203.6	187.4
1941	180.5	197.7	178.7	229.4	136.0	207.3	227.6	210.3	183.4	108.4	189.7	201.7
1940	175.9	205.0	178.0	240.3	144.0	204.4	235.0	206.9	200.9	121.4	196.7	205.5
1939	183.6	193.2	179.3	246.0	162.2	219.1	240.8	217.3	207.6	129.3	199.4	207.4
1938	178.5	188.7	172.1	231.4	156.0	203.2	236.7	211.0	198.5	120.9	202.2	209.2
1937	168.7	186.1	159.4	225.7	145.0	183.6	219.0	193.4	186.4	113.8	202.4	188.8
1936	167.7	183.0	167.0	220.6	140.3	190.1	214.4	198.2	191.7	122.6	192.9	198.5
1935	163.5	177.4	164.6	216.5	139.3	197.3	209.4	196.9	190.4	121.6	182.2	190.8
1934	161.6	184.4	149.0	213.8	126.2	199.3	223.8	201.7	184.9	130.1	184.8	204.4
1933	152.6	180.0	161.0	208.3	124.2	188.6	214.3	191.4	180.6	130.2	186.4	192.9
1932	159.8	179.2	164.6	210.2	121.6	186.4	201.1	199.6	179.7	123.1	173.9	188.5
1931	152.2	186.4	159.4	214.5	122.4	196.1	201.5	198.0	187.0	125.2	185.6	198.7
1930	148.7	197.1	165.4	202.6	124.4	182.1	193.4	188.5	184.7	128.4	175.5	179.8
1929	152.9	199.0	155.2	221.4	122.4	190.6	211.7	191.4	194.8	130.3	183.4	191.6
1928	161.2	199.8	148.5	210.0	126.7	171.9	186.8	177.0	185.7	128.4	172.3	187.0
1927	152.3	199.6	161.3	206.8	125.9	169.0	204.7	185.7	198.3	126.5	172.4	179.0
1926	149.3	193.3	153.4	205.2	115.7	164.9	210.1	174.9	197.0	128.9	173.1	177.3
1925	153.7	221.8	161.5	207.3	114.6	169.2	200.4	174.6	179.0	123.3	165.0	179.6
1924	150.8	242.2	165.6	227.0	117.7	178.4	227.5	191.0	214.1	128.4	182.6	179.6
1923	150.8	224.6	156.7	209.3	117.2	165.2	201.3	176.6	184.2	122.3	174.7	173.2
1922	182.6	223.7	163.4	204.9	109.9	164.6	197.1	173.1	187.0	116.4	182.6	162.1
1921	146.6	221.6	167.7	202.3	107.7	173.4	196.6	193.6	181.6	125.7	170.2	160.3
1920	141.4	213.0	154.0	197.0	100.0	164.7	200.8	181.9	177.1	124.3	172.6	154.4

TABLE 18. - 1/DEATH RATES FOR TEN SELECTED CAUSES BY PREFECTURE: JAPAN 1920-1949 - Cont'd

3/INTRACRANIAL LESIONS OF VASCULAR ORIGIN (83) - Cont'd

Year	Shiga	Shimane	Shizuoka	Tochigi	Tokushima	Tokyo	Tottori	Toyama	Wakayama	Yamagata	Yamaguchi	Yamanashi
1949	121.1	148.6	116.9	142.3	108.0	94.4	133.8	127.8	116.8	172.6	118.3	122.9
1948	117.0	133.1	109.1	140.6	106.6	86.5	130.4	127.4	109.4	170.6	115.4	119.5
1947	142.9	149.1	119.2	136.0	125.4	86.3	138.2	132.7	122.4	185.2	145.9	126.6
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	188.6	202.7	166.0	218.7	163.7	122.0	201.9	202.2	197.1	242.7	194.6	174.6
1941	193.1	232.3	169.5	213.9	168.4	124.7	207.6	185.8	182.6	258.4	192.6	183.2
1940	197.7	230.6	171.7	221.6	169.2	133.9	208.1	211.4	192.2	267.6	204.2	188.1
1939	198.6	241.3	165.6	207.8	178.1	143.8	241.0	201.1	196.1	249.7	230.1	190.2
1938	189.1	241.7	168.1	206.8	166.7	139.1	209.4	205.5	184.8	247.9	237.2	185.8
1937	184.1	215.6	156.0	186.8	159.1	125.8	195.4	190.4	175.5	235.0	215.6	163.0
1936	196.0	223.9	160.1	194.2	168.2	130.2	191.3	193.5	171.2	238.3	214.3	176.5
1935	180.6	202.6	153.5	185.3	172.6	129.1	195.5	184.8	180.8	222.1	203.1	179.1
1934	199.8	216.4	156.9	194.2	156.6	128.8	186.1	186.4	174.6	235.0	198.9	160.9
1933	173.1	213.5	150.9	195.6	152.8	130.3	184.4	177.3	170.1	222.7	197.1	172.0
1932	181.4	217.3	152.8	189.8	148.8	128.0	179.6	177.7	155.0	219.5	198.4	163.7
1931	185.7	196.1	155.1	194.3	150.0	134.4	174.9	187.2	160.4	225.2	189.4	153.4
1930	184.9	210.3	147.7	188.4	143.3	132.0	168.4	165.4	169.4	219.7	197.2	158.8
1929	199.5	217.5	154.8	199.7	164.9	138.8	182.3	182.2	163.9	218.0	197.3	164.8
1928	193.7	194.7	153.6	195.0	145.6	136.5	179.9	167.7	164.2	216.3	186.6	158.7
1927	181.1	192.7	144.2	180.1	164.4	135.5	173.4	164.9	158.5	217.9	190.0	152.8
1926	186.1	187.1	149.6	189.7	147.7	131.2	159.7	161.2	151.4	208.1	186.5	150.7
1925	177.4	182.6	145.9	186.2	141.2	130.2	148.7	155.8	147.3	208.8	188.6	139.8
1924	197.9	219.6	154.3	187.2	172.2	142.2	189.6	211.0	171.5	218.3	229.9	155.4
1923	190.9	215.6	150.4	174.2	140.3	137.5	169.6	177.8	153.2	200.3	209.3	135.2
1922	172.5	172.8	144.9	170.3	145.4	134.9	154.8	186.7	143.6	207.3	194.3	136.6
1921	190.7	184.1	147.8	171.6	150.4	132.1	158.3	158.8	152.2	192.3	182.2	130.8
1920	176.0	159.5	147.1	179.7	134.4	131.8	142.1	174.8	146.3	189.7	188.3	127.2

See footnotes at end of table.

TABLE 18. - 1/DEATH RATES FOR TEN SELECTED CAUSES BY PREFECTURE: JAPAN 1920-1949 - Cont'd
4/DISEASES OF THE HEART (90-95)

Year	All Japan	Aichi	Akita	Aomori	Chiba	Ehime	Fukui	Fukuoka	Fukushima	Gifu	Gumma
1949	64.3	66.5	69.3	53.0	74.9	63.8	69.2	54.4	64.4	67.3	75.0
1948	61.2	61.8	68.0	53.0	73.2	64.5	58.7	49.3	66.6	64.3	70.2
1947	62.3	65.8	66.2	47.8	69.0	62.5	57.6	52.9	63.9	76.2	69.5
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	61.1	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	59.1	56.2	64.4	50.8	80.0	60.9	71.3	45.9	63.0	66.9	74.8
1941	58.2	55.3	67.7	55.9	70.4	57.6	74.7	46.7	60.9	65.1	71.2
1940	63.3	60.3	76.7	54.7	87.3	66.5	82.8	52.8	62.6	72.5	69.8
1939	66.5	63.1	67.6	58.0	89.0	65.5	80.9	59.9	69.9	70.0	71.1
1938	66.8	65.1	63.6	51.4	95.0	70.6	82.9	61.4	68.3	77.8	74.9
1937	60.5	60.4	62.1	45.9	84.1	58.7	82.5	54.0	56.5	68.5	66.4
1936	61.3	59.4	58.4	62.3	75.9	65.4	71.9	54.8	60.3	74.2	66.6
1935	57.7	55.6	58.7	54.1	77.2	61.5	75.9	47.8	59.1	68.3	56.8
1934	62.5	60.2	54.9	58.1	79.6	66.3	91.3	50.7	65.7	80.6	67.2
1933	59.7	55.4	65.3	47.5	80.6	60.9	87.7	48.3	59.5	70.4	58.9
1932	58.9	55.3	63.0	50.4	78.7	56.8	82.8	50.7	61.2	71.0	58.6
1931	64.3	60.4	63.7	54.8	81.5	68.4	90.3	54.3	62.8	76.3	59.9
1930	64.1	55.8	69.7	55.6	83.9	61.1	91.6	55.6	65.4	83.3	62.6
1929	65.7	61.1	88.2	66.8	85.2	60.2	90.3	51.7	65.9	73.9	60.8
1928	63.9	65.1	70.2	57.5	81.2	64.1	81.9	50.9	61.4	79.2	67.0
1927	62.5	60.7	74.3	53.4	77.6	66.0	85.2	52.1	58.2	73.0	63.9
1926	62.8	60.8	76.5	65.6	83.3	61.8	76.5	49.1	57.9	67.2	58.4
1925	67.1	66.7	81.3	62.4	84.4	67.3	79.4	52.0	63.2	75.8	66.0
1924	69.4	69.0	79.4	68.5	90.4	70.3	75.7	50.1	71.6	78.7	67.3
1923	72.9	74.9	112.2	78.3	97.8	74.8	88.4	50.5	72.0	86.2	67.5
1922	65.4	61.6	86.4	73.8	90.1	63.8	85.6	47.8	63.0	77.7	63.7
1921	66.5	71.0	77.6	62.9	85.7	70.1	75.8	50.3	63.3	77.5	64.2
1920	63.9	61.9	74.5	62.0	77.2	62.4	68.3	48.8	63.6	71.7	59.3

TABLE 18. - 1/DEATH RATES FOR TEN SELECTED CAUSES BY PRE-STRUCTURES: JAPAN 1920-1949 - Cont'd

4/DISEASES OF THE HEART (90-95) - Cont'd

Year	Firoshima	Hokkaido	Hyogo	Ibaraki	Ishikawa	Iwate	Kagawa	Kagoshima	Kanagawa	Kochi	Kumamoto	Kyoto
1949	63.1	54.6	57.5	77.8	73.7	65.2	65.2	62.0	53.2	51.4	67.7	57.6
1948	67.0	49.7	52.6	73.4	68.2	65.8	62.7	58.6	50.0	47.1	61.4	59.0
1947	69.3	46.1	59.4	70.1	71.9	58.7	64.7	58.6	49.6	50.9	65.3	62.6
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	66.6	42.6	48.1	73.0	70.3	67.9	68.0	58.5	51.1	54.6	63.1	58.0
1941	61.3	47.3	47.4	69.5	67.9	67.9	56.7	54.7	50.5	47.2	62.6	53.8
1940	71.7	53.8	52.1	73.8	86.3	73.2	67.2	56.9	52.9	55.8	68.1	64.5
1939	69.6	59.2	58.3	76.8	86.3	78.6	74.9	56.1	57.8	55.5	71.5	59.6
1938	73.5	53.2	60.7	74.7	90.2	67.2	72.2	60.3	58.1	55.3	74.0	58.9
1937	66.4	49.8	53.8	66.5	86.1	64.8	65.4	51.9	52.0	47.0	63.9	56.4
1936	66.7	50.8	54.8	64.0	83.7	65.6	66.2	53.3	54.1	55.5	67.4	52.1
1935	62.6	45.8	50.8	60.9	87.8	57.5	65.2	50.7	54.3	56.2	61.7	52.1
1934	67.1	46.3	58.1	65.0	96.4	62.5	73.8	52.6	59.0	53.7	62.2	63.4
1933	67.9	47.8	56.5	66.8	87.3	64.0	72.6	44.1	57.5	51.1	59.8	54.2
1932	65.7	45.6	59.4	67.7	87.6	56.5	60.1	49.4	55.4	46.8	56.9	56.4
1931	71.3	48.2	59.2	71.2	86.5	64.1	73.6	50.3	61.0	50.9	67.7	63.7
1930	73.6	48.3	60.5	64.3	91.8	66.6	69.2	55.8	59.2	57.6	61.2	63.6
1929	74.2	58.2	62.7	70.5	100.1	63.9	74.3	50.3	65.6	52.9	63.6	66.9
1928	71.7	47.3	61.5	71.3	94.4	63.7	74.1	53.0	62.2	61.1	57.4	63.0
1927	73.6	51.9	60.7	68.1	92.3	66.2	71.5	48.3	63.4	63.6	58.1	65.2
1926	75.1	48.4	58.7	65.1	92.4	66.6	70.8	58.1	60.6	64.4	62.1	65.6
1925	82.6	47.6	68.6	73.7	99.1	72.7	73.1	50.5	65.9	65.2	57.3	75.4
1924	92.2	41.8	63.8	75.1	104.0	69.5	81.5	57.4	68.7	70.9	65.0	77.1
1923	90.4	46.7	60.8	72.0	117.7	72.2	77.5	60.8	70.3	65.6	62.6	82.1
1922	80.9	38.6	50.8	62.5	105.5	68.6	69.6	55.2	62.9	68.0	55.6	77.1
1921	82.2	41.3	61.2	66.3	95.0	61.9	69.2	57.5	66.0	64.0	65.6	74.2
1920	78.4	42.8	59.8	66.1	85.9	56.2	64.8	56.8	56.7	63.1	58.1	75.4

TABLE 18. - 1/DEATH RATES FOR TEN SELECTED CAUSES BY PREFECTURE: JAPAN 1920-1949 - Cont'd

4/DISEASES OF THE HEART (90-95) - Cont'd

Year	Mie	Miyazaki	Nagano	Nagasaki	Nara	Niigata	Oita	Okayama	Osaka	Saga	Saitama
1949	69.8	66.4	71.3	62.0	65.2	66.2	71.3	63.7	59.6	64.5	73.6
1948	68.4	59.0	55.2	60.8	65.0	65.7	65.1	60.7	56.3	67.9	65.7
1947	68.0	59.6	60.5	56.8	71.0	64.7	64.4	59.5	62.2	68.0	63.5
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	72.8	70.6	56.0	70.6	72.7	69.5	70.5	56.9	48.2	65.4	67.0
1941	68.5	67.5	54.5	58.7	69.6	64.5	64.5	58.6	43.8	64.8	64.8
1940	72.2	64.7	55.3	63.4	77.6	71.8	65.4	62.5	50.4	70.8	70.2
1939	72.5	67.1	66.9	70.2	77.2	76.8	69.9	72.6	52.6	75.9	73.3
1938	84.5	62.7	66.0	71.0	69.1	67.3	71.0	65.9	53.1	73.4	72.2
1937	71.5	59.0	57.3	62.7	70.0	64.6	70.5	62.2	50.1	60.5	64.0
1936	69.7	65.8	53.2	63.1	68.6	70.4	68.7	63.1	50.4	66.7	63.5
1935	65.0	59.8	55.7	55.4	70.1	64.5	62.7	61.1	48.9	65.1	61.7
1934	71.3	61.4	52.6	56.2	80.2	72.1	68.0	65.0	57.2	63.9	65.6
1933	64.3	58.9	55.1	63.3	66.9	61.5	66.6	57.6	57.6	51.5	66.6
1932	63.7	59.1	55.3	52.9	72.8	61.2	66.2	61.2	53.9	61.4	63.4
1931	79.6	58.2	67.2	67.2	75.8	69.6	70.4	64.3	59.5	59.2	66.4
1930	69.9	60.5	60.5	57.9	74.3	68.8	70.2	63.4	62.3	67.4	61.1
1929	71.5	62.9	63.2	59.0	70.4	77.1	67.2	66.3	57.3	62.2	62.9
1928	79.3	62.0	63.3	56.0	66.9	68.5	61.6	63.8	59.6	60.3	58.6
1927	72.8	67.3	56.6	54.4	64.5	73.3	70.2	66.1	59.3	51.6	62.7
1926	70.4	59.2	63.1	51.4	63.1	63.2	61.2	61.2	57.9	56.0	62.0
1925	74.7	65.5	59.2	52.9	75.5	78.4	57.9	66.7	64.6	61.5	65.8
1924	77.3	72.5	60.9	54.2	77.1	82.0	66.6	70.0	67.3	63.4	69.5
1923	88.4	77.9	61.9	59.4	61.6	83.8	69.9	78.1	70.1	64.9	73.5
1922	71.6	70.4	55.7	53.4	59.6	74.2	69.2	69.5	68.1	63.4	59.5
1921	76.6	69.2	63.6	53.8	67.5	69.3	73.4	70.0	69.6	60.6	63.7
1920	67.3	69.5	59.9	57.5	60.8	71.0	72.3	67.5	70.1	58.8	63.0

TABLE 18. - 1/DEATH RATES FOR TEN SELECTED CAUSES BY PREFECTURE: JAPAN 1920-1949 - Cont'd
4/DISEASES OF THE HEART (90-95) - Cont'd

Year	Shira	Shirane	Shizuoka	Tochigi	Tokushima	Tokyo	Tottori	Toyama	Wakayama	Yamagata	Yamaguchi	Yamanashi
1949	73.5	80.2	59.9	68.0	76.6	56.4	62.7	68.4	59.5	72.1	63.1	73.8
1948	74.3	70.6	59.7	67.6	79.5	48.6	62.1	68.3	57.9	71.9	60.5	77.1
1947	84.2	75.6	62.6	66.7	92.4	45.6	63.7	65.0	63.2	71.1	70.9	71.4
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	84.6	75.7	58.5	68.5	80.6	42.2	64.3	64.6	62.0	75.6	61.7	74.8
1941	79.6	75.6	59.0	72.4	79.7	45.6	66.5	62.5	55.0	77.4	59.4	74.2
1940	83.7	85.3	62.0	70.3	87.9	47.8	68.5	77.3	64.3	79.7	65.7	73.3
1939	87.3	84.3	65.1	73.9	90.9	52.3	74.9	75.9	68.7	76.6	79.9	77.6
1938	93.6	84.7	69.1	70.9	85.1	53.1	80.5	89.4	68.1	74.1	81.9	76.7
1937	76.6	81.1	61.2	64.7	78.4	47.9	69.3	81.6	57.4	65.0	76.6	68.4
1936	87.1	77.5	61.0	57.2	87.2	45.8	73.4	75.9	56.9	64.7	71.9	75.8
1935	76.3	72.0	60.8	57.3	76.0	45.0	63.2	73.9	59.0	64.0	69.8	69.4
1934	90.8	78.1	58.6	59.5	85.7	45.8	70.5	81.0	61.5	64.4	72.7	73.5
1933	79.4	83.7	63.6	63.5	83.0	46.5	60.5	72.2	60.1	61.2	71.2	65.2
1932	70.3	85.8	61.6	57.2	80.2	43.5	62.6	71.9	58.7	57.0	74.7	68.8
1931	84.2	88.6	63.7	62.6	98.4	49.4	61.1	80.9	63.5	67.9	80.1	69.9
1930	87.8	95.2	64.9	62.9	82.9	48.1	64.6	90.7	63.4	67.2	84.1	68.0
1929	63.8	91.6	66.0	60.3	85.7	53.2	69.2	87.2	61.5	68.4	73.1	68.4
1928	83.8	84.7	63.7	59.6	91.9	51.8	65.7	75.8	63.0	63.0	71.7	72.0
1927	76.6	92.8	63.6	53.9	89.2	51.5	74.9	85.9	61.9	60.7	70.6	63.6
1926	91.2	91.9	64.5	54.4	78.5	50.1	62.8	82.4	57.9	69.4	68.8	62.3
1925	83.0	97.5	67.1	65.5	89.4	52.4	79.0	81.8	50.4	73.8	70.0	69.8
1924	77.6	97.0	69.5	66.6	90.6	56.9	76.5	86.4	62.8	78.8	76.6	64.5
1923	86.5	111.4	71.7	67.2	94.6	59.2	77.6	92.7	61.2	79.6	78.0	61.9
1922	77.3	77.0	61.7	59.5	91.6	55.8	70.1	83.3	59.0	66.8	73.5	64.4
1921	73.6	81.5	68.1	58.5	93.8	55.5	70.7	83.0	59.9	65.9	74.6	58.4
1920	73.4	82.7	62.3	61.3	83.2	55.1	57.6	75.4	54.8	67.1	67.3	65.8

See footnotes at end of table.

TABLE 18. - 1/DEATH RATES FOR TEN SELECTED CAUSES BY PREFECTURE: JAPAN 1920-1949 - Cont'd
5/PNEUMONIA, ALL FORMS (107-109)

Year	All Japan	Aichi	Akita	Aomori	Chiba	Ehime	Fukui	Fukuoka	Fukushima	Gifu	Gunma
1949	68.9	68.3	63.3	92.0	71.5	71.3	65.2	73.9	86.3	62.8	81.4
1948	66.1	55.9	66.2	96.9	61.2	65.7	60.7	60.5	68.7	59.1	76.7
1947	130.3	124.9	106.5	136.1	117.6	134.8	121.7	124.5	129.5	128.3	132.3
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	156.8	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	143.6	130.6	156.3	208.1	158.2	128.4	142.9	147.7	154.1	121.3	145.9
1941	142.0	159.2	150.9	188.5	153.1	136.0	190.9	157.1	140.2	155.4	160.1
1940	153.2	162.4	163.2	200.3	160.0	156.7	181.1	152.7	160.1	148.9	161.7
1939	183.7	160.6	186.7	225.8	186.8	186.9	195.3	235.2	189.2	150.1	202.6
1938	168.0	186.1	143.4	184.0	177.1	162.2	199.0	220.2	176.2	166.4	182.1
1937	152.2	169.1	138.4	200.7	144.6	152.7	206.2	192.4	161.7	146.4	171.8
1936	159.6	161.5	163.8	207.5	139.2	178.7	199.5	230.7	166.5	152.9	146.3
1935	151.6	131.5	160.4	202.9	160.8	137.8	195.0	174.3	155.7	142.9	146.1
1934	181.7	240.0	141.7	217.0	173.2	188.6	212.4	213.9	164.9	200.4	206.2
1933	157.4	159.8	155.3	203.3	145.8	168.4	190.6	188.7	150.4	131.0	140.5
1932	169.7	187.0	172.6	203.0	166.9	170.1	215.6	199.9	176.5	146.2	158.3
1931	198.1	202.5	152.4	246.1	189.9	188.7	237.4	267.8	222.8	203.6	212.8
1930	156.7	181.3	152.2	223.2	138.4	168.8	189.9	204.3	182.6	150.6	147.9
1929	190.6	193.3	177.4	196.9	165.1	191.1	206.6	282.6	180.7	149.5	178.3
1928	200.8	229.9	160.2	233.8	201.3	176.8	232.8	246.5	218.5	226.4	238.3
1927	194.6	231.5	171.3	226.0	182.8	179.6	220.4	236.7	206.8	188.5	237.7
1926	177.8	189.3	170.2	200.3	155.1	200.1	183.2	264.7	184.9	146.3	153.0
1925	215.5	229.3	179.4	256.2	219.8	196.0	230.6	259.8	213.6	195.0	231.5
1924	209.8	235.3	187.9	232.2	199.9	244.8	214.6	243.0	204.5	189.9	213.1
1923	205.5	231.5	160.4	222.9	231.7	183.4	221.3	247.4	222.6	202.4	230.2
1922	196.7	202.8	174.2	255.3	198.3	186.1	218.8	226.1	206.9	179.5	243.3
1921	200.7	211.8	173.1	217.1	195.2	168.8	169.5	305.1	191.7	154.0	229.8
1920	312.8	347.8	247.4	376.9	323.6	281.0	297.1	406.9	362.3	286.8	320.3

TABLE 18. - 1/DEATH RATES FOR TEN SELECTED CAUSES BY PREFECTURE: JAPAN 1920-1949 - Cont'd

5/PNEUMONIA, ALL FORMS (107-109) - Cont'd

Year	Hiroshima	Hokkaido	Hyogo	Ibarak	Ishikawa	Iwate	Kagawa	Kaposhima	Kanagawa	Kochi	Kumamoto	Kyoto
1949	55.0	79.5	55.6	62.4	77.4	103.1	61.6	69.9	69.4	54.3	68.8	54.1
1948	60.9	79.4	56.7	58.1	83.5	107.7	72.5	88.2	62.2	69.2	72.7	51.9
1947	127.6	141.6	116.4	112.1	148.1	160.6	130.7	163.6	126.6	125.1	154.2	116.8
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	140.1	156.2	125.1	147.8	190.0	196.5	129.9	172.1	142.7	115.1	159.9	100.4
1941	130.8	140.3	117.5	159.7	183.7	174.4	138.9	143.5	132.2	162.9	169.4	114.2
1940	147.5	157.6	131.1	168.6	217.1	215.1	163.1	170.8	128.4	132.5	174.2	142.3
1939	186.5	161.3	185.3	196.4	251.7	183.9	174.5	198.0	179.7	161.2	186.0	140.4
1938	163.8	150.5	140.4	185.7	208.4	177.9	154.8	188.9	157.9	185.2	180.8	119.4
1937	145.6	149.0	139.4	143.1	206.8	178.1	134.8	155.1	130.7	145.7	179.5	116.1
1936	187.9	133.5	141.1	161.6	231.3	226.9	193.5	172.6	118.2	173.5	178.8	154.4
1935	165.9	142.8	134.3	168.0	238.5	178.4	137.6	161.8	141.0	125.9	159.8	104.7
1934	181.0	163.8	155.1	188.7	249.9	187.8	219.6	190.0	162.0	156.3	202.4	143.6
1933	153.4	160.8	139.8	159.9	208.2	189.2	167.7	164.6	140.9	161.9	171.0	135.2
1932	211.1	148.7	158.7	159.2	243.6	185.1	190.5	176.9	151.6	177.0	189.3	140.3
1931	193.1	162.2	181.6	206.8	247.2	201.3	209.0	207.4	170.7	163.9	230.2	172.9
1930	162.0	174.1	137.3	151.6	188.9	175.1	174.9	172.5	125.3	138.3	171.5	132.9
1929	170.8	175.0	190.2	169.4	260.3	179.9	188.2	187.9	201.0	203.6	208.1	161.8
1928	204.6	199.7	171.4	231.5	244.9	222.0	210.0	201.1	215.1	182.4	202.0	160.1
1927	212.4	213.8	168.3	186.5	238.4	202.5	193.1	183.8	178.0	158.3	196.3	152.3
1926	194.9	172.9	165.5	145.7	241.2	185.0	208.6	203.9	147.1	184.8	219.3	145.5
1925	215.9	214.4	191.4	247.2	252.5	208.1	184.9	220.3	208.4	205.8	204.8	170.5
1924	264.0	183.6	202.6	195.0	247.5	231.6	224.7	220.5	212.5	169.1	247.6	152.3
1923	234.5	201.4	167.2	239.3	229.7	185.0	212.0	197.9	184.2	176.6	185.6	179.2
1922	192.3	173.1	175.9	211.9	225.0	210.7	221.7	223.9	208.5	165.3	231.3	126.2
1921	198.1	194.5	174.8	195.7	195.6	165.6	221.4	223.9	208.5	138.0	230.5	154.0
1920	335.3	330.0	358.2	279.8	302.1	236.7	261.1	262.7	287.6	265.3	279.3	255.6

TABLE 18. 1/DEATH RATES FOR TEN SELECTED CAUSES BY PREFECTURE: JAPAN 1920-1949 - Cont'd
5/PNEUMONIA, ALL FORMS (107-109) - Cont'd

Year	Mie	Miyagi	Miyazaki	Nagano	Nagasaki	Nara	Miigata	Oita	Okayama	Osaka	Saga	Saitama
1949	59.7	76.4	62.9	59.7	78.9	61.3	75.3	73.3	51.8	75.8	58.9	71.9
1948	62.9	76.4	66.8	57.8	75.1	51.7	76.2	68.4	63.9	58.7	59.2	61.9
1947	119.3	140.3	137.8	116.7	144.7	123.4	116.1	146.7	128.6	156.0	132.8	127.5
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	132.9	167.3	152.9	117.3	170.2	117.3	148.8	178.0	110.8	136.4	167.7	157.3
1941	143.7	153.0	139.1	125.2	155.7	121.5	147.1	172.9	111.3	134.3	182.5	163.8
1940	142.0	171.1	130.1	129.3	188.4	130.2	171.6	178.2	126.5	157.0	165.1	178.6
1939	185.7	175.5	209.0	153.9	206.6	154.8	178.9	214.6	169.5	215.9	220.0	209.2
1938	207.9	153.9	172.6	148.7	227.0	128.5	154.2	195.2	134.1	155.3	204.8	171.5
1937	161.4	154.3	151.0	121.5	186.9	146.1	154.3	218.1	120.8	160.8	193.6	164.6
1936	189.6	181.8	173.5	149.4	178.8	147.9	152.9	205.0	155.6	144.0	208.1	148.4
1935	163.1	178.7	171.5	133.7	157.6	153.6	167.8	165.8	137.7	154.9	150.1	174.6
1934	205.1	168.0	167.9	139.5	184.8	196.2	176.1	209.5	172.3	169.5	208.0	204.9
1933	137.0	185.2	175.3	106.5	183.7	137.2	149.6	182.7	159.1	162.8	181.8	163.9
1932	163.4	166.8	195.4	128.1	220.1	168.5	175.8	229.1	153.6	164.8	203.1	181.8
1931	226.7	187.1	184.2	158.0	203.3	162.7	192.4	237.0	154.1	200.1	215.3	231.6
1930	273.6	189.7	144.3	186.2	194.3	120.4	168.7	178.5	134.0	135.9	208.5	143.2
1929	190.3	173.2	248.3	137.6	220.9	177.7	180.0	284.4	159.4	226.1	239.7	198.4
1928	224.0	224.0	180.8	177.2	222.5	151.9	212.2	216.1	157.6	174.4	214.9	237.4
1927	196.9	217.1	153.6	157.5	192.9	203.5	191.4	227.3	163.9	227.3	193.8	221.3
1926	165.9	192.4	196.1	125.2	203.3	133.2	175.6	242.1	173.2	171.4	233.7	162.9
1925	274.0	222.2	200.7	133.8	201.2	206.2	204.5	205.9	152.4	249.6	227.2	233.6
1924	206.0	253.0	191.3	176.7	235.9	185.4	224.7	243.9	182.0	210.7	237.8	214.9
1923	186.2	241.1	163.5	161.3	182.0	176.2	197.6	234.3	167.3	232.9	244.9	237.9
1922	197.1	244.7	189.9	175.3	185.1	151.2	225.1	202.7	150.4	196.1	234.7	221.3
1921	133.8	212.2	180.2	145.6	238.1	170.8	199.6	236.5	155.0	243.4	258.3	221.5
1920	281.3	397.4	218.4	240.7	322.7	296.3	319.5	293.9	212.4	448.6	301.8	354.7

TABLE 18. - 1/DEATH RATES FOR TEN SELECTED CAUSES BY PREFECTURE: JAPAN 1920-1949 - Cont'd
5/PNEUMONIA, ALL FORMS (107-109) - Cont'd

Year	Shira	Shimane	Shizuoka	Tochigi	Tokushima	Tokyo	Tottori	Toyama	Yakayama	Yamagata	Yamaguchi	Yamanashi
1949	58.4	62.3	61.5	70.0	69.6	65.9	50.5	79.6	56.0	77.4	68.5	64.1
1948	58.5	74.6	57.4	56.4	86.2	57.3	59.9	77.3	59.3	72.3	67.0	54.8
1947	129.2	132.5	111.0	125.4	153.7	129.7	105.5	124.4	113.6	126.2	121.8	129.1
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	114.7	150.9	120.7	152.7	149.8	135.3	121.0	208.1	116.1	132.6	150.7	147.4
1941	124.4	143.9	118.8	146.3	153.8	124.9	132.1	195.0	98.6	143.2	129.9	132.9
1940	135.9	153.7	126.5	156.3	178.8	124.8	134.0	234.3	123.1	146.6	140.2	130.8
1939	149.6	227.5	137.7	198.2	229.0	151.6	152.1	250.0	158.0	167.3	198.4	160.2
1938	154.3	202.1	141.6	155.3	199.4	147.0	120.6	225.0	117.2	145.5	179.0	168.3
1937	134.1	147.9	121.3	144.3	193.0	121.5	129.4	211.5	110.0	152.9	146.3	135.0
1936	146.0	183.0	131.4	149.4	209.0	107.4	129.9	209.3	149.1	154.8	193.9	124.0
1935	121.7	162.8	139.9	153.6	176.9	117.8	124.2	222.2	135.4	174.2	164.3	169.3
1934	174.5	190.5	169.0	186.5	246.2	157.5	157.1	269.2	150.4	178.7	164.7	163.1
1933	131.3	156.8	140.8	174.2	206.7	138.2	122.7	214.0	111.5	156.2	158.3	141.4
1932	141.3	192.3	145.6	164.2	230.4	129.0	117.0	228.9	158.7	173.1	162.5	144.1
1931	211.2	223.7	168.1	222.7	264.0	180.1	159.9	254.8	158.7	179.5	197.4	178.3
1930	119.9	169.0	147.3	146.0	183.9	116.1	119.8	210.3	112.4	183.7	176.3	144.2
1929	171.1	206.0	149.9	172.7	227.3	169.5	159.7	270.8	162.8	175.0	188.0	157.2
1928	155.0	187.4	169.2	215.8	243.1	185.8	118.8	263.6	136.8	184.1	184.4	213.5
1927	161.6	198.7	155.8	213.1	235.0	175.3	136.1	254.2	146.9	190.5	180.2	175.7
1926	157.0	182.6	136.7	151.0	231.1	140.0	144.0	201.7	133.4	188.5	214.2	149.7
1925	186.5	186.6	219.8	242.9	262.7	211.4	173.9	242.9	189.2	208.0	188.0	202.6
1924	166.2	236.0	180.9	190.4	268.8	171.9	164.1	275.2	161.2	202.3	245.5	221.4
1923	205.6	193.6	157.9	221.9	275.2	207.8	149.6	309.0	134.3	207.2	185.8	176.4
1922	160.2	187.1	171.2	205.8	253.0	176.1	141.1	285.9	140.2	230.1	182.6	214.5
1921	140.5	156.3	171.9	206.7	236.0	204.3	129.9	232.4	156.0	171.4	218.7	132.5
1920	207.1	232.0	270.7	296.0	413.2	298.7	240.4	347.1	196.8	338.1	294.3	332.5

See footnotes at end of table.

TABLE 18. 1/DEATH RATE FOR TEN SELECTED CAUSES BY PREFECTURE: JAPAN 1920-1949 - Cont'd
2/DIARRHEA, ENTERITIS AND ULCERATION OF INTESTINES (All ages) (117-120)

Year	All Japan	Aichi	Akita	Aomori	Chiba	Ehime	Fukui	Fukuoka	Fukushima	Gifu	Gumma
1949	87.8	71.9	125.8	144.0	64.8	79.3	144.6	85.3	84.6	101.4	75.6
1948	103.8	76.8	182.7	206.3	82.8	84.4	198.6	78.4	105.0	112.7	102.2
1947	130.1	107.1	190.1	196.1	104.0	129.8	211.9	120.0	113.7	139.8	115.7
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	142.4	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	130.7	96.2	201.1	252.9	125.5	118.8	233.2	101.3	159.6	154.5	149.2
1941	131.7	129.9	200.2	241.1	136.8	119.3	256.8	90.3	156.8	174.5	158.9
1940	149.4	151.1	228.0	267.3	154.0	132.0	297.2	108.1	196.6	192.5	216.7
1939	156.9	166.7	224.3	279.2	147.7	147.7	272.8	118.7	189.5	195.9	208.6
1938	163.5	168.2	202.7	284.9	163.5	135.0	316.7	134.0	195.7	225.4	239.1
1937	168.3	174.2	263.2	295.1	153.3	143.4	332.3	135.4	218.4	209.0	225.3
1936	171.5	167.5	239.8	355.6	159.8	133.2	375.9	149.2	192.2	223.5	211.8
1935	161.1	152.0	232.0	281.6	152.4	125.4	281.0	131.5	197.3	207.1	186.2
1934	186.6	201.8	276.1	375.2	159.8	141.3	417.9	133.6	230.5	256.8	219.8
1933	193.4	178.7	276.8	303.8	194.9	179.4	354.0	158.1	227.9	229.3	219.6
1932	206.3	191.2	348.9	494.7	198.3	174.9	386.6	169.8	260.2	248.5	232.9
1931	213.6	207.7	320.2	372.3	203.8	173.8	385.9	203.6	250.9	270.5	251.1
1930	220.7	211.7	325.8	455.4	212.4	184.8	381.6	200.9	259.5	271.7	246.7
1929	246.4	247.6	350.5	446.9	250.9	194.0	445.3	235.9	302.2	292.1	311.7
1928	232.4	256.3	363.6	474.7	232.7	184.9	406.2	216.8	266.8	299.5	285.3
1927	242.5	250.4	344.9	429.5	220.3	215.1	448.1	202.9	286.9	284.7	332.6
1926	231.7	231.6	350.3	410.9	222.9	200.3	396.7	193.8	245.8	274.2	298.0
1925	237.2	252.6	741.0	431.3	208.4	184.2	398.6	189.9	243.9	293.7	336.0
1924	257.5	282.4	367.0	514.3	247.9	221.5	405.0	207.7	275.3	314.3	328.8
1923	284.1	308.3	427.5	505.0	306.9	221.2	461.5	210.7	299.6	330.3	382.5
1922	267.6	258.4	378.2	517.0	343.4	225.4	453.0	198.3	274.3	297.3	334.5
1921	273.0	324.4	349.0	457.3	255.6	250.1	451.4	212.8	272.3	336.5	326.3
1920	253.5	299.4	262.9	436.2	235.2	225.0	420.1	213.1	245.2	304.8	330.8

TABLE 18. - 1/DEATH RATE FOR TEN SELECTED CAUSES BY PREFECTURE: JAPAN 1920-1949 - Cont'd
G/MIAR-ITA, ENTERITIS AND ULCERATION OF INTESTINES (all ages) (119-120)

Year	Firoshima	Hokkaido	Hyogo	Ibaraki	Ishikawa	Iwate	Karawa	Kasoshima	Kanagawa	Kochi	Kumamoto	Kyoto
1949	79.4	106.4	97.9	80.2	120.9	119.3	73.5	100.3	46.1	72.0	92.0	70.1
1948	98.4	130.5	103.3	87.2	172.7	162.1	84.7	118.4	60.9	91.5	102.1	85.1
1947	132.6	145.7	154.6	NA	174.9	137.7	98.7	142.9	74.2	117.2	124.5	135.8
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	115.8	150.7	111.1	134.9	196.7	212.0	122.6	212.8	82.3	113.4	134.3	105.4
1941	108.1	164.0	118.7	144.0	219.1	208.2	107.5	161.8	83.7	103.4	127.0	97.6
1940	124.7	176.3	121.9	164.5	217.1	263.9	116.2	208.7	91.7	107.2	130.4	93.7
1939	156.2	208.0	137.5	171.6	206.8	247.6	150.0	183.8	98.4	114.7	136.9	102.6
1938	145.6	193.3	151.0	191.5	231.1	275.9	148.6	181.3	103.1	124.9	157.4	113.9
1937	127.4	213.8	147.4	175.8	237.0	310.0	129.8	198.9	113.2	119.5	152.8	104.5
1936	152.7	213.7	154.6	170.3	243.2	245.4	141.1	189.0	108.7	120.6	149.8	132.8
1935	130.3	207.0	148.4	169.9	222.8	234.8	126.8	189.3	119.5	142.2	148.7	122.1
1934	158.7	202.0	171.1	183.9	315.6	283.3	161.5	197.3	136.9	132.5	159.8	138.3
1933	171.9	235.0	186.1	178.2	266.1	284.7	176.8	189.7	149.7	160.0	160.1	142.0
1932	170.2	260.8	192.6	192.7	288.1	366.7	173.1	200.0	155.1	167.9	179.1	163.2
1931	167.8	240.7	187.3	214.0	277.5	286.5	173.5	233.6	158.4	158.4	199.1	183.0
1930	168.0	267.7	204.9	218.9	270.1	385.1	165.5	227.3	170.2	188.7	208.9	173.9
1929	191.3	292.3	217.9	275.3	303.2	306.5	196.9	234.7	217.3	198.8	213.7	172.4
1928	192.7	279.4	212.3	287.3	271.6	366.6	169.4	216.6	195.9	201.9	195.2	160.3
1927	217.1	296.6	222.7	263.5	307.5	312.4	186.1	203.1	238.5	232.4	190.5	192.5
1926	205.8	218.5	213.2	244.0	267.3	301.1	175.3	240.0	205.9	222.9	171.6	200.5
1925	205.0	281.3	227.2	255.4	277.6	285.7	172.1	215.3	232.3	188.1	197.4	205.7
1924	239.8	271.4	225.4	279.6	269.4	369.5	248.3	213.3	237.5	216.3	205.6	210.2
1923	235.8	246.7	263.4	291.7	320.2	347.4	227.5	247.4	319.8	264.1	226.1	242.8
1922	236.0	288.2	267.3	285.4	377.6	377.6	218.8	332.6	248.2	220.8	231.2	232.5
1921	225.7	244.8	254.5	271.5	290.3	352.4	216.5	204.1	279.3	232.2	238.5	266.5
1920	216.9	264.4	255.1	271.6	249.3	321.6	205.8	163.7	242.8	216.4	203.6	223.4

TABLE 18. - 1/DEATH RATES FOR TEN SELECTED CAUSES BY PREFECTURE: JAPAN 1920-1949 - Cont'd
 6/DIARRHEA, ENTERITIS AND ULCERATION OF INTESTINES (all ages) (119-120) - Cont'd

Year	Mie	Miyagi	Miyazaki	Nagano	Nagasaki	Nara	Niigata	Oita	Okayama	Osaka	Saga	Saitama
1949	97.5	76.2	93.6	65.8	88.4	101.8	112.3	103.7	87.2	81.5	97.9	90.5
1948	111.1	103.2	104.4	90.5	99.9	130.1	140.7	113.7	109.0	84.0	114.9	105.3
1947	162.9	96.7	130.6	114.4	129.6	156.6	154.7	156.3	159.3	120.6	152.7	133.2
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	162.5	138.9	148.4	155.2	127.6	147.5	200.0	154.2	135.2	86.4	127.6	164.9
1941	171.2	123.9	127.5	144.2	110.4	163.0	189.3	142.2	130.7	89.5	134.3	204.1
1940	166.5	138.0	152.2	162.1	118.2	155.6	244.1	157.1	149.5	86.6	134.3	259.1
1939	176.6	178.4	177.4	169.6	133.8	171.6	257.9	151.6	169.7	92.1	147.8	242.5
1938	208.3	180.1	157.6	163.4	138.6	184.6	258.6	151.5	179.3	87.2	141.2	264.0
1937	182.5	205.7	170.5	172.4	164.6	180.0	266.7	184.1	167.1	96.5	162.2	240.3
1936	197.6	186.1	149.0	166.9	166.0	171.9	279.9	159.6	166.4	91.3	204.9	280.7
1935	177.0	181.7	163.0	148.2	142.9	177.6	217.3	156.6	155.2	109.0	166.6	210.2
1934	227.8	202.2	156.6	173.1	137.6	222.9	303.7	165.1	161.0	173.3	173.7	249.6
1933	221.5	213.3	180.8	175.3	147.3	226.6	282.4	173.7	208.6	138.2	194.0	256.0
1932	215.4	268.4	194.8	181.4	170.6	230.8	281.8	193.4	179.8	127.7	235.9	280.4
1931	243.1	264.5	212.9	179.1	188.0	232.7	317.8	205.5	186.0	151.6	253.4	310.2
1930	235.4	281.9	186.1	223.5	174.2	209.0	293.3	190.1	185.2	155.9	251.9	281.7
1929	261.8	334.1	208.0	224.8	195.5	245.6	362.1	201.6	225.7	161.8	249.5	334.6
1928	273.0	290.9	183.9	206.4	172.0	213.3	331.4	186.4	206.2	165.4	201.4	329.8
1927	251.9	296.2	219.3	221.1	177.1	224.0	317.2	203.8	228.2	185.3	232.3	321.8
1926	243.6	272.2	176.2	222.9	133.4	229.3	341.0	182.8	202.9	197.8	225.3	287.7
1925	265.8	254.8	166.3	219.3	138.2	243.2	334.1	175.8	180.3	202.0	202.8	322.1
1924	272.5	308.6	187.5	242.2	156.8	239.6	370.4	194.4	222.5	236.5	235.6	342.7
1923	305.0	317.2	222.3	243.1	164.0	255.6	387.9	194.7	240.7	247.4	237.6	333.6
1922	259.5	285.8	207.9	253.5	161.7	258.8	389.6	169.6	219.9	290.1	221.9	323.1
1921	323.3	278.4	187.9	277.9	165.4	284.5	388.5	211.7	233.9	257.6	241.7	364.9
1920	296.8	251.0	170.2	253.7	147.3	278.6	319.5	158.2	222.1	238.2	212.5	328.8

TABLE 18 - $\frac{1}{2}$ DEATH RATES FOR TEN SELECTED CAUSES BY PREFECTURE: JAPAN 1920-1949 - Cont'd
 $\frac{6}{2}$ DIARRHEA, ENTERITIS AND ULCERATION OF INTESTINES (all ages) (119-120) - Cont'd

Year	Shiga	Shimane	Shizuoka	Tochigi	Tokushima	Tokyo	Tottori	Torima	Wakayama	Yamagata	Yamaguchi	Yamanashi
1949	134.6	101.4	74.2	101.9	95.8	40.4	100.1	171.7	98.4	125.2	79.3	88.1
1948	145.2	110.1	85.5	107.4	108.7	48.9	113.5	185.9	97.0	149.6	85.0	105.0
1947	225.7	128.4	106.1	130.8	148.4	64.0	147.6	246.7	133.9	158.1	133.6	133.2
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	194.3	147.0	113.5	178.0	124.8	49.1	149.7	270.5	144.1	218.5	113.2	150.6
1941	205.9	137.3	113.7	185.1	120.3	43.1	129.9	246.1	128.1	215.7	96.6	188.2
1940	212.1	149.9	121.6	228.8	128.2	49.4	136.9	308.3	143.0	258.5	103.8	195.9
1939	165.2	190.0	124.7	235.8	135.9	51.2	169.9	304.1	146.6	272.0	120.9	195.5
1938	230.2	156.7	139.1	256.3	140.9	55.2	160.1	329.4	157.3	264.7	129.4	205.7
1937	224.4	151.0	149.7	232.4	167.8	59.5	172.9	348.6	141.5	281.8	117.5	216.8
1936	220.0	161.5	175.5	244.4	121.7	78.8	178.1	296.2	148.7	307.1	126.3	214.0
1935	209.2	153.5	146.7	239.3	150.3	66.4	169.6	294.0	198.4	265.4	129.2	214.2
1934	253.3	175.9	161.6	285.3	157.9	77.4	187.9	362.0	172.3	328.9	132.3	200.7
1933	246.9	185.6	184.6	285.6	170.8	102.5	190.0	307.4	201.8	339.5	138.4	257.7
1932	249.8	224.0	183.7	251.0	179.8	88.4	236.2	329.2	191.9	338.5	156.8	226.3
1931	280.4	197.5	186.9	299.1	190.9	103.9	213.7	362.9	190.4	330.1	165.3	233.1
1930	270.8	221.4	220.7	276.1	166.8	106.8	255.7	313.0	206.4	355.8	171.8	229.6
1929	281.6	259.6	222.6	323.1	206.4	142.6	249.1	367.3	202.5	389.6	184.3	272.1
1928	279.9	211.2	219.7	300.9	188.9	136.5	233.2	325.3	210.0	367.7	161.6	233.1
1927	308.1	207.5	220.9	302.9	216.6	137.6	230.2	329.2	210.9	376.3	190.3	266.2
1926	290.1	208.9	231.3	294.5	206.3	148.7	236.3	340.9	218.6	399.4	174.5	251.5
1925	313.7	205.6	244.4	297.8	174.8	153.5	212.8	333.3	217.4	336.4	166.4	330.1
1924	300.4	221.8	231.7	300.2	237.8	199.6	246.5	339.1	237.5	373.4	201.2	301.5
1923	309.1	241.8	240.4	304.4	254.8	224.7	277.4	395.5	267.2	418.9	215.3	312.0
1922	319.8	190.7	230.2	325.1	259.4	229.6	246.2	350.8	286.1	402.0	183.0	333.8
1921	405.7	189.4	269.8	335.0	252.5	202.8	249.0	308.9	268.5	388.9	185.0	353.1
1920	373.6	190.8	247.8	311.7	247.5	207.4	263.3	263.2	246.5	350.3	184.5	333.2

See footnotes at end of table.

TABLE 18. - 1/DEATH RATES FOR TEN SELECTED CAUSES BY PREFECTURE: JAPAN 1920-1949 - Cont'd
7/NEPHRITIS, ALL FORMS (130-132)

Year	All Japan	Alchi	Akita	Aomori	Chiba	Ehime	Fukui	Fukuoka	Fukushima	Gifu	Gumma
1949	41.2	30.6	51.2	30.7	49.1	37.0	36.7	43.5	35.5	33.3	41.5
1948	45.6	36.8	58.7	38.6	55.2	38.4	41.3	46.5	41.5	36.7	44.6
1947	57.7	46.9	63.9	40.2	63.0	49.4	43.8	70.0	49.4	48.1	54.2
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	66.9	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	69.1	51.9	94.4	65.8	76.5	54.4	63.4	74.5	64.9	60.0	69.3
1941	68.1	49.1	94.7	63.4	79.0	55.4	65.7	69.0	60.5	63.5	64.9
1940	75.7	56.3	119.5	74.9	81.8	64.7	77.9	76.3	63.6	62.8	69.6
1939	82.2	58.3	102.3	74.4	84.8	68.9	85.1	94.9	66.4	69.8	74.1
1938	87.2	63.2	91.3	72.1	102.2	71.3	101.1	112.3	73.7	72.1	77.7
1937	79.3	57.5	90.2	64.4	88.5	67.6	89.3	104.0	66.3	77.7	74.1
1936	82.2	60.1	95.0	96.7	90.4	67.4	85.9	101.1	68.2	70.9	73.8
1935	80.8	60.7	93.4	64.1	87.6	74.2	89.4	96.0	64.3	69.5	78.1
1934	87.2	65.6	95.1	75.8	93.9	73.4	108.2	92.8	76.0	72.7	81.0
1933	80.1	67.3	98.7	66.3	105.0	73.0	91.8	99.3	71.6	75.4	81.5
1932	92.6	69.2	108.3	78.8	107.5	77.5	96.8	107.2	77.9	76.0	87.9
1931	88.4	68.4	110.1	73.6	113.7	78.2	113.9	117.6	77.1	84.6	89.2
1930	88.5	68.8	117.3	91.1	114.0	77.8	116.2	117.3	82.0	78.6	85.7
1929	103.2	74.9	166.5	118.0	121.2	76.1	98.9	114.3	80.7	79.0	90.6
1928	107.5	76.6	113.7	88.0	131.6	77.6	97.0	114.3	82.9	80.8	86.0
1927	87.5	69.3	135.6	85.1	106.2	77.8	98.2	105.2	79.5	71.7	87.0
1926	98.2	72.1	153.8	95.7	110.4	80.5	90.7	109.0	75.8	75.4	81.8
1925	107.3	70.3	140.9	86.7	120.8	77.2	93.3	111.0	78.3	74.8	89.2
1924	105.8	74.4	131.8	85.4	124.9	90.0	90.2	104.6	86.7	78.6	100.7
1923	107.7	76.5	203.9	114.6	133.5	93.5	96.4	103.7	85.9	82.5	97.2
1922	106.4	74.1	158.8	122.1	120.3	88.4	94.5	103.2	93.1	78.5	94.6
1921	105.7	80.4	121.9	90.5	121.0	85.2	87.7	110.4	91.8	83.6	91.9
1920	99.2	73.0	105.1	78.5	107.3	78.1	93.1	118.1	88.2	72.8	85.8

TABLE 18. - 1/DEATH RATES FOR TEN SELECTED CAUSES BY PREFECTURE: JAPAN 1920-1949 - Cont'd

7/NEPHRITIS, ALL FORMS (130-132) - Cont'd

Year	Firoshima	Hokkaido	Hyoogo	Ibaraki	Ishikawa	Iwate	Yamaguchi	Kagoshima	Kanazawa	Kochi	Kumamoto	Kyoto
1949	38.7	27.1	35.8	51.3	38.2	38.1	43.8	57.8	32.5	48.0	47.6	30.8
1948	45.1	28.5	42.3	56.1	45.9	44.2	47.0	57.5	37.8	47.6	60.0	36.3
1947	58.5	31.6	56.5	65.3	56.7	53.2	61.3	78.4	49.6	62.6	74.6	51.1
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	65.2	46.1	61.7	82.5	68.0	62.4	79.6	106.8	62.4	71.6	90.5	66.2
1941	59.9	30.6	59.6	80.8	59.6	67.3	68.4	101.1	61.8	101.1	84.2	64.7
1940	68.9	55.6	67.2	89.1	73.5	80.9	76.4	108.3	65.5	71.8	92.8	72.0
1939	76.4	64.0	78.0	92.8	82.6	76.1	85.5	106.0	81.8	81.7	104.8	72.4
1938	75.4	57.3	80.4	95.9	88.8	76.9	90.8	110.1	91.0	84.9	109.1	80.6
1937	66.4	57.4	74.3	88.3	85.9	72.5	75.6	95.4	80.6	76.6	94.9	69.9
1936	74.7	62.4	77.8	90.4	82.8	73.8	81.7	100.0	83.2	83.2	105.0	80.9
1935	74.3	56.7	74.5	85.6	80.2	76.5	79.5	99.7	83.2	89.4	96.4	74.3
1934	79.4	60.4	78.9	97.3	99.5	83.1	80.8	98.4	90.0	80.2	101.3	81.5
1933	79.7	64.3	84.7	99.3	86.6	83.1	81.9	96.5	101.2	83.9	104.3	81.4
1932	84.9	63.2	80.5	103.0	92.3	85.1	74.9	101.4	105.3	87.5	106.1	88.9
1931	91.8	71.3	86.6	105.5	91.3	89.5	87.9	101.9	98.7	88.9	114.2	95.8
1930	93.7	66.1	82.7	107.1	990.6	100.1	87.1	105.8	103.2	95.2	107.2	94.3
1929	88.5	87.9	82.4	110.1	93.6	105.6	87.3	104.3	119.2	100.4	107.3	95.9
1928	89.2	65.0	88.0	107.8	88.9	93.5	77.3	97.0	115.2	107.4	107.7	103.6
1927	89.7	75.8	84.4	102.7	91.0	94.0	90.4	97.9	106.8	92.1	104.6	103.8
1926	92.6	69.4	81.0	103.1	84.3	96.3	84.4	99.3	108.1	86.9	110.2	96.1
1925	93.2	68.4	91.6	117.1	81.1	110.2	89.1	97.4	118.2	78.7	99.6	107.3
1924	103.0	66.3	98.2	115.0	83.7	120.5	94.6	98.5	125.8	91.8	101.9	115.8
1923	101.2	67.0	91.5	106.4	100.4	130.4	91.5	99.7	113.4	92.2	105.7	115.0
1922	96.0	64.2	92.2	102.9	95.9	133.0	89.1	106.8	124.1	100.9	117.5	107.7
1921	94.9	69.1	92.8	102.9	88.6	112.6	84.9	106.0	122.2	106.1	109.0	117.9
1920	85.7	68.5	86.8	101.1	74.7	109.0	88.1	105.3	109.3	96.7	101.5	106.5

TABLE 18. - $\frac{1}{\text{DEATH RATES FOR TEN SELECTED CAUSES BY PREFECTURE: JAPAN 1920-1949}}$ - Cont'd $\frac{7}{\text{NEPHRITIS, ALL FORMS (130-132)}} - \text{Cont'd}$

Year	Mie	Miyagi	Miyazaki	Nagano	Nagasaki	Nara	Niigata	Oita	Okayama	Osaka	Saga	Saitama
1949	41.5	37.8	57.7	44.4	43.4	49.4	46.3	51.8	39.0	43.9	60.7	46.5
1948	43.3	39.1	53.5	52.4	46.5	51.4	54.4	59.0	42.7	46.9	67.7	50.0
1947	56.1	48.8	70.5	57.0	60.4	67.7	63.3	67.2	53.0	67.3	80.3	63.6
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	65.6	77.6	87.2	77.4	72.2	75.8	77.1	87.8	52.7	65.6	88.3	88.2
1941	68.1	71.7	81.3	75.1	75.4	81.1	77.3	80.3	55.1	60.9	97.0	83.4
1940	76.0	71.9	90.4	76.0	79.3	80.2	88.1	94.3	61.7	73.2	106.4	97.5
1939	78.7	78.1	96.7	87.7	88.7	94.0	93.3	92.4	66.3	78.7	109.4	96.8
1938	89.6	83.2	97.3	83.6	93.7	103.5	98.0	97.4	71.3	82.9	124.2	102.7
1937	75.7	86.1	86.3	77.7	88.1	91.2	87.0	90.9	59.6	80.3	125.6	89.5
1936	85.0	80.4	93.1	79.3	86.5	91.8	90.3	92.3	62.9	84.3	116.7	95.6
1935	85.2	84.5	93.9	84.4	89.6	95.9	87.2	90.2	64.5	81.7	106.4	92.2
1934	89.5	86.0	102.6	90.8	85.7	102.8	112.2	92.9	64.5	93.5	100.9	113.2
1933	88.0	91.5	96.1	88.9	88.7	99.2	99.7	97.2	71.4	98.1	123.4	116.0
1932	90.3	93.1	113.0	92.1	90.9	106.5	98.9	97.8	64.7	103.0	127.6	115.6
1931	108.7	106.6	115.6	97.4	94.1	105.7	112.0	93.3	71.6	107.0	122.8	134.2
1930	93.4	124.0	116.0	109.0	92.2	101.0	113.1	100.2	69.9	107.5	127.8	123.2
1929	100.5	123.1	113.8	108.5	92.1	103.1	125.6	99.4	77.5	106.2	111.8	124.6
1928	109.1	117.0	115.3	99.7	92.7	107.6	108.3	92.8	77.6	110.3	122.1	126.4
1927	90.9	112.7	112.4	95.8	90.6	105.2	112.3	93.8	73.4	105.0	118.4	117.1
1926	85.2	118.9	104.1	100.7	85.8	96.6	123.2	91.1	70.3	105.3	113.1	115.2
1925	103.1	129.9	95.6	105.4	82.0	103.1	113.1	83.9	76.0	112.7	110.2	125.4
1924	97.2	140.9	107.8	109.5	81.5	100.6	132.6	96.2	80.4	122.2	117.0	133.0
1923	102.9	155.6	110.8	110.8	87.7	108.3	126.3	89.1	75.2	114.5	119.4	120.7
1922	89.3	148.3	104.6	119.6	84.2	94.7	126.2	93.4	77.7	121.8	137.1	118.6
1921	107.8	132.8	110.3	108.0	89.1	123.1	114.2	95.2	74.7	122.8	123.4	123.3
1920	101.8	129.4	100.3	106.9	81.7	112.1	112.8	90.1	68.2	119.1	120.8	106.0

TABLE 18. - 1/DEATH RATES FOR TEN SELECTED CAUSES BY PREFECTURE: JAPAN 1920-1949 - Cont'd

7/NEPHRITIS, ALL FORMS (130-132) - Cont'd

Year	Shiga	Shimane	Shizuoka	Tochigi	Tokushima	Tokyo	Fukushima	Tochigi	Tokyo	Yamanashi	Yamanashi
1949	47.5	46.9	35.6	42.2	44.6	37.2	61.6	43.9	26.6	48.3	48.3
1943	57.7	49.8	39.7	45.6	49.3	38.5	59.7	50.8	31.8	57.9	56.3
1947	67.5	57.0	52.0	54.6	57.0	46.5	79.2	66.2	45.4	71.7	69.1
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	82.2	78.5	61.3	75.9	76.0	57.4	110.0	82.2	48.6	84.9	81.5
1941	84.7	87.9	61.3	66.3	73.6	58.4	106.7	83.3	53.3	93.9	83.7
1940	92.2	100.9	67.6	74.1	83.5	66.1	115.6	92.3	58.8	105.4	85.4
1939	90.7	108.7	75.4	77.4	95.0	76.2	133.6	104.6	63.5	100.8	101.4
1938	105.3	115.4	82.1	80.0	92.0	83.1	131.6	111.1	65.5	99.8	111.4
1937	96.9	97.5	73.7	72.6	87.1	74.1	112.0	100.3	58.1	91.7	95.8
1936	95.3	103.2	72.1	70.9	88.3	73.8	108.7	99.5	61.6	91.5	108.1
1935	93.3	98.2	72.6	67.4	96.2	74.9	111.7	97.4	66.1	99.1	98.3
1934	111.3	105.3	75.9	77.4	98.9	81.0	106.6	122.1	65.3	111.7	101.2
1933	109.8	114.7	78.6	86.1	92.5	87.8	103.3	102.8	59.8	105.1	103.0
1932	106.7	111.5	87.5	84.1	100.6	92.8	113.2	112.9	68.9	112.6	105.2
1931	106.4	120.4	84.4	91.9	115.0	102.8	121.6	119.0	79.6	111.5	119.6
1930	107.1	124.5	88.9	86.9	108.2	101.1	117.5	112.5	69.8	115.8	114.8
1929	109.1	119.5	87.9	90.4	117.8	115.9	115.5	118.8	67.5	135.4	112.5
1928	107.9	110.2	85.5	88.6	117.3	114.6	114.1	109.1	74.8	124.4	109.7
1927	107.9	115.7	80.5	82.5	109.3	116.2	119.2	119.8	67.1	119.4	120.0
1926	105.5	111.9	84.4	85.8	103.5	122.4	108.6	118.7	72.8	114.1	108.0
1925	117.0	115.3	89.0	85.2	113.5	118.7	108.4	106.1	76.7	113.4	110.7
1924	118.2	120.2	96.1	95.7	120.8	134.2	118.7	116.8	73.9	117.6	126.7
1923	125.3	135.8	100.1	92.6	113.4	137.1	125.2	126.1	78.6	120.6	133.9
1922	108.6	109.4	93.3	93.2	107.8	140.2	107.7	101.9	74.6	110.0	115.5
1921	124.8	113.8	100.5	90.0	107.7	146.2	107.7	109.0	76.3	98.0	108.5
1920	114.1	102.7	84.0	85.1	105.8	139.7	80.3	103.1	73.7	106.9	98.0

See footnotes at end of table.

TABLE 13. - $\frac{1}{\text{DEATH RATES FOR TEN SELECTED CAUSES BY PREFECTURE: JAPAN 1920-1949}}$ - Cont'd
CONGENITAL DEBILITY (158)

Year	All Japan	Aichi	Arita	Aomori	Chiba	Ehime	Fukui	Fukuoka	Fukushima	Yifu	Gumma
1949	45.4	50.0	86.7	89.2	41.1	44.4	65.9	45.3	45.4	59.0	36.8
1948	47.6	57.2	83.0	90.2	46.4	48.5	75.4	42.1	52.1	62.1	42.9
1947	51.8	58.5	97.0	90.2	52.0	46.7	73.8	52.6	56.9	66.0	45.5
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	69.9	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	71.1	68.9	135.8	122.5	73.2	72.9	107.8	23.3	83.0	96.2	63.5
1941	72.1	70.9	127.9	118.5	76.1	71.2	114.5	66.6	81.9	95.1	67.8
1940	76.8	76.8	153.0	139.9	82.5	73.8	126.5	88.7	87.3	108.1	71.6
1939	80.5	72.9	149.4	132.1	86.6	78.7	102.7	78.1	97.9	112.0	69.3
1938	85.8	83.4	159.9	151.5	87.0	74.8	130.3	87.6	103.4	117.0	74.6
1937	90.0	85.7	164.9	145.4	98.5	86.8	127.1	91.4	105.8	121.7	75.8
1936	97.7	97.9	175.9	179.8	103.5	98.1	152.1	97.6	111.2	137.8	77.2
1935	93.7	91.2	177.1	164.4	101.1	89.1	131.9	83.9	112.5	127.3	77.6
1934	101.3	103.3	180.2	174.4	99.8	99.6	175.0	94.9	129.2	149.7	85.4
1933	101.0	96.2	178.7	178.2	105.8	98.1	154.9	101.3	118.2	139.0	77.0
1932	90.2	89.9	156.6	156.2	100.1	96.1	122.1	92.2	111.3	125.4	70.9
1931	94.8	96.6	173.3	166.3	107.9	90.0	149.6	88.3	116.2	122.9	76.6
1930	92.8	89.9	168.8	173.2	107.5	90.2	173.5	86.2	110.1	124.6	79.8
1929	124.0	100.2	186.5	165.8	108.4	97.2	155.4	97.0	123.0	134.6	83.8
1928	102.9	105.0	181.2	137.2	117.0	86.7	147.5	96.9	117.5	128.7	96.2
1927	104.9	105.7	188.6	158.8	114.2	90.0	169.5	95.0	128.2	132.3	97.2
1926	111.7	108.1	198.1	184.1	129.4	95.0	164.2	99.3	130.0	143.7	99.6
1925	113.2	118.1	206.2	190.5	127.4	92.9	158.6	100.6	134.5	146.7	108.3
1924	119.3	127.1	216.1	195.5	139.8	102.2	151.3	99.0	144.0	147.3	112.2
1923	128.0	133.2	245.7	202.9	159.0	98.9	177.1	106.4	151.7	168.8	123.7
1922	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1921	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1920	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA

TABLE 18. - 1/DEATH RATES FOR TEN SELECTED CAUSES BY PREFECTURE: JAPAN 1920-1949 - Cont'd

CONGENITAL DEBILITY (158) - Cont'd

Year	Hiroshima	Hokkaido	Hyogo	Ibaraki	Ishikawa	Iwate	Kagawa	Karoshima	Kanagawa	Kochi	Kumamoto	Kyoto
1949	38.4	39.2	35.9	53.5	50.6	70.7	59.1	39.1	23.0	50.8	44.1	36.1
1948	42.5	41.9	42.8	56.6	60.3	70.2	65.2	39.0	26.7	48.8	42.1	38.2
1947	51.3	42.7	46.8	66.0	65.0	76.4	85.5	41.5	32.7	57.1	43.6	43.7
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	67.8	59.0	64.1	89.2	102.9	128.1	104.2	49.2	46.4	77.9	64.8	67.5
1941	66.0	59.5	62.7	96.1	94.6	125.6	97.5	44.4	47.1	72.8	67.0	66.0
1940	68.5	65.7	62.4	109.8	109.2	142.6	107.6	46.8	52.9	72.5	66.4	69.7
1939	72.3	71.0	74.2	110.1	96.7	132.4	111.5	49.0	59.3	78.8	66.8	69.3
1938	82.9	73.5	77.9	120.3	109.9	141.1	115.2	51.7	61.3	92.5	71.1	80.6
1937	86.0	72.6	78.5	130.3	120.3	148.6	120.4	49.2	61.9	108.1	70.4	85.2
1936	92.8	77.6	93.2	134.1	133.9	156.0	127.4	53.6	62.8	101.4	78.5	92.3
1935	89.6	78.1	85.6	129.0	113.1	144.1	112.1	50.1	67.4	108.7	74.1	83.8
1934	97.9	81.1	96.9	136.9	130.5	159.4	125.8	52.6	66.6	106.4	81.5	92.7
1933	99.6	86.5	99.8	133.1	125.3	167.3	131.9	48.9	71.6	119.2	79.4	91.7
1932	85.5	72.0	90.3	125.1	105.6	149.9	117.7	44.3	63.8	110.9	71.1	80.3
1931	87.7	82.7	91.2	132.8	120.7	155.4	121.5	44.9	68.8	111.3	75.4	82.4
1930	87.8	79.5	89.6	130.0	99.6	152.2	115.6	49.3	71.9	111.0	68.2	82.4
1929	94.7	93.4	107.4	144.1	121.2	169.5	131.8	42.6	85.6	142.5	74.4	91.3
1928	95.6	89.6	103.2	146.6	119.8	179.8	127.1	40.4	84.7	138.2	71.5	86.9
1927	99.6	93.2	104.6	143.7	132.5	174.2	140.7	48.0	88.9	128.0	72.8	91.9
1926	106.0	96.6	113.7	158.5	129.1	180.7	150.3	45.0	86.4	149.5	72.5	104.1
1925	106.8	101.2	108.0	164.4	127.2	181.8	140.1	42.4	107.4	148.7	66.7	106.7
1924	113.5	96.2	118.2	177.7	136.6	210.9	152.3	41.0	107.6	141.0	78.0	107.9
1923	120.6	98.8	127.0	183.7	142.5	225.7	155.2	45.1	102.3	165.9	79.6	112.2
1922	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1921	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1920	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA

TABLE 18. - 1/DEATH RATES FOR TEN SELECTED CAUSES BY PREFECTURE: JAPAN 1920-1949 - Cont'd

CONGENITAL DEBILITY (158) - Cont'd

Year	Mie	Miyagi	Miyazaki	Nagano	Narasaki	Nara	Niigata	Oita	Okayama	Osaka	Saga	Saitama
1949	46.9	48.8	54.2	30.1	55.4	51.1	41.9	55.4	52.6	38.4	70.1	52.0
1948	52.0	55.3	45.6	35.2	48.6	62.5	44.2	55.3	60.6	40.3	72.2	51.9
1947	61.9	53.8	50.4	39.8	55.2	67.8	46.7	71.6	71.1	47.6	87.4	56.7
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	78.9	85.6	68.1	54.1	80.1	95.1	58.5	99.7	92.3	55.8	107.4	80.1
1941	82.6	80.1	66.0	59.9	72.6	98.8	61.4	97.6	92.2	58.5	109.3	93.5
1940	92.3	99.0	69.7	62.8	76.1	102.9	65.5	93.3	96.7	54.6	118.8	98.7
1939	95.8	90.0	73.8	63.4	84.6	109.5	67.6	101.1	103.4	63.1	129.3	103.7
1938	117.4	101.4	75.0	68.6	88.2	119.7	65.7	109.7	110.3	61.6	136.7	105.0
1937	114.1	108.6	80.1	66.4	91.2	121.4	71.5	125.7	121.4	67.6	144.8	111.1
1936	115.0	115.2	91.9	74.7	101.9	125.6	84.4	134.7	135.8	69.4	164.5	119.5
1935	113.1	126.8	79.8	76.3	89.6	122.0	76.3	124.0	121.0	72.7	151.9	108.5
1934	122.5	127.0	80.1	76.3	97.9	135.1	80.6	128.6	134.0	81.3	158.9	121.6
1933	113.3	121.5	82.9	72.8	94.5	135.7	77.8	129.0	142.6	79.4	162.4	125.8
1932	102.6	109.4	71.6	66.3	80.9	122.9	62.7	112.8	125.7	75.2	139.5	111.3
1931	99.8	111.1	69.1	68.9	76.9	121.9	73.7	108.9	130.5	79.7	133.8	122.3
1930	101.5	110.5	67.2	71.0	79.1	123.3	68.5	102.9	121.8	80.8	136.8	111.4
1929	109.9	128.4	79.8	77.7	86.8	139.0	76.6	109.0	146.6	87.8	148.2	132.1
1928	110.3	116.1	66.3	74.5	84.1	130.8	74.6	104.4	142.5	86.9	146.9	132.7
1927	111.5	128.3	70.7	80.1	85.5	141.0	77.4	104.5	142.3	93.3	145.3	134.5
1926	126.1	136.4	77.6	88.3	85.2	143.0	81.4	116.1	158.4	95.4	152.7	142.8
1925	127.5	142.8	73.1	91.4	83.1	141.5	80.8	112.9	145.2	101.0	138.6	149.4
1924	125.8	159.6	77.6	87.7	79.2	161.6	91.8	129.4	164.3	109.0	149.5	155.6
1923	142.8	174.8	73.1	91.7	86.4	154.5	91.4	129.3	171.9	117.5	163.5	185.7
1922	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1921	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1920	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA

TABLE 18. - 1/DEATH RATES FOR TEN SELECTED CAUSES BY PREFECTURE: JAPAN 1920-1949 - Cont'd

CONGENITAL DEBILITY (158) - Cont'd

Year	Shira	Shirane	Shizuoka	Tochi	Tokushima	Tokyo	Tottori	Toyama	Wakayama	Yamagata	Yamaguchi	Yamanashi
1949	54.7	49.7	41.6	37.9	60.8	24.8	48.2	58.3	34.6	65.6	43.8	31.7
1948	53.4	53.8	41.5	37.8	68.5	28.3	51.6	55.0	37.0	64.8	42.5	33.9
1947	64.7	67.9	45.4	46.6	72.6	30.8	58.3	69.8	48.5	70.2	51.5	35.5
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	99.5	94.7	69.0	64.2	96.8	45.5	71.0	91.6	58.8	106.3	57.6	54.2
1941	91.0	100.7	71.8	65.4	102.9	47.2	86.8	86.1	88.5	106.6	87.2	57.3
1940	100.6	98.0	77.6	77.2	105.2	47.1	84.8	96.9	64.9	110.2	54.7	59.9
1939	93.0	115.6	83.3	74.7	110.7	49.5	77.4	86.8	70.2	114.9	63.7	61.8
1938	99.0	120.0	91.1	74.3	118.9	50.7	92.3	96.0	76.6	119.6	69.1	64.4
1937	106.9	123.1	91.9	78.8	122.3	53.8	91.8	96.7	80.7	131.7	80.0	67.1
1936	119.7	127.8	105.1	87.9	127.1	59.0	104.2	101.4	89.0	130.9	81.4	64.2
1935	108.5	124.3	103.6	79.5	120.5	57.0	90.3	97.4	84.3	139.6	82.1	77.5
1934	126.6	129.9	111.6	85.0	130.1	59.7	104.3	107.0	90.7	145.0	83.5	69.2
1933	115.3	137.4	103.7	84.1	127.4	63.7	95.9	108.1	96.4	137.7	83.1	69.8
1932	90.9	116.8	98.3	72.0	118.8	61.1	94.5	79.5	84.5	131.9	63.6	64.5
1931	99.9	115.6	93.8	74.7	125.4	67.7	96.4	90.5	78.7	140.5	72.2	62.5
1930	99.2	109.8	91.5	75.1	116.1	66.6	94.8	82.9	81.9	137.2	68.6	70.7
1929	99.6	119.1	103.8	86.8	125.0	80.1	101.4	86.3	86.5	150.2	76.6	70.9
1928	102.2	112.1	106.8	77.3	124.1	82.7	96.8	92.8	90.3	164.4	69.0	73.0
1927	104.5	113.9	100.2	85.9	110.7	79.6	101.4	94.3	86.8	166.4	72.3	68.2
1926	120.6	117.6	113.3	87.6	127.6	81.4	101.9	99.3	82.6	178.4	82.3	77.8
1925	120.5	124.4	110.9	89.8	122.4	87.4	113.2	95.7	92.3	181.4	71.5	84.6
1924	121.2	121.2	120.7	89.3	132.6	95.9	121.3	99.3	90.3	197.0	78.1	78.7
1923	137.5	127.7	124.3	103.5	140.0	99.1	123.1	97.5	108.8	210.4	90.0	86.5
1922	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1921	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1920	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA

See footnotes at end of table.

TABLE 18. - 1/DEATH RATES FOR TEN SELECTED CAUSES BY PREFECTURE: JAPAN 1920-1949 - Cont'd
8/SENILITY (162)

Year	All Japan	Aichi	Akita	Aomori	Chiba	Ehime	Fukui	Fukuoka	Fukushima	Gifu	Gumma
1949	81.1	81.2	78.8	77.3	99.8	117.0	119.1	69.2	77.0	89.5	69.1
1948	79.9	81.6	82.1	85.2	107.6	116.8	116.8	61.5	77.2	85.7	65.0
1947	101.2	100.1	95.6	85.2	122.7	139.6	138.4	91.7	102.5	112.1	89.9
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	134.1	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	130.4	110.2	132.4	113.6	192.0	181.7	186.1	101.4	141.1	156.5	122.9
1941	123.0	114.6	124.8	106.3	189.4	162.3	183.5	92.7	132.6	156.5	116.4
1940	124.4	119.2	141.6	127.0	173.2	171.1	189.3	90.6	137.5	152.3	125.2
1939	133.3	120.7	136.4	131.7	181.4	189.1	185.5	106.5	136.4	150.2	130.6
1938	138.7	137.9	121.7	120.6	213.7	194.8	193.3	114.3	148.4	187.4	134.9
1937	119.4	115.9	104.1	100.9	172.0	174.3	177.4	99.4	122.3	147.5	103.8
1936	130.7	127.9	110.5	130.2	181.9	180.2	197.2	103.6	135.8	173.1	125.4
1935	114.2	118.3	106.8	92.2	167.0	169.0	172.7	94.1	113.2	145.1	107.7
1934	127.8	135.2	107.1	105.5	181.9	179.1	231.5	95.1	131.4	168.6	119.4
1933	123.4	122.0	105.2	103.4	177.6	179.3	194.3	95.7	118.0	153.7	107.8
1932	116.6	121.1	102.9	98.0	158.3	168.7	184.4	92.9	113.7	143.2	111.1
1931	130.8	144.1	111.5	110.4	184.2	182.7	198.5	108.3	134.1	182.2	123.6
1930	118.6	119.1	103.8	94.7	163.7	168.6	189.8	105.5	118.4	139.4	111.1
1929	124.6	129.2	127.6	120.8	170.4	175.5	207.3	110.4	122.6	141.8	115.9
1928	123.2	142.9	111.8	104.9	185.0	167.5	183.3	108.0	116.6	152.4	107.5
1927	120.8	130.4	121.7	103.1	172.4	164.9	190.1	102.2	123.8	139.7	113.5
1926	111.0	119.5	125.8	107.1	155.7	152.4	154.4	96.1	108.1	119.2	99.4
1925	117.0	135.9	124.2	91.4	166.3	157.6	170.1	98.3	112.8	143.7	108.2
1924	125.5	135.0	127.3	95.5	174.8	176.1	160.8	101.8	122.0	144.2	115.6
1923	131.5	163.4	167.5	114.9	215.4	174.7	172.9	100.5	116.0	154.4	118.0
1922	132.4	141.4	162.7	138.9	196.5	178.5	172.0	103.6	116.0	143.7	119.8
1921	133.6	186.1	141.5	110.2	206.6	186.1	169.7	110.2	119.6	149.0	115.6
1920	131.1	163.3	131.1	93.2	198.3	180.8	175.4	118.5	118.7	149.7	112.6

TABLE 18. - 1/DEATH RATES FOR TEN SELECTED CAUSES BY PREFECTURE: JAPAN 1920-1949 - Cont'd
8/SENILITY (162) - Cont'd

Year	Hiroshima	Hokkaido	Hyogo	Ibaraki	Ishikawa	Iwate	Yamaguchi	Kagoshima	Kanagawa	Kochi	Kumamoto	Kyoto
1949	96.5	55.1	78.3	89.3	111.5	83.2	110.3	91.9	58.9	132.8	92.5	77.5
1948	88.8	53.9	82.5	93.0	97.8	81.7	94.9	79.1	57.2	135.8	83.1	79.9
1947	118.4	85.0	114.9	110.6	107.8	94.0	137.5	103.3	67.3	170.1	106.0	107.0
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	145.3	70.9	131.9	158.9	173.1	149.5	190.0	172.2	87.9	230.0	171.3	129.9
1941	135.3	71.2	116.0	151.1	170.5	137.6	169.6	141.7	86.9	226.8	147.8	117.2
1940	147.1	71.0	119.7	152.7	184.3	143.5	179.2	144.3	93.6	213.6	143.2	122.0
1939	164.6	93.3	136.0	160.0	175.2	149.4	206.7	145.6	93.4	224.7	147.6	124.4
1938	167.9	89.9	145.3	168.9	191.5	140.7	195.7	131.9	101.1	231.6	150.7	130.7
1937	140.1	76.1	125.3	133.8	179.0	133.2	173.2	118.7	85.2	207.3	137.9	104.6
1936	152.4	86.3	137.8	151.0	196.9	145.8	168.2	133.6	97.6	198.6	155.1	117.0
1935	135.9	72.0	123.7	131.8	162.0	121.5	158.7	118.3	74.5	184.5	139.9	101.4
1934	139.5	69.7	143.3	157.3	222.4	137.9	173.4	119.9	89.7	188.5	135.3	112.4
1933	144.9	77.0	144.7	141.1	171.8	148.1	167.3	112.4	89.5	196.9	131.9	112.8
1932	131.6	63.0	134.0	131.9	142.5	142.5	145.2	112.3	79.4	202.7	125.3	111.2
1931	145.2	73.9	148.0	139.8	197.4	148.0	171.8	120.2	92.4	203.3	143.5	128.2
1930	134.3	64.3	136.7	120.6	171.0	138.6	151.5	116.5	81.4	186.8	136.6	118.7
1929	137.2	78.6	135.7	129.1	200.7	141.3	158.2	111.2	89.2	188.2	133.8	121.3
1928	125.9	67.7	129.6	128.8	178.7	143.6	132.8	107.8	96.2	208.2	129.7	119.0
1927	122.1	70.6	132.7	131.7	201.2	146.3	157.2	106.2	84.1	182.3	118.7	115.6
1926	116.8	65.1	121.0	112.2	178.0	129.7	134.9	113.0	86.4	181.4	113.6	108.1
1925	128.0	66.8	124.0	128.9	172.1	144.3	135.2	101.8	80.4	199.9	108.9	112.3
1924	144.6	57.3	139.2	132.2	189.2	144.8	170.3	103.0	98.1	215.0	125.3	115.4
1923	141.5	54.6	143.6	137.7	211.6	140.0	164.1	99.5	118.7	201.2	116.5	118.7
1922	146.6	55.5	142.6	137.7	215.0	164.3	191.1	107.5	95.6	226.2	125.3	118.9
1921	139.4	54.4	141.8	126.8	173.6	141.5	182.2	109.0	104.8	211.6	134.8	116.9
1920	134.5	56.2	149.3	138.4	182.9	131.2	174.8	98.5	101.3	218.5	119.8	132.9

TABLE 18. - $\frac{1}{2}$ DEATH RATES FOR TEN SELECTED CAUSES BY PREFECTURE: JAPAN 1920-1949 - Cont'd

g/SENILITY (162) - Cont'd

Year	Mie	Miyagi	Miyazaki	Nagano	Narasaki	Nara	Miigata	Oita	Okayama	Osaka	Saga	Saitama
1949	84.9	65.4	86.0	81.2	106.5	89.4	69.3	117.0	116.6	56.4	89.4	81.0
1948	92.4	69.7	71.3	82.3	96.4	88.9	73.9	108.8	118.4	63.0	91.0	81.5
1947	119.2	74.5	102.0	101.8	129.7	108.2	83.8	130.8	143.7	84.9	124.5	87.1
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	168.9	117.9	151.6	154.0	170.2	154.4	125.2	192.0	209.8	82.2	162.6	123.3
1941	163.1	110.5	136.6	150.0	152.1	147.5	121.3	175.1	198.8	71.3	144.3	138.2
1940	164.5	108.9	127.9	144.3	149.4	140.5	130.7	177.4	195.4	71.9	139.0	135.3
1939	154.0	112.6	138.8	159.5	173.7	146.7	131.8	185.7	214.8	75.7	142.6	134.8
1938	187.5	110.9	129.5	162.9	178.1	151.5	131.9	194.9	231.3	71.0	160.9	147.3
1937	147.3	90.5	121.4	140.3	157.7	118.2	120.4	169.0	191.3	65.4	141.6	117.7
1936	164.3	105.8	124.4	149.2	182.4	156.7	129.5	189.9	211.3	68.0	183.8	125.6
1935	142.7	86.8	124.0	130.5	153.4	126.2	101.7	167.4	176.0	61.5	146.8	102.1
1934	157.3	96.3	121.1	143.3	155.7	128.3	140.5	174.0	195.3	72.1	146.6	125.5
1933	144.0	96.0	118.5	130.2	161.1	131.6	119.4	182.7	207.8	71.9	141.7	115.1
1932	137.2	87.6	122.9	125.7	156.5	124.3	112.7	173.0	173.0	66.5	141.7	115.6
1931	158.9	101.6	121.9	136.7	165.6	137.8	125.8	194.4	193.3	78.3	154.2	131.2
1930	131.2	93.7	115.5	125.2	164.2	132.8	103.9	175.8	176.4	71.2	158.3	109.1
1929	132.2	94.0	116.2	124.2	168.3	127.6	128.2	177.2	187.5	72.0	150.1	121.6
1928	147.9	95.2	125.0	125.4	169.3	121.6	117.0	175.5	179.0	72.4	147.9	118.3
1927	130.4	93.7	122.4	125.4	162.5	121.8	119.6	171.1	182.9	73.4	129.1	111.5
1926	114.5	78.1	118.8	110.2	152.2	110.2	111.5	165.5	161.4	69.1	135.7	95.3
1925	138.8	84.2	110.5	127.3	141.2	120.2	121.1	162.3	169.1	73.3	119.7	107.8
1924	121.2	91.7	130.0	123.0	154.8	131.6	123.4	201.5	200.3	82.8	135.1	115.3
1923	148.0	97.2	110.0	125.6	159.0	121.1	128.1	187.1	193.1	85.8	142.9	116.2
1922	131.0	106.0	129.5	130.9	163.0	112.4	130.8	211.4	186.5	92.1	144.3	114.8
1921	153.0	96.4	135.8	125.1	170.3	122.7	115.7	219.0	204.6	94.7	146.0	120.4
1920	149.1	90.5	121.5	121.2	159.3	132.8	117.3	215.5	187.6	103.4	142.8	108.9

Table 18, pg.35

TABLE 18. - 1/DEATH RATE'S FOR TEN SELECTED CAUSES BY PREFECTURE: JAPAN 1920-1949 - Cont'd

8/SENILITY (162) - Cont'd

Year	Shira	Shirane	Shizuoka	Tochigi	Tokushima	Tokyo	Tottori	Toyama	Wakayama	Yamagata	Yamaguchi	Yamanashi
1949	97.3	104.5	75.4	76.2	116.3	46.8	120.2	64.9	96.8	73.7	103.1	71.7
1948	98.7	93.9	72.0	72.0	112.5	43.4	121.6	63.3	94.3	74.7	95.1	74.0
1947	127.3	147.1	100.4	79.4	152.4	55.6	158.3	78.3	131.8	84.6	141.0	98.5
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	186.9	224.2	142.6	113.9	220.0	64.1	227.2	115.4	184.3	126.0	164.1	132.3
1941	177.5	221.6	139.7	118.8	202.5	57.5	227.9	111.2	159.0	126.0	153.9	130.3
1940	184.6	235.3	174.9	115.4	200.9	55.7	208.3	125.0	155.3	134.4	146.1	126.8
1939	172.9	273.9	141.8	119.9	236.9	58.4	237.3	117.6	166.0	137.5	189.8	128.2
1938	208.5	266.2	154.3	127.6	234.1	63.2	221.4	124.6	151.6	128.5	201.2	145.0
1937	151.0	207.5	134.1	113.1	216.1	54.2	201.1	117.2	136.9	121.6	166.6	116.8
1936	178.2	237.6	152.2	112.0	228.0	56.0	199.9	115.4	154.0	118.6	186.8	136.0
1935	148.0	204.0	129.6	96.6	209.0	47.6	193.9	96.5	149.1	98.4	160.4	119.7
1934	172.5	238.0	150.2	118.0	211.3	50.7	206.1	130.1	155.4	118.0	184.0	136.9
1933	164.0	221.2	145.5	112.0	202.8	53.2	188.4	99.6	148.7	111.9	166.8	133.0
1932	147.0	211.7	136.8	96.5	190.0	47.1	194.5	104.6	130.4	107.8	161.4	115.6
1931	175.4	250.6	144.3	115.3	218.7	52.3	203.4	124.5	160.2	120.0	176.2	116.5
1930	155.0	261.5	128.4	93.0	192.6	45.9	188.4	97.2	144.0	103.8	177.0	111.4
1929	143.9	264.7	140.4	93.7	184.1	52.1	196.1	100.6	133.1	119.4	174.1	108.2
1928	148.4	247.8	150.2	95.9	193.0	55.9	174.0	94.9	151.5	111.4	162.1	104.2
1927	139.8	233.7	131.3	93.0	207.0	50.3	196.6	97.7	122.6	108.7	175.1	115.2
1926	119.1	221.4	119.5	94.6	173.8	46.5	152.2	88.3	116.5	105.2	153.1	102.0
1925	141.5	224.0	136.0	92.5	184.1	50.6	169.8	90.2	132.6	105.9	147.9	105.9
1924	142.3	250.8	131.9	93.7	215.0	60.7	198.3	91.0	144.1	100.5	184.8	111.8
1923	162.4	277.7	164.7	98.0	196.1	69.3	211.1	97.2	128.3	110.7	193.1	114.9
1922	147.9	244.2	140.1	98.2	215.4	66.9	195.3	104.5	149.7	117.2	189.3	119.3
1921	166.1	240.5	159.9	96.4	203.7	65.8	209.8	85.1	145.7	98.1	184.3	119.7
1920	146.2	231.8	150.1	93.7	197.3	66.8	196.0	86.3	146.3	107.0	184.2	121.9

See footnotes at end of table.

TABLE 10. - 1/DEATH RATES FOR TEN SELECTED CAUSES BY PREFECTURE: JAPAN 1920-1949 - Cont'd

ACCIDENTAL DEATHS (169-195)

Year	All Japan	Alchi	Akita	Aomori	Chiba	Chime	Fukui	Fukuoka	Fukushima	Gifu	Gumma
1949	42.0	35.2	44.6	39.3	32.8	50.7	57.0	57.3	44.9	59.0	38.2
1948	48.6	35.1	47.5	49.3	37.5	38.3	520.1	60.5	47.2	38.5	42.0
1947	49.3	39.3	50.1	45.9	38.5	47.6	53.6	68.3	50.2	45.8	71.1
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	45.1	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	42.3	34.2	47.7	43.1	39.9	45.2	40.4	56.4	47.2	40.6	34.2
1941	39.4	38.4	42.4	38.9	38.5	42.7	39.6	55.4	43.6	40.3	34.5
1940	39.3	28.7	42.3	40.2	37.4	38.4	52.0	53.6	45.3	37.9	37.6
1939	41.1	31.7	49.8	41.1	37.8	37.9	47.3	66.4	47.8	42.0	37.0
1938	44.6	32.5	50.3	41.4	45.2	41.4	39.1	64.6	52.5	43.6	44.6
1937	42.5	33.7	50.2	42.5	34.3	42.9	46.3	61.2	49.2	41.0	47.0
1936	43.0	31.7	76.7	48.6	40.2	45.2	51.1	66.2	53.0	43.7	43.1
1935	41.8	32.9	47.2	43.7	36.0	48.0	45.0	63.6	55.5	41.1	56.8
1934	46.8	33.8	44.7	41.6	32.8	42.6	52.3	56.6	44.6	43.6	37.0
1933	44.6	31.1	47.3	47.0	36.6	46.3	44.8	54.7	47.7	42.7	31.8
1932	40.3	32.9	43.8	43.5	32.2	43.9	48.6	51.9	46.1	43.7	40.1
1931	38.7	32.3	40.6	43.2	34.6	45.8	42.1	52.8	46.8	37.7	33.7
1930	40.8	32.5	41.9	44.0	32.4	41.2	44.5	64.1	47.5	36.9	35.3
1929	42.0	34.0	48.9	39.4	32.0	45.0	43.2	61.2	51.1	42.8	35.0
1928	41.6	32.8	44.6	44.3	34.6	45.3	46.5	62.3	45.8	36.5	33.7
1927	47.6	33.9	45.1	45.3	36.8	43.6	54.1	60.7	59.6	41.5	34.9
1926	42.2	34.9	45.5	44.8	36.6	47.5	44.8	59.8	47.9	35.3	40.7
1925	41.9	37.2	42.2	42.3	33.5	46.9	48.3	59.9	51.1	39.1	36.4
1924	43.8	38.2	50.2	46.8	36.7	42.7	58.1	56.0	57.8	43.4	40.3
1923	123.6	39.1	52.3	45.0	133.2	45.1	48.4	57.7	53.9	45.7	40.2
1922	44.6	36.3	51.1	52.1	41.1	48.6	51.4	47.0	65.6	41.0	38.1
1921	43.3	35.8	45.3	39.2	39.5	42.6	50.8	56.2	51.1	42.1	43.8
1920	46.8	36.3	54.8	51.4	40.0	52.0	44.2	64.4	59.1	42.9	43.2

TABLE 18. - 1/DEATH RATES FOR TEN SELECTED CAUSES BY PREFECTURE: JAPAN 1920-1949 - Cont'd
ACCIDENTAL DEATHS (162-195) - Cont'd

Year	Hiroshima	Hokkaido	Hyogo	Ibaraki	Ishikawa	Iwate	Yamagata	Kagoshima	Yamaguchi	Kochi	Kumamoto	Kyoto
1949	44.7	54.2	42.1	34.3	41.2	55.2	46.1	47.6	45.6	46.9	41.6	31.3
1948	49.5	55.7	43.0	34.2	46.4	86.5	51.3	39.6	47.9	48.3	37.8	34.5
1947	62.5	55.5	54.3	38.7	49.6	60.5	50.8	36.9	52.1	61.0	42.9	39.3
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	51.5	62.7	43.4	36.5	39.4	48.5	63.2	40.3	41.9	51.8	38.3	30.8
1941	45.9	66.2	36.6	40.7	36.3	44.5	47.0	36.0	43.3	50.9	32.7	29.1
1940	44.0	66.2	38.6	30.9	40.8	50.3	40.9	35.2	37.0	44.7	35.7	25.1
1939	42.6	67.1	43.0	35.2	38.0	60.5	45.0	34.7	40.2	52.2	35.0	25.9
1938	49.4	67.0	60.2	43.0	42.9	59.2	51.9	57.0	47.7	53.4	40.4	25.8
1937	48.7	65.4	43.8	37.4	42.2	62.7	52.5	32.2	45.0	52.7	35.3	30.6
1936	46.5	61.1	44.9	38.5	36.4	52.6	49.0	38.2	42.2	55.9	35.3	31.9
1935	43.7	60.2	45.8	34.4	35.9	53.2	61.8	29.7	40.4	52.3	37.5	32.4
1934	48.2	114.8	53.2	33.1	44.1	45.0	46.1	30.3	37.8	59.2	36.5	48.9
1933	43.2	59.7	44.6	36.2	43.7	283.7	45.5	29.1	42.5	46.8	41.8	31.3
1932	42.9	60.2	44.9	32.7	35.8	47.7	47.5	29.3	41.0	44.3	35.2	32.4
1931	40.0	57.9	41.2	32.5	40.8	49.8	50.9	31.0	38.3	46.9	34.1	32.1
1930	41.0	59.5	40.2	33.3	41.0	48.1	50.8	34.5	38.8	40.4	38.7	32.8
1929	43.3	67.2	43.5	37.0	42.9	47.2	54.7	33.3	43.1	45.8	39.4	34.1
1928	42.2	60.2	44.8	34.0	40.3	48.1	49.1	30.3	42.1	46.2	42.8	30.0
1927	41.7	70.5	45.6	34.3	39.9	46.4	51.3	32.7	36.8	41.1	66.8	231.0
1926	52.4	71.0	43.7	32.5	35.3	47.0	49.3	33.5	40.1	43.4	41.3	32.6
1925	45.4	67.6	60.5	35.0	36.1	48.6	48.7	27.0	41.0	42.6	37.8	31.3
1924	44.4	66.0	44.4	33.7	38.8	55.4	56.7	29.7	46.0	41.9	37.6	31.0
1923	53.3	69.1	51.4	38.9	37.6	59.3	47.0	30.2	1195.4	41.5	42.3	36.5
1922	47.1	73.3	46.6	38.2	45.0	54.9	56.0	29.2	41.1	45.0	39.1	35.2
1921	49.3	70.3	48.2	33.8	30.6	54.4	47.8	31.8	38.8	40.5	36.9	36.0
1920	52.4	94.6	45.4	45.9	40.9	69.3	53.6	30.0	50.0	65.4	39.3	32.4

TABLE 18. - 1/DEATH RATES FOR TEN SELECTED CAUSES BY PREFECTURE: JAPAN 1920-1949 - Cont'd

ACCIDENTAL DEATHS (169-195) - Cont'd

Year	Mie	Miyagi	Miyazaki	Nagano	Nagasaki	Nara	Niigata	Oita	Okayama	Osaka	Saga	Saitama
1949	35.2	41.8	43.2	37.5	51.2	31.7	41.5	42.9	41.1	35.5	55.7	30.0
1948	35.8	44.8	42.1	35.4	52.5	33.7	42.1	44.0	59.7	38.2	61.1	34.1
1947	41.8	47.9	45.2	41.1	57.9	42.1	46.2	45.9	53.2	49.1	51.6	52.3
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	41.8	42.2	48.1	39.1	55.4	36.4	46.5	46.7	48.3	28.5	57.8	36.4
1941	36.8	35.7	47.4	38.1	48.7	35.0	41.0	45.5	43.7	28.1	55.5	28.7
1940	36.2	33.3	39.2	37.2	49.7	35.1	44.0	44.0	42.8	32.3	47.0	35.2
1939	37.3	37.0	50.1	39.5	48.5	30.0	52.0	50.2	40.1	29.4	47.4	37.9
1938	37.4	37.6	47.7	44.1	48.3	31.0	50.2	42.8	46.7	32.4	49.9	39.0
1937	37.1	41.0	41.4	44.3	42.2	33.3	48.2	46.3	47.5	32.1	45.1	35.0
1936	37.4	37.4	46.5	42.1	46.6	32.2	42.5	48.3	46.4	31.0	52.2	35.5
1935	39.1	38.7	42.7	41.1	40.8	38.5	47.3	45.5	43.2	32.9	44.7	31.6
1934	36.0	35.3	41.0	40.7	45.6	37.4	46.2	40.3	50.1	79.6	43.7	30.7
1933	34.6	55.8	42.2	38.8	44.0	33.1	46.0	47.0	43.5	41.3	44.3	72.8
1932	37.4	37.9	37.4	41.0	36.6	39.9	46.7	40.6	45.1	36.2	44.9	37.2
1931	38.4	37.7	44.0	36.7	41.1	30.4	41.6	41.7	36.3	37.2	40.3	31.4
1930	37.8	37.9	46.4	36.8	42.2	32.4	46.9	47.3	44.0	31.5	51.2	32.3
1929	36.8	39.0	38.4	43.2	51.6	37.7	44.9	41.2	43.7	32.4	46.6	37.0
1928	36.8	42.9	41.9	45.0	42.7	31.4	44.7	50.7	44.1	32.9	52.8	32.7
1927	35.8	41.5	43.4	44.5	48.5	31.6	52.1	41.7	43.9	33.1	51.2	37.2
1926	35.7	39.3	37.3	41.3	35.1	32.9	32.9	43.9	44.4	34.2	53.0	31.9
1925	38.1	40.5	40.5	39.0	51.6	38.4	45.1	41.7	43.0	30.2	47.7	32.1
1924	33.2	41.0	42.5	42.9	50.9	35.7	46.8	49.5	44.9	33.5	48.8	34.5
1923	36.8	43.2	42.0	49.5	47.0	35.8	51.2	44.1	46.8	36.2	52.0	52.8
1922	35.9	46.8	37.6	41.8	47.9	33.1	59.8	45.6	42.9	37.3	48.6	35.0
1921	43.4	41.9	44.0	44.5	42.1	40.8	44.5	53.3	46.3	33.9	47.1	35.5
1920	36.8	43.6	42.5	42.8	43.9	33.1	50.9	39.3	48.2	33.8	47.6	34.1

TABLE 18. - 1/DEATH RATES FOR TEN SELECTED CAUSES BY PREFECTURE: JAPAN 1920-1949 - Cont'd

ACCIDENTAL DEATHS (169-195) - Cont'd

Year	Shira	Shimane	Shizuoka	Tochigi	Tokushima	Tokyo	Tottori	Toyama	Yakayama	Yamagata	Yamaguchi	Yamanashi
1949	29.4	44.7	35.9	35.5	45.9	34.2	38.8	55.7	33.7	42.2	57.6	42.8
1948	33.8	51.1	37.0	34.9	44.9	38.6	37.5	54.1	37.4	44.1	60.9	42.8
1947	46.1	48.0	40.0	61.1	52.5	40.1	48.7	56.5	41.7	47.5	64.4	41.1
1946	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1945	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1944	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1943	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1942	36.4	47.9	39.5	38.2	55.9	24.4	47.6	47.3	40.3	44.7	96.3	41.7
1941	33.2	45.9	38.0	35.0	45.5	23.7	43.3	44.1	41.4	42.1	55.6	40.1
1940	37.2	42.7	32.2	35.0	48.1	25.1	39.9	53.7	38.0	40.3	51.2	43.7
1939	38.2	45.4	37.8	40.8	50.6	26.4	40.9	54.7	30.8	41.4	54.3	35.3
1938	37.3	51.0	42.0	42.4	56.5	27.6	46.5	57.4	40.0	45.0	57.0	43.3
1937	37.8	46.0	39.7	40.3	52.6	28.4	47.4	59.3	45.7	44.2	58.5	39.2
1936	36.4	50.0	41.3	42.2	50.0	26.4	40.8	59.1	36.8	46.7	52.5	37.4
1935	36.1	44.2	37.4	39.1	54.8	27.6	38.3	55.5	37.5	39.7	51.7	41.9
1934	44.0	47.7	36.9	38.3	56.6	27.8	58.8	49.8	41.9	40.1	48.7	34.3
1933	33.9	40.6	41.0	36.6	51.5	29.1	43.8	51.0	35.5	41.7	50.4	34.7
1932	35.2	43.4	37.5	37.6	45.5	31.4	38.9	46.7	37.2	46.2	46.5	37.6
1931	33.8	45.2	32.5	33.0	45.0	25.8	35.1	46.0	34.8	40.9	52.9	36.7
1930	35.9	44.5	47.0	36.2	50.2	28.1	45.6	49.3	32.6	41.4	51.0	39.5
1929	36.5	50.6	39.5	40.8	54.6	30.8	36.1	48.0	39.2	43.3	46.2	37.1
1928	33.0	47.0	36.4	36.6	51.7	30.4	37.7	50.4	31.7	44.3	44.0	42.3
1927	35.2	43.4	32.9	36.3	43.0	30.1	39.2	47.5	37.7	45.2	51.5	37.6
1926	37.9	44.6	37.2	35.0	47.3	28.0	44.2	45.6	37.1	46.3	58.3	41.5
1925	32.2	39.6	38.2	35.8	53.1	31.6	39.0	45.4	33.9	41.8	47.2	35.1
1924	35.1	46.1	36.6	39.8	58.0	38.5	43.9	53.1	33.4	50.0	57.8	36.0
1923	33.6	47.2	62.7	39.0	57.1	716.9	46.1	50.3	39.4	48.3	54.2	44.4
1922	36.9	49.4	41.4	41.1	50.8	28.4	42.6	51.5	37.8	45.4	52.5	40.8
1921	40.8	47.6	42.1	38.1	52.9	28.5	42.5	46.7	43.8	49.9	47.9	44.2
1920	38.6	49.1	41.0	50.6	60.7	31.8	37.8	48.2	36.1	53.7	49.6	52.4

See footnotes on next page.

TABLE 18. - 1/DEATH RATES FOR TEN SELECTED CAUSES BY PREFECTURE: JAPAN 1920-1949 - Cont'd

Footnotes:

- 1/ Data for 1949 are provisional.
Rates are the number of deaths per 100,000 population, estimated as of 1 October 1920-1945 and 1 July 1946-1949.
Estimated populations, 1920-1945 included civilian and military personnel in Japan only, except in 1943 when Okinawa and Karafuto were included. Estimated population, 1946-1949 are of the civilian population of Japan Proper.
 - 2/ Tuberculosis (all forms). 1920-1943: includes chronic pleuro pneumonia and all adenopathy; excludes spondylitis. 1944-1946: are estimated of the Ministry of Welfare. 1947-1949: excludes chronic pleuro pneumonia and nontubercular adenopathy; includes spondylitis.
 - 3/ Intracranial lesions of vascular origin. 1920-1932: includes alcoholic cerebral apoplexy, congestion and a small amount of paralysis not due to intracranial lesions of vascular origin. 1933-1943: includes alcoholic cerebral apoplexy and congestion. 1947-1949: excludes alcoholic apoplexy and congestion.
 - 4/ Diseases of the heart. 1920-1943: includes diphtheric endocarditis. 1947-1949: excludes diphtheric endocarditis.
 - 5/ Pneumonia (all forms). 1920-1943: excludes typhoid pneumonia. 1947-1949: includes typhoid pneumonia.
 - 6/ Diarrhea, enteritis and ulceration of intestines (all areas). 1920-1943: includes malnutrition 1 year of age and over. 1947-1949: excludes malnutrition 1 year of age and over.
 - 7/ Nephritis (all forms) 1920-1943: includes alcoholic nephritis. 1947-1949: excludes alcoholic nephritis.
 - 8/ Senility. 1920-1922: excludes senility with mention of senile dementia, psychosis and paresis. 1923-1943 and 1947-1949: includes senility with mention of senile dementia.
- "NA" indicates data are not available.
- Sources: Rates were computed by Public Health and Welfare Section, HQ, SCAP.
- Sources of original data were: 1920-1943: Final Vital Statistics Annual Report, Cabinet Bureau of Statistics.
1947-1948: Final Vital Statistics Annual Reports, Ministry of Welfare.
1949: Monthly Vital Statistics Schedule Reports, Ministry of Welfare.

TABLE 19. - 1/DEATHS FROM THE TEN LEADING CAUSES OF DEATH: JAPAN, 1948-1949

List No.	Cause	*1949	1948
12-22	2/ Tuberculosis (all forms)	138,764	143,909
83	3/ Intracranial lesions of vascular origin	100,650	94,329
119-120	4/ Diarrhea, enteritis and ulceration of intestines	72,186	83,264
162	5/ Senility	66,699	64,074
45-55	Cancer and other malignant tumors	58,890	55,677
107-109	6/ Pneumonia (all forms)	56,636	52,978
90-95	7/ Diseases of the heart	52,825	49,046
158	Conrenital debility	37,285	38,204
169-195	Accidental deaths	34,498	38,960
130-132	8/ Nephritis (all forms)	33,859	36,561

TABLE 19. - 1/ DEATHS FROM THE TEN LEADING CAUSES OF DEATH: JAPAN, 1948-1949, Cont'd

Footnotes:

- * Data for 1949 are provisional
- 1/ Data refer to deaths of Japanese nationals in Japan. Data for 1948 are from Final Annual Report, Ministry of Welfare, and for 1949 from Monthly Schedule Reports, Ministry of Welfare.
- 2/ Includes spondylitis and excludes chronic pleuro pneumonia and nontuberculous adenopathy.
- 3/ Excludes alcoholic epilepsy and congestion.
- 4/ Excludes malnutrition 1 year of age and over.
- 5/ Includes senility with mention of senile dementia.
- 6/ Includes typhoid pneumonia.
- 7/ Excludes diphtheric endocarditis.
- 8/ Excludes alcoholic nephritis.

This Table corresponds to Table 15, Statistical Annex, Public Health and Welfare in Japan, 1948.

TABLE 20. - 1/DEATH RATES FOR THE TEN LEADING CAUSES OF DEATH: JAPAN, 1948-1949
(Rates per 100,000 population)

List No.	Cause	1949	1948
13-22	<u>2</u> / Tuberculosis (all forms)	168.8	179.4
83	<u>3</u> / Intracranial lesions of vascular origin	122.4	117.6
119-120	<u>4</u> / Diarrhea, enteritis and ulceration of intestines	87.8	103.8
162	<u>5</u> / Senility	81.1	79.9
45-55	Cancer and other malignant tumors	71.6	69.4
107-109	<u>6</u> / Pneumonia (all forms)	68.9	66.1
90-95	<u>7</u> / Diseases of the heart	64.3	61.2
158	Congenital debility	45.4	47.6
169-195	Accidental deaths	42.0	48.6
130-132	<u>9</u> / Nephritis (all forms)	41.2	45.6

TABLE 20. - 1/DEATH RATES FOR THE TEN LEADING CAUSES OF DEATH: JAPAN, 1948-1949, Cont'd
(Rates per 100,000 population)

Footnotes:

* Data for 1949 are provisional.

1/Rates are the number of deaths per 100,000 population estimated as of 1 July of the year concerned and were calculated by the Public Health and Welfare Section, HQ, SCAP. Data refer to deaths of Japanese nationals in Japan. Data for 1948 are from Final Annual Report, "Ministry of Welfare, and for 1949 from Monthly Schedule Reports, Ministry of Welfare.

2/Includes spondylitis and excludes chronic pleuro pneumonia and nontuberculous adenomathy.

3/Excludes alcoholic epilepsy and convulsion.

4/Excludes malnutrition 1 year of age and over.

5/Includes senility with mention of senile dementia.

6/Includes typhoid pneumonia.

7/Excludes diphtheric endocarditis.

8/Excludes alcoholic nephritis.

This Table corresponds to Table 16, Statistical Annex, Public Health and Welfare in Japan, 1948.

TABLE 21. - RANK ORDER OF THE TEN LEADING CAUSES OF DEATH: JAPAN, 1948-1949

List No.	Cause	1949	1948
13-22	2/ Tuberculosis (all forms)	1	1
83	3/ Intracranial Lesions of vascular origin	2	2
119-120	4/ Diarrhea, enteritis and ulceration of intestines	3	3
162	5/ Senility	4	4
45-55	Cancer and other malignant tumors	5	5
107-109	6/ Pneumonia (all forms)	6	6
90-95	7/ Diseases of the heart	7	7
158	Congenital debility	8	9
169-195	Accidental deaths	9	8
130-132	8/ Nephritis (all forms)	10	10

TABLE 21. - RANK ORDER OF THE TEN LEADING CAUSES OF DEATH: JAPAN, 1948-1949, Cont'd

Footnotes:

- * Data for 1949 are provisional.
- 2/ Includes spondylitis and excludes chronic pleuro pneumonia and nontuberculous adenopathy.
- 3/ Excludes alcoholic epilepsy and congestion.
- 4/ Excludes malnutrition 1 year of age and over.
- 5/ Includes senility with mention of senile dementia.
- 6/ Includes typhoid pneumonia.
- 7/ Excludes diphtheric endocarditis.
- 8/ Excludes alcoholic nephritis.

This Table corresponds to Table 17, Statistical Annex, Public Health and Welfare in Japan, 1948.

TABLE 22. - DEATHS BY AGE: JAPAN, 1948

All ages	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60 & over	Unknown Age
950,610	245,001	22,291	13,620	29,575	50,546	40,210	33,990	33,677	32,823	36,981	42,359	50,243	318,631	753

Footnotes: Data refer to deaths of Japanese nationals in Japan. Data are from Final Annual Report, Ministry of Welfare. This Table corresponds to Table 18, Statistical Annex, Public Health and Welfare in Japan, 1948.

TABLE 23. - DEATH RATES BY AGE: JAPAN 1948
(Rates per 1,000 population)

All ages	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60 & over
11.8	24.1	2.4	1.5	3.5	6.8	7.0	6.5	6.8	7.6	9.2	12.9	18.7	52.5

Footnotes: Rates are per 1,000 population as of 1 October, and refer only to deaths of Japanese nationals in Japan. All rates were computed by Public Health and Welfare Section, GHQ, SCAP. Data are from Final Annual Report, Ministry of Welfare.

This Table corresponds to Table 19, Statistical Annex, Public Health and Welfare in Japan, 1948.

TABLE 24. 1/DEATHS FROM TUBERCULOSIS (ALL FORMS) BY AGE: JAPAN, 1948-1949

Year	All ages	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60 & over	Unknown Age
*1949	138,764	7,172	3,174	2,787	12,608	26,108	21,565	15,525	12,629	9,477	7,695	6,480	5,295	8,199	50
1948	143,909	5,961	3,163	3,222	14,411	29,304	22,323	16,660	12,862	9,506	7,838	6,335	5,079	7,183	62

Footnotes * 1949 data are provisional.

1/ Tuberculosis (all forms) (International List Nos. 13-22) includes spondylitis and excludes chronic pleuro pneumonia and nontuberculous adenomathy 1948-1949. Data refer to Japanese nationals in Japan. Data for 1948 are from Final Annual Report, Ministry of Welfare, and for 1949 from Special Report, Health Statistics, Ministry of Welfare.

This Table corresponds to Table 20, Statistical Annex, Public Health and Welfare in Japan, 1948.

TABLE 25. - 1/DEATH RATES FOR TUBERCULOSIS (ALL FORMS) BY AGE: JAPAN, 1948-1949
(Rates per 100,000 population)

Year	All ages	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60 & over
*1949	168.0	67.8	32.2	31.7	146.5	338.1	357.2	303.5	250.4	212.7	190.0	192.1	195.5	130.3
1948	178.3	56.7	33.5	36.3	169.2	393.5	368.9	320.9	258.0	220.6	196.5	192.7	189.0	118.4

Footnotes: * 1949 data are provisional.

1/ Rates are per 100,000 population as of 1 October of each year and were computed by Public Health and Welfare Section, GHQ, SCAP. Data refer to Japanese nationals in Japan. Data for 1948 are from Final Annual Report, Ministry of Welfare, and for 1949 from Special Report, Health Statistics, Ministry of Welfare.

This Table corresponds to Table 21, Statistical Annex, Public Health and Welfare in Japan, 1948.

TABLE 26. - LIVE BIRTHS, MATERNAL DEATHS AND MATERNAL DEATH RATES:
JAPAN, 1947-1949

Year	Live Births	(140-150) Maternal Deaths	Maternal death rates per 1,000 live births
*1949	2,723,873	4,495	1.7
*1948	2,709,871	4,437	1.6
1947	2,678,792	4,488	1.7

Footnotes:

- * Live births data for 1948 and 1949 are provisional.
Maternal death data for 1949 are provisional.
Data refer to events occurring to Japanese nationals in Japan.
Rates are per 1,000 live births in same period and were computed
by Public Health and Welfare Section, GHQ, SCAP.
Data for 1947 and maternal death data for 1948 are from Final
Annual Report, Ministry of Welfare.
Data for 1949 and birth data for 1948 are from Monthly Schedule
Report, Ministry of Welfare.
This Table corresponds to Table 22, Statistical Annex, Public
Health and Welfare in Japan, 1948.

TABLE 27. - 1/CASES OF SELECTED COMMUNICABLE DISEASES: JAPAN, 1948-1949

Year	Population	Typhoid fever	Paratyphoid fever	Plague	Cholera	Epidemic meningitis	Scarlet fever	Diphtheria	Japanese		
									Dysentery	Malaria	"B" encephalitis
									Small pox	Small pox	typhus fever
1949	82,200,000	6,489	2,226	-	-	1,467	4,667	14,825	24,001	3,732	1,284
1948	80,200,000	9,426	2,892	-	-	2,035	2,924	16,198	14,628	4,940	7,208
											121
											474

Footnotes: * Data for 1949 are provisional.

1/Data refer to cases occurring among civilian population in Japan, and are from Weekly Report, Ministry of Welfare.

2/Excludes croup.

3/Includes ekiri.

A dash (-) indicates that no cases were reported.

This Table corresponds to Table 23, Statistical Annex, Public Health and Welfare in Japan, 1948.

TABLE 28. - 1/CASE RATES FOR SELECTED COMMUNICABLE DISEASES: JAPAN, 1948-1949
(Rates per 100,000 population)

Year	Typhoid fever	Paratyphoid fever	Plague	Cholera	Epidemic meningitis	Scarlet fever	Diphtheria	Dysentery	Japanese		
									Malaria	Small pox	"B" encephalitis
									Small pox	Small pox	typhus fever
1949	7.9	2.7	-	-	1.8	5.7	18.0	29.2	4.5	0.2	1.6
1948	11.8	3.6	-	-	2.6	3.7	20.3	18.3	6.2	0.0	9.0
											0.1
											0.6

Footnotes: * Data for 1949 are provisional.

1/Rates are the number of cases per 100,000 population and were computed by Public Health and Welfare Section, GHQ, SCAP. Case data refer to civilian population in Japan, and are from Weekly Reports, Ministry of Welfare.

2/Excludes croup.

3/Includes ekiri. "NA" indicates data not available.

A dash (-) indicates that no cases were reported. A rate of 0.0 indicates that there were some cases but the rate was less than 0.1.

This Table corresponds to Table 24, Statistical Annex, Public Health and Welfare in Japan, 1948.

TABLE 29. - 1/CASES OF SELECTED COMMUNICABLE DISEASES BY MONTH: JAPAN, 1948-1949

TYPHOID FEVER

Year	Total Cases	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	2/Unknown
* 1949	6,489	3/524	375	249	3/327	354	605	3/872	865	781	3/684	376	3/477	-
1948	9,426	3/553	403	358	425	3/928	802	3/596	1,244	1,013	3/895	548	661	-

PARATYPHOID FEVER

Year	Total Cases	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	2/Unknown
* 1949	2,226	3/219	117	96	3/118	124	252	3/318	319	244	3/175	119	3/125	-
1948	2,892	3/191	117	109	145	3/321	298	3/458	401	311	3/248	120	173	-

PLAGUE

Year	Total Cases	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	2/Unknown
* 1949	-	3/-	-	-	3/-	3/-	-	3/-	-	-	3/-	-	3/-	-
1948	-	3/-	-	-	-	3/-	-	3/-	-	-	3/-	-	-	-

CHOLERA

Year	Total Cases	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	2/Unknown
1949	-	3/-	-	-	3/-	3/-	-	3/-	-	-	3/-	-	3/-	-
1948	-	3/-	-	-	-	3/-	-	3/-	-	-	3/-	-	-	-

TABLE 29. - 1/CASES OF SELECTED COMMUNICABLE DISEASES BY MONTH: JAPAN, 1948-1949, Cont'd

EPIDEMIC MENINGITIS

Year	Total Cases	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	2/ Unknown
*1949	1,467	3/112	109	159	3/189	122	91	3/126	104	187	3/137	57	3/ 74	-
1948	2,035	3/161	177	258	261	3/200	126	3/139	289	171	3/112	59	82	-

SCARLET FEVER

Year	Total Cases	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	2/ Unknown
*1949	4,667	3/518	317	316	3/438	480	493	3/426	208	191	3/319	394	3/567	-
1948	2,924	3/290	168	164	253	3/359	230	3/287	154	151	3/229	249	390	-

DIPHTHERIA

Year	Total Cases	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	2/ Unknown
*1949	14,825	3/1,149	1,597	1,468	3/1,568	1,066	875	3/875	542	591	3/1,273	1,324	3/1,797	-
1948	16,198	3/2,101	1,596	1,590	1,401	3/1,510	927	3/736	594	799	3/1,713	1,565	1,666	-

DYSENTERY

Year	Total Cases	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	2/ Unknown
*1949	24,001	3/168	137	218	3/212	549	1,286	3/4,883	7,055	5,371	3/3,000	593	3/429	-
1948	14,628	3/126	154	292	290	3/607	1,093	3/3,852	3,925	2,547	3/1,228	342	179	-

TABLE 29. - 1/CASES OF SELECTED COMMUNICABLE DISEASES BY MONTH: JAPAN, 1948-1949, Cont'd
SMALLPOX

Year	Total Cases	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	2/Unknown
*1949	124	3/2	2	12	3/25	56	17	3/6	-	-	3/-	3	3/1	-
1948	29	3/2	2	1	4	3/5	1	3/7	-	1	3/4	1	1	-

TYPHUS FEVER

Year	Total Cases	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	2/Unknown
*1949	121	3/31	24	9	3/12	6	3	3/7	-	2	3/5	5	3/17	-
1948	474	3/73	58	34	138	3/36	54	3/23	2	3	3/3	13	3/37	-

MALARIA

Year	Total Cases	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	2/Unknown
*1949	3,732	3/114	86	107	3/207	411	645	3/897	507	377	3/211	85	3/85	-
1948	4,940	3/262	212	232	293	3/429	661	3/1,003	788	514	3/322	116	108	-

JAPANESE "B" ENCEPHALITIS

Year	Total Cases	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	2/Unknown
*1949	1,284	3/1	-	-	3/-	-	2	3/12	70	717	3/436	43	3/3	-
1948	7,208	3/-	-	-	-	3/-	4	3/20	4,907	2,097	3/180	-	-	-

TABLE 29. - 1/CASES OF SELECTED COMMUNICABLE DISEASES BY MONTH: JAPAN, 1948-1949, Cont'd

Footnotes:

- * Data for 1949 are provisional.
- 1/ Data refer to cases of communicable disease among civilian population in Japan, and are from Weekly Reports, Ministry of Welfare.
- 2/ Delays and corrections for the 12 month period not contained in any monthly summary.
- 3/ Monthly total for five-week period. All other monthly totals are for four-week periods.
- "NA" indicates that data are not available.
- A dash (-) indicates that no cases were reported.

This Table corresponds to Table 25, Statistical Annex, Public Health and Welfare in Japan, 1948.

TABLE 30. - 1/CASE RATES FOR SELECTED COMMUNICABLE DISEASES BY MONTH: JAPAN, 1948-1949
(Rates per 100,000 population)

TYPHOID FEVER

Year	Annual	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
* 1949	7.9	6.8	6.0	4.0	4.2	5.7	9.8	11.2	13.9	12.6	8.8	6.1	6.2
1948	11.8	7.2	6.6	5.8	6.9	12.1	13.1	20.8	20.3	16.5	11.7	8.9	10.8

PARATYPHOID FEVER

Year	Annual	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
* 1949	2.7	2.8	1.9	1.5	1.5	2.0	4.1	4.1	5.1	3.9	2.3	1.9	1.6
1948	3.6	2.5	1.9	1.8	2.4	4.2	4.9	6.0	6.5	5.1	3.2	2.0	2.8

PLAQUE

Year	Annual	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
* 1949	-	-	-	-	-	-	-	-	-	-	-	-	-
1948	-	-	-	-	-	-	-	-	-	-	-	-	-

CHOLERA

Year	Annual	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
* 1949	-	-	-	-	-	-	-	-	-	-	-	-	-
1948	-	-	-	-	-	-	-	-	-	-	-	-	-

TABLE 30. - 1/CASE RATES FOR SELECTED COMMUNICABLE DISEASES BY MONTH: JAPAN, 1948-1949, Cont'd
(Rates per 100,000 population)

EPIDEMIC MENINGITIS

Year	Annual	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
*1949	1.8	1.4	1.8	2.6	2.4	2.0	1.5	1.6	1.7	3.0	1.8	0.9	1.0
1948	2.6	2.1	2.9	4.2	4.3	2.6	2.1	1.8	4.7	2.8	1.5	1.0	1.3

SCARLET FEVER

Year	Annual	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
*1949	5.7	6.7	5.1	5.1	5.6	7.7	7.9	5.5	7.4	3.1	4.1	6.4	7.3
1948	3.7	3.8	2.7	2.7	4.1	4.7	3.7	3.7	2.5	2.5	3.0	4.1	6.4

DIPHTHERIA

Year	Annual	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
*1949	18.0	23.8	25.7	23.7	20.2	17.2	14.1	11.3	8.7	9.5	16.4	21.3	23.2
1948	20.3	27.4	26.0	25.9	22.8	19.7	15.1	9.6	9.7	13.0	22.3	25.5	27.2

DYSENTERY

Year	Annual	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
*1949	29.2	2.2	2.2	3.5	4.0	8.8	20.7	63.0	113.7	86.6	38.7	9.6	5.5
1948	18.3	1.6	2.5	4.8	4.7	7.8	17.8	50.2	64.0	41.5	16.0	5.6	2.9

TABLE 30. - 1/CASE RATES FOR SELECTED COMMUNICABLE DISEASES BY MONTH: JAPAN, 1946-1949, Cont'd
(Rates per 100,000 population)

SMALLPOX

Year	Annual	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
* 1949	0.2	0.0	0.0	0.2	0.3	0.9	0.3	0.1	-	-	-	0.0	0.0
1948	0.0	0.0	0.0	0.0	0.1	0.1	0.0	0.1	-	0.0	0.1	0.0	0.0

TYPHUS FEVER

Year	Annual	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
* 1949	0.1	0.4	0.4	0.1	0.2	0.1	0.0	0.1	-	0.0	0.1	0.1	0.2
1948	0.6	1.0	0.9	0.6	2.2	0.5	0.9	0.3	0.0	0.0	0.0	0.2	0.6

MALARIA

Year	Annual	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
* 1949	4.5	1.5	1.4	1.7	2.7	6.6	10.4	11.6	8.2	6.1	2.7	1.4	1.1
1948	6.2	3.4	3.5	3.8	4.8	5.6	10.8	13.1	12.8	8.4	4.2	1.9	1.8

JAPANESE "B" ENCEPHALITIS

Year	Annual	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
* 1949	1.6	0.0	-	-	-	-	0.0	0.2	1.1	11.6	5.6	0.7	0.0
1948	9.0	-	-	-	-	-	0.1	0.3	80.0	34.2	2.3	-	-

TABLE 30. - 1/CASE RATES FOR SELECTED COMMUNICABLE DISEASES BY MONTH: JAPAN, 1948-1949, Cont'd

Footnotes:

* Data for 1949 are provisional.

1/Rates are the number of cases per 100,000 civilian population in Japan, and are computed on an annual basis by Public Health and Welfare Section, GHQ, SCAP. Case data are from Weekly Reports, Ministry of Welfare.

"NA" indicates that data are not available.

A dash (-) indicates that no cases were reported.

A rate of 0.0 indicates that there were some cases but the rates was less than 0.1.

This Table corresponds to Table 26, Statistical Annex, Public Health and Welfare in Japan, 1948.

TABLE 31.- 1/ CASES OF SELECTED COMMUNICABLE DISEASES BY PREFECTURE, JAPAN, 1920 - 1949
TYPHOID FEVER

Year	All Japan	Hokkaido	Aomori	Iwate	Miyagi	Akita	Yamagata	Fukushima	Ibaraki	Tochigi	Gumma
•1949	6,489	192	87	83	210	55	60	114	86	67	57
1948	9,426	393	135	87	203	89	75	181	156	137	127
1947	17,820	756	238	230	395	156	350	443	421	420	278
1946	44,421	2,635	812	729	930	453	964	1,851	1,041	1,556	698
1945	57,933	4,494	896	1,035	784	747	922	1,561	822	2,200	733
1944	57,448	3,440	664	504	987	570	866	1,017	557	1,023	386
1943	52,519	2,188	582	852	1,374	266	528	899	460	915	386
1942	35,589	1,364	600	703	1,164	392	375	727	356	505	286
1941	40,595	1,266	513	1,016	1,007	1,052	607	927	561	503	426
1940	40,706	1,470	681	1,148	1,237	1,135	1,088	810	675	607	328
1939	37,837	1,548	546	1,728	1,183	1,029	882	1,111	510	491	336
1938	42,074	1,392	530	1,378	2,015	726	856	1,234	598	694	739
1937	38,124	1,285	484	1,022	1,160	937	572	977	568	495	485
1936	36,938	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1935	38,357	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1934	42,420	1,893	628	1,705	2,030	937	704	1,092	660	434	340
1933	38,408	2,209	559	743	1,602	669	760	794	898	401	425
1932	35,437	1,770	451	792	1,455	701	741	930	638	580	585
1931	38,202	1,810	466	652	1,341	520	809	809	471	767	630
1930	41,365	2,541	946	840	1,440	776	887	1,125	486	1,036	744
1929	37,262	1,856	713	522	1,143	417	691	1,193	462	688	517
1928	41,990	2,476	588	864	1,027	476	783	1,087	739	825	602
1927	37,499	2,275	896	823	717	596	934	1,176	811	740	775
1926	43,899	2,025	1,069	533	881	823	836	1,109	737	505	541
1925	45,762	3,418	754	702	1,440	989	528	1,475	769	622	628
1924	58,344	2,431	593	754	1,549	643	632	1,767	1,030	859	596
1923	52,584	2,108	731	840	1,383	961	992	1,917	1,195	1,028	688
1922	52,339	2,610	1,285	1,274	1,113	1,793	1,296	2,479	1,061	824	801
1921	50,094	1,710	722	1,226	1,489	960	1,252	2,277	1,259	910	914
1920	53,779	3,218	747	1,227	2,120	1,378	1,328	2,212	942	962	790

See footnotes at end of table.

TABLE 31.- 1/ CASES OF SELECTED COMMUNICABLE DISEASES BY PREFECTURE: JAPAN, 1920 - 1949 - Cont'd
TYPHOID FEVER - Cont'd

Year	Saitama	Chiba	Tokyo	Kanagawa	Niigata	Yoyama	Ishikawa	Fukui	Yamanashi	Nagano	Gifu	Shizuoka
*1949	199	155	896	272	309	91	46	136	11	116	210	249
1948	245	242	1,257	455	276	152	73	125	22	447	380	326
1947	498	407	1,386	665	619	388	194	162	135	332	600	646
1946	1,244	1,323	2,249	1,617	1,409	885	301	309	329	867	1,100	1,522
1945	1,700	1,383	2,540	2,851	2,597	1,039	439	353	928	1,683	2,206	2,515
1944	952	1,104	6,674	4,288	2,016	924	472	364	373	804	914	1,590
1943	1,056	964	6,904	4,012	1,348	837	632	457	199	874	1,005	1,498
1942	579	592	3,131	2,407	1,761	539	915	370	280	504	405	1,123
1941	761	596	3,421	2,023	1,314	512	403	455	537	499	474	1,218
1940	915	706	3,104	1,569	1,704	623	517	326	273	599	421	957
1939	431	647	2,356	1,322	1,446	491	399	127	266	536	413	1,528
1938	594	737	2,634	992	1,511	740	482	145	234	780	571	1,298
1937	606	641	2,301	982	966	438	291	243	452	526	833	949
1936	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1935	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1934	633	424	2,710	837	1,469	367	366	168	259	470	596	998
1933	513	361	3,273	974	1,239	251	279	223	356	585	461	915
1932	464	396	3,203	845	1,342	155	296	201	334	593	378	738
1931	578	332	4,194	1,124	1,183	256	422	224	271	675	427	878
1930	614	609	3,839	1,018	1,657	265	465	116	421	805	468	937
1929	402	406	3,942	665	739	311	193	113	312	1,021	480	1,027
1928	524	545	4,159	800	856	237	228	122	799	767	504	888
1927	424	336	3,986	1,107	980	156	237	321	400	804	472	1,016
1926	561	394	5,257	865	975	209	292	252	244	1,013	639	1,553
1925	506	579	5,681	1,000	975	368	285	115	494	815	508	1,248
1924	816	752	9,657	3,309	1,410	247	243	156	644	920	503	2,375
1923	865	798	7,741	3,237	1,057	372	364	305	393	1,174	678	1,774
1922	994	797	5,439	1,376	2,062	496	545	160	811	1,230	624	2,873
1921	1,145	453	5,120	2,239	1,490	203	395	228	813	1,283	717	2,562
1920	1,723	656	5,218	2,064	977	138	275	189	1,457	1,530	590	2,296

See footnotes at end of table.

TABLE 31.- 1/ CASES OF SELECTED COMMUNICABLE DISEASES BY PREFECTURE: JAPAN, 1920 - 1949 - Cont'd
TYPHOID FEVER - Cont'd

Year	Aichi	Mie	Shiga	Kyoto	Osaka	Hyogo	Nara	Wakayama	Tottori	Shimane	Okayama	Hiroshima
*1949	332	262	31	164	253	298	87	91	66	79	116	288
1948	438	296	53	271	353	382	119	123	93	122	179	285
1947	975	776	127	403	579	994	137	465	154	279	353	721
1946	1,282	1,138	341	1,370	1,328	2,547	509	823	369	736	1,224	1,683
1945	2,011	920	449	1,877	1,100	6,261	742	829	365	597	1,030	2,043
1944	2,072	933	337	1,465	4,043	7,501	496	636	253	431	698	2,409
1943	2,743	650	358	1,274	4,450	4,442	462	636	293	370	776	2,245
1942	1,541	421	285	709	2,501	3,189	330	650	226	377	814	1,567
1941	1,502	571	360	931	3,224	3,695	347	605	467	409	870	1,757
1940	1,473	588	369	874	3,345	3,179	271	511	294	272	811	1,901
1939	1,445	507	319	1,391	3,448	2,573	513	536	171	316	599	1,513
1938	1,547	540	302	1,261	4,075	3,133	406	638	342	223	734	1,777
1937	1,555	506	372	1,025	3,433	3,400	338	755	250	282	598	2,458
1936	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1935	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1934	2,219	599	440	1,637	4,660	2,624	288	457	324	235	684	1,566
1933	2,075	636	492	1,299	3,700	2,594	392	706	171	223	554	1,158
1932	1,896	448	328	996	3,093	2,002	231	467	146	368	766	1,045
1931	2,038	480	282	1,324	2,952	1,827	251	444	259	222	383	1,579
1930	1,740	508	303	982	2,579	1,993	231	413	207	793	589	1,360
1929	2,383	444	275	1,212	2,283	1,957	222	450	234	632	686	1,602
1928	2,283	578	369	1,388	3,214	2,468	388	505	371	369	1,330	597
1927	1,719	231	206	1,171	2,411	1,866	250	465	354	310	559	1,111
1926	1,765	526	285	1,578	2,398	1,878	237	445	434	422	816	823
1925	1,968	717	327	1,813	2,404	2,474	261	529	424	414	524	1,575
1924	2,083	610	244	2,740	3,526	2,459	339	417	550	448	561	2,903
1923	1,968	554	312	2,676	2,233	2,115	321	307	457	406	729	1,820
1922	1,639	742	165	2,125	2,876	2,056	258	294	479	406	898	1,123
1921	1,902	522	212	2,103	1,994	1,739	477	387	268	489	1,059	1,239
1920	1,861	549	207	1,780	1,970	2,371	522	312	363	742	1,322	2,352

See footnotes at end of table.

TABLE 31.- 1/ CASES OF SELECTED COMMUNICABLE DISEASES BY PREFECTURE: JAPAN, 1920 - 1949 - Cont'd
TYPHOID FEVER - Cont'd

Year	Yamaguchi	Tokushima	Kagawa	Ehime	Kochi	Fukuoka	Saga	Nagasaki	Kumamoto	Oita	Miyazaki	Kagoshima	Unknown
*1949	51	66	42	58	100	174	44	60	26	33	34	13	-
1948	89	127	53	165	159	195	55	69	25	116	56	20	-
1947	116	255	198	202	416	356	82	110	101	107	166	29	-
1946	380	794	602	519	774	1,301	275	446	230	237	493	196	-
1945	438	559	627	320	751	1,509	283	244	131	251	148	20	-
1944	463	774	609	352	938	1,305	261	315	101	305	169	93	-
1943	445	475	458	509	645	1,423	233	446	89	342	229	290	-
1942	300	440	475	511	550	1,084	238	174	91	323	142	163	-
1941	368	462	822	547	552	1,279	534	513	165	237	172	115	-
1940	386	490	655	935	563	1,268	319	576	191	234	219	89	-
1939	255	440	486	719	627	1,364	192	590	135	230	61	111	-
1938	370	451	469	854	775	1,943	293	395	251	208	99	108	-
1937	422	343	523	799	509	1,618	360	426	422	241	126	150	-
1936	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1935	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1934	327	376	704	939	382	1,985	255	452	402	552	227	376	-
1933	348	190	428	901	418	1,755	328	416	354	387	165	228	-
1932	360	308	420	577	396	1,599	547	723	372	388	91	282	-
1931	447	293	576	579	399	1,556	147	2,047	717	361	114	285	-
1930	403	375	585	565	438	2,162	273	805	903	357	185	581	-
1929	449	328	728	604	390	2,091	182	570	861	343	186	337	-
1928	458	260	665	707	538	1,995	244	921	1,573	362	125	272	-
1927	502	158	472	458	372	2,120	211	730	1,130	326	104	174	-
1926	708	149	457	473	787	3,695	104	1,134	2,663	355	122	398	-
1925	821	306	523	641	435	2,566	214	948	1,004	399	151	425	-
1924	1,575	463	740	541	326	2,568	264	755	1,071	479	138	658	-
1923	1,670	191	492	426	467	2,041	184	736	1,088	1,249	231	310	- 1,000
1922	781	132	746	448	458	2,190	107	916	715	327	187	276	52
1921	602	114	806	519	548	2,747	73	960	1,019	350	144	276	178
1920	614	172	645	466	486	2,628	76	536	911	357	138	339	23

See footnotes at end of table.

TABLE 31.- 1/ CASES OF SELECTED COMMUNICABLE DISEASES BY PREFECTURE; JAPAN, 1920 - 1949 - Cont'd
PARATYPHOID FEVER

Year	All Japan	Hokkaido	Aomori	Iwate	Miyagi	Akita	Yamagata	Fukushima	Ibaraki	Tochigi	Gumma
1949	2,226	82	40	33	116	16	21	136	48	23	31
1948	2,892	122	52	54	137	10	28	78	84	51	56
1947	4,735	230	54	65	260	46	106	102	192	103	121
1946	9,090	916	173	114	272	49	435	313	284	287	111
1945	10,269	1,584	65	148	383	146	172	266	298	322	290
1944	14,819	780	131	132	552	257	108	387	297	262	93
1943	12,382	459	94	180	899	109	152	289	241	229	103
1942	6,218	350	149	111	344	99	103	145	121	369	94
1941	6,233	256	69	106	201	81	175	220	608	98	130
1940	6,251	289	160	201	90	147	246	180	171	101	91
1939	5,227	242	80	137	64	85	289	127	147	88	48
1938	6,100	297	83	99	57	83	171	161	195	110	152
1937	4,439	237	92	168	98	93	251	110	52	94	53
1936	4,775	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1935	4,207	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1934	4,462	293	46	103	140	155	148	277	104	125	68
1933	5,279	795	42	77	38	53	156	129	388	144	126
1932	4,694	798	87	110	22	45	327	96	56	139	163
1931	4,042	379	48	25	51	28	279	86	66	133	128
1930	4,467	436	73	54	42	43	419	84	85	151	246
1929	4,189	307	78	53	96	30	201	201	127	153	158
1928	4,887	511	45	82	56	49	190	114	92	213	165
1927	4,735	580	52	61	69	44	70	137	189	119	250
1926	4,422	475	37	74	69	77	87	128	90	124	162
1925	5,050	727	49	114	76	57	60	135	164	159	196
1924	5,330	430	38	68	57	51	68	135	140	182	329
1923	5,288	396	96	155	137	84	87	114	115	160	327
1922	7,111	517	172	138	69	153	99	229	187	196	211
1921	6,287	365	91	114	98	116	195	193	87	131	274
1920	7,697	708	90	133	111	102	106	254	106	226	711

See footnotes at end of table.

TABLE 31.- 1/ CASES OF SELECTED COMMUNICABLE DISEASES BY PREFECTURE: JAPAN, 1920 - 1949 - Cont'd

Year	Saitama	Chiba	Tokyo	Kanagawa	Niigata	Toyama	Ishikawa	Fukui	Yamanashi	Nagano	Gifu	Shizuoka
*1949	55	44	488	68	80	43	31	30	5	50	27	103
1948	44	59	681	117	110	28	17	22	12	70	81	165
1947	98	133	487	162	189	114	50	42	49	137	138	160
1946	185	179	689	306	332	100	67	59	100	342	199	467
1945	289	153	989	281	414	156	129	48	316	752	108	422
1944	230	143	5,506	628	290	344	65	73	59	290	183	358
1943	154	167	3,408	547	406	104	59	62	32	347	172	352
1942	191	45	892	218	344	125	42	121	25	175	34	100
1941	137	31	860	244	193	35	34	65	89	267	56	194
1940	301	69	675	247	313	7	40	43	99	329	75	164
1939	96	60	436	179	300	20	60	36	92	193	158	383
1938	143	61	478	108	238	5	143	59	91	360	63	204
1937	100	64	326	155	124	12	34	35	20	172	136	159
1936	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1935	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1934	125	25	564	114	252	19	11	8	33	128	29	121
1933	91	71	505	88	294	16	23	9	84	106	58	213
1932	76	33	606	74	185	7	17	15	25	99	21	130
1931	72	32	593	79	197	19	22	13	7	117	18	133
1930	132	39	478	120	189	22	18	8	26	146	31	158
1929	164	38	380	102	164	29	28	12	41	139	15	238
1928	170	66	415	270	175	30	24	17	22	150	39	159
1927	117	52	403	134	186	14	33	18	20	271	52	215
1926	104	48	512	94	150	11	150	16	31	175	34	148
1925	246	106	434	95	219	25	26	12	11	101	21	115
1924	130	99	535	145	239	35	16	33	31	162	35	255
1923	188	103	496	187	243	50	27	51	43	134	57	254
1922	355	131	390	199	586	129	61	71	99	137	77	256
1921	203	93	400	233	431	9	31	45	112	159	38	466
1920	367	97	428	272	347	56	51	17	115	140	105	604

See footnotes at end of table.

TABLE 31.- 1/ CASES OF SELECTED COMMUNICABLE DISEASES BY PREFECTURE: JAPAN, 1920 - 1949 - Cont'd

Year	Aichi	Mie	Shiga	Kyoto	Osaka	Hyogo	Nara	Wakayama	Tottori	Shimane	Okayama	Hiroshima
1949	89	54	19	58	51	32	12	33	15	16	21	81
1948	93	63	15	67	67	36	14	16	13	66	16	64
1947	198	130	29	97	276	109	16	65	33	114	21	165
1946	247	136	31	171	206	278	57	89	81	145	38	243
1945	193	49	60	121	99	645	79	64	49	130	35	211
1944	345	65	47	85	553	813	34	57	48	-	37	459
1943	373	72	40	219	740	668	15	46	36	47	50	403
1942	260	46	28	106	200	178	11	92	53	48	74	229
1941	210	84	57	143	223	353	32	59	26	55	37	165
1940	194	58	28	200	302	231	32	48	21	44	69	144
1939	194	57	17	115	203	162	9	55	9	52	31	198
1938	216	72	21	150	211	285	7	49	26	19	15	175
1937	200	43	34	61	335	193	10	32	14	43	39	294
1936	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1935	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1934	146	68	28	85	272	173	8	36	24	63	24	36
1933	264	30	23	111	325	282	9	69	53	15	19	41
1932	146	31	44	133	255	149	4	63	10	40	22	69
1931	162	27	34	187	241	136	18	55	38	53	19	43
1930	123	45	40	217	206	177	10	88	13	44	30	54
1929	134	29	21	117	202	209	7	97	16	50	20	53
1928	142	42	25	131	286	279	9	45	35	83	46	47
1927	138	44	22	147	258	283	28	71	69	75	39	69
1926	138	70	20	231	291	181	34	44	46	30	67	48
1925	128	64	11	414	351	275	36	19	24	30	35	48
1924	109	32	24	272	470	269	31	66	24	56	48	37
1923	107	29	31	268	309	233	53	35	27	70	38	41
1922	143	87	32	452	457	522	92	51	26	106	34	49
1921	164	94	29	343	411	314	154	49	18	75	55	52
1920	253	97	50	312	383	410	124	73	10	56	86	106

See footnotes at end of table.

TABLE 31.- 1/ CASES OF SELECTED COMMUNICABLE DISEASES BY PREFECTURE: JAPAN, 1920 - 1949 - Cont'd
PARATYPHOID FEVER - Cont'd

Year	Yamaguchi	Tokushima	Kagawa	Ehime	Kochi	Fukuoka	Saga	Nagasaki	Kumamoto	Oita	Miyazaki	Kagoshima	Unknown
1949	10	13	5	17	8	46	6	6	33	10	15	6	-
1948	12	8	33	33	35	56	17	19	21	23	19	8	-
1947	31	38	65	34	43	63	29	39	24	12	48	18	-
1946	88	51	213	59	67	341	122	192	48	53	116	59	-
1945	48	81	46	105	52	323	20	31	-	31	112	14	-
1944	94	90	50	142	23	194	64	49	13	58	30	301	-
1943	62	130	56	46	36	343	17	49	24	79	51	215	-
1942	65	85	37	48	51	226	3	41	27	32	15	16	-
1941	29	39	27	52	86	195	13	36	12	73	46	32	-
1940	43	49	26	29	43	197	16	354	41	15	9	20	-
1939	26	32	23	32	44	443	12	144	24	67	11	7	-
1938	23	47	22	21	26	998	27	78	46	180	8	17	-
1937	42	27	21	20	40	210	24	23	72	42	17	18	-
1936	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1935	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1934	67	177	21	40	19	101	27	24	36	14	23	62	-
1933	45	52	23	49	16	162	11	101	40	11	4	10	-
1932	128	37	25	35	7	141	18	28	152	20	4	2	-
1931	42	20	19	26	13	234	4	28	76	13	7	20	-
1930	56	24	19	17	16	123	27	34	57	27	18	2	-
1929	75	37	49	16	9	120	21	37	51	47	13	5	-
1928	71	67	41	44	21	136	34	31	121	39	39	19	-
1927	26	43	64	50	23	68	20	38	46	15	10	3	-
1926	36	24	39	25	11	119	20	38	37	15	13	3	-
1925	41	64	31	12	20	123	42	63	30	29	5	7	-
1924	67	114	33	17	29	127	61	55	74	88	8	6	-
1923	68	76	24	17	16	103	32	64	89	36	7	11	-
1922	40	62	40	29	14	136	33	86	75	48	16	16	3
1921	43	69	22	20	18	212	5	54	134	22	29	16	1
1920	49	41	12	22	42	173	11	63	77	33	33	35	-

See footnotes at end of table.

TABLE 31.-- 1/ CASES OF SELECTED COMMUNICABLE DISEASES BY PREFECTURE: JAPAN, 1920 - 1949 - Cont'd
PLAGUE

Year	All Japan	Hokkaido	Aomori	Iwate	Miyagi	Akita	Yamagata	Fukushima	Ibaraki	Tochigi	Guma
*1949	-	-	-	-	-	-	-	-	-	-	-
1948	-	-	-	-	-	-	-	-	-	-	-
1947	-	-	-	-	-	-	-	-	-	-	-
1946	-	-	-	-	-	-	-	-	-	-	-
1945	-	-	-	-	-	-	-	-	-	-	-
1944	-	-	-	-	-	-	-	-	-	-	-
1943	-	-	-	-	-	-	-	-	-	-	-
1942	-	-	-	-	-	-	-	-	-	-	-
1941	-	-	-	-	-	-	-	-	-	-	-
1940	-	-	-	-	-	-	-	-	-	-	-
1939	-	-	-	-	-	-	-	-	-	-	-
1938	-	-	-	-	-	-	-	-	-	-	-
1937	-	-	-	-	-	-	-	-	-	-	-
1936	-	-	-	-	-	-	-	-	-	-	-
1935	-	-	-	-	-	-	-	-	-	-	-
1934	-	-	-	-	-	-	-	-	-	-	-
1933	-	-	-	-	-	-	-	-	-	-	-
1932	-	-	-	-	-	-	-	-	-	-	-
1931	-	-	-	-	-	-	-	-	-	-	-
1930	-	-	-	-	-	-	-	-	-	-	-
1929	-	-	-	-	-	-	-	-	-	-	-
1928	-	-	-	-	-	-	-	-	-	-	-
1927	-	-	-	-	-	-	-	-	-	-	-
1926	3	-	-	-	-	-	-	-	-	-	-
1925	-	-	-	-	-	-	-	-	-	-	-
1924	7	-	-	-	-	-	-	-	-	-	-
1923	1	-	-	-	-	-	-	-	-	-	-
1922	118	-	-	-	-	-	-	-	-	-	-
1921	2	-	-	-	-	-	-	-	-	-	-
1920	22	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

TABLE 31... 1/ CASES OF SELECTED COMMUNICABLE DISEASES BY PREFECTURE, JAPAN, 1920 - 1949 - Cont'd
 FLAGUE - Cont'd

Year	Saitama	Chiba	Tokyo	Kanagawa	Niigata	Toyama	Ishikawa	Fukui	Yamanashi	Nagano	Gifu	Shizuoka
*1949	-	-	-	-	-	-	-	-	-	-	-	-
1948	-	-	-	-	-	-	-	-	-	-	-	-
1947	-	-	-	-	-	-	-	-	-	-	-	-
1946	-	-	-	-	-	-	-	-	-	-	-	-
1945	-	-	-	-	-	-	-	-	-	-	-	-
1944	-	-	-	-	-	-	-	-	-	-	-	-
1943	-	-	-	-	-	-	-	-	-	-	-	-
1942	-	-	-	-	-	-	-	-	-	-	-	-
1941	-	-	-	-	-	-	-	-	-	-	-	-
1940	-	-	-	-	-	-	-	-	-	-	-	-
1939	-	-	-	-	-	-	-	-	-	-	-	-
1938	-	-	-	-	-	-	-	-	-	-	-	-
1937	-	-	-	-	-	-	-	-	-	-	-	-
1936	-	-	-	-	-	-	-	-	-	-	-	-
1935	-	-	-	-	-	-	-	-	-	-	-	-
1934	-	-	-	-	-	-	-	-	-	-	-	-
1933	-	-	-	-	-	-	-	-	-	-	-	-
1932	-	-	-	-	-	-	-	-	-	-	-	-
1931	-	-	-	-	-	-	-	-	-	-	-	-
1930	-	-	-	-	-	-	-	-	-	-	-	-
1929	-	-	-	-	-	-	-	-	-	-	-	-
1928	-	-	-	-	-	-	-	-	-	-	-	-
1927	-	-	-	-	-	-	-	-	-	-	-	-
1926	-	-	-	8	-	-	-	-	-	-	-	-
1925	-	-	-	-	-	-	-	-	-	-	-	-
1924	-	-	-	-	-	-	-	-	-	-	-	4
1923	-	-	-	-	-	-	-	-	-	-	-	-
1922	-	-	-	-	-	-	-	-	-	-	-	-
1921	-	-	-	-	-	-	-	-	-	-	-	-
1920	-	-	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table

TABLE 31.- 1/ CASES OF SELECTED COMMUNICABLE DISEASES BY PREFECTURE: JAPAN, 1920 - 1949 - Cont'd

PLAGUE - Cont'd

Year	Aichi	Mie	Shiga	Kyoto	Osaka	Hyogo	Nara	Wakayama	Tottori	Shimane	Okayama	Hiroshima
1949	-	-	-	-	-	-	-	-	-	-	-	-
1948	-	-	-	-	-	-	-	-	-	-	-	-
1947	-	-	-	-	-	-	-	-	-	-	-	-
1946	-	-	-	-	-	-	-	-	-	-	-	-
1945	-	-	-	-	-	-	-	-	-	-	-	-
1944	-	-	-	-	-	-	-	-	-	-	-	-
1943	-	-	-	-	-	-	-	-	-	-	-	-
1942	-	-	-	-	-	-	-	-	-	-	-	-
1941	-	-	-	-	-	-	-	-	-	-	-	-
1940	-	-	-	-	-	-	-	-	-	-	-	-
1939	-	-	-	-	-	-	-	-	-	-	-	-
1938	-	-	-	-	-	-	-	-	-	-	-	-
1937	-	-	-	-	-	-	-	-	-	-	-	-
1936	-	-	-	-	-	-	-	-	-	-	-	-
1935	-	-	-	-	-	-	-	-	-	-	-	-
1934	-	-	-	-	-	-	-	-	-	-	-	-
1933	-	-	-	-	-	-	-	-	-	-	-	-
1932	-	-	-	-	-	-	-	-	-	-	-	-
1931	-	-	-	-	-	-	-	-	-	-	-	-
1930	-	-	-	-	-	-	-	-	-	-	-	-
1929	-	-	-	-	-	-	-	-	-	-	-	-
1928	-	-	-	-	-	-	-	-	-	-	-	-
1927	-	-	-	-	-	-	-	-	-	-	-	-
1926	-	-	-	-	-	-	-	-	-	-	-	-
1925	-	-	-	-	-	-	-	-	-	-	-	-
1924	-	-	-	-	-	3	-	-	-	-	-	-
1923	-	-	-	-	-	-	-	-	-	-	-	-
1922	-	-	-	-	81	-	2	-	-	-	-	-
1921	-	-	-	-	-	2	-	-	-	-	-	-
1920	-	-	-	-	-	-	-	22	-	-	-	-

See footnotes at end of table.

TABLE 31.- 1/ CASES OF SELECTED COMMUNICABLE DISEASES BY PREFECTURE: JAPAN, 1920 - 1949 Cont'd

PLAGUE - Cont'd

Year	Yamaguchi	Tokushima	Kagawa	Khine	Kochi	Fukuoka	Saga	Nagasaki	Kumamoto	Oita	Miyazaki	Kagoshima ^{2/}	Unknown
*1949	-	-	-	-	-	-	-	-	-	-	-	-	-
1948	-	-	-	-	-	-	-	-	-	-	-	-	-
1947	-	-	-	-	-	-	-	-	-	-	-	-	-
1946	-	-	-	-	-	-	-	-	-	-	-	-	-
1945	-	-	-	-	-	-	-	-	-	-	-	-	-
1944	-	-	-	-	-	-	-	-	-	-	-	-	-
1943	-	-	-	-	-	-	-	-	-	-	-	-	-
1942	-	-	-	-	-	-	-	-	-	-	-	-	-
1941	-	-	-	-	-	-	-	-	-	-	-	-	-
1940	-	-	-	-	-	-	-	-	-	-	-	-	-
1939	-	-	-	-	-	-	-	-	-	-	-	-	-
1938	-	-	-	-	-	-	-	-	-	-	-	-	-
1937	-	-	-	-	-	-	-	-	-	-	-	-	-
1936	-	-	-	-	-	-	-	-	-	-	-	-	-
1935	-	-	-	-	-	-	-	-	-	-	-	-	-
1934	-	-	-	-	-	-	-	-	-	-	-	-	-
1933	-	-	-	-	-	-	-	-	-	-	-	-	-
1932	-	-	-	-	-	-	-	-	-	-	-	-	-
1931	-	-	-	-	-	-	-	-	-	-	-	-	-
1930	-	-	-	-	-	-	-	-	-	-	-	-	-
1929	-	-	-	-	-	-	-	-	-	-	-	-	-
1928	-	-	-	-	-	-	-	-	-	-	-	-	-
1927	-	-	-	-	-	-	-	-	-	-	-	-	-
1926	-	-	-	-	-	-	-	-	-	-	-	-	-
1925	-	-	-	-	-	-	-	-	-	-	-	-	-
1924	-	-	-	-	-	-	-	-	-	-	-	-	-
1923	-	-	-	-	1	-	-	-	-	-	-	-	-
1922	-	-	-	-	35	-	-	-	-	-	-	-	-
1921	-	-	-	-	-	-	-	-	-	-	-	-	-
1920	-	-	-	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

TABLE 31.- 1/ CASES OF SELECTED COMMUNICABLE DISEASES BY PREFECTURE; JAPAN, 1920 - 1949 - Cont'd
CHOLERA

Year	All Japan	Hokkaido	Aomori	Iwate	Miyagi	Akita	Yamagata	Fukushima	Ibaraki	Tochigi	Gumma
1949	-	-	-	-	-	-	-	-	-	-	-
1938	-	-	-	-	-	-	-	-	-	-	-
1947	-	-	-	-	-	-	-	-	-	-	-
1946	1,229	-	-	-	-	-	-	6	-	-	-
1945	-	-	-	-	-	-	-	-	-	-	-
1944	-	-	-	-	-	-	-	-	-	-	-
1943	-	-	-	-	-	-	-	-	-	-	-
1942	-	-	-	-	-	-	-	-	-	-	-
1941	-	-	-	-	-	-	-	-	-	-	-
1940	-	-	-	-	-	-	-	-	-	-	-
1939	-	-	-	-	-	-	-	-	-	-	-
1938	18	-	-	-	-	-	-	-	-	-	-
1937	57	-	-	-	-	-	-	-	-	-	-
1936	-	-	-	-	-	-	-	-	-	-	-
1935	-	-	-	-	-	-	-	-	-	-	-
1934	-	-	-	-	-	-	-	-	-	-	-
1933	-	-	-	-	-	-	-	-	-	-	-
1932	4	-	-	-	-	-	-	-	-	-	-
1931	-	-	-	-	-	-	-	-	-	-	-
1930	-	-	-	-	-	-	-	-	-	-	-
1929	205	-	-	-	-	-	-	-	-	-	-
1928	1	-	-	-	-	-	-	-	-	-	-
1927	2	-	-	-	-	-	-	-	-	-	-
1926	25	-	-	-	-	-	-	-	-	-	-
1925	624	-	-	-	-	-	-	-	-	1	-
1924	-	-	-	-	-	-	-	-	-	-	-
1923	4	-	-	-	-	-	-	-	-	-	-
1922	749	-	-	-	-	-	-	-	-	-	-
1921	29	-	34	10	1	-	-	4	79	-	2
1920	4,985	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

TABLE 31.- 1/ CASES OF SELECTED COMMUNICABLE DISEASES BY PREFECTURE: JAPAN, 1920 - 1949 - Cont'd

CHOLERA - Cont'd

Year	Saitama	Chiba	Tokyo	Kenagawa	Niigata	Toyama	Ishikawa	Fukui	Yamanashi	Nagano	Gifu	Shizuoka
*1949	-	-	-	-	-	-	-	-	-	-	-	-
1948	-	-	-	-	-	-	-	-	-	-	-	-
1947	-	-	-	-	-	-	-	-	-	-	-	-
1946	1	33	12	29	27	77	22	14	-	2	-	2
1945	-	-	-	-	-	-	-	-	-	-	-	-
1944	-	-	-	-	-	-	-	-	-	-	-	-
1943	-	-	-	-	-	-	-	-	-	-	-	-
1942	-	-	-	-	-	-	-	-	-	-	-	-
1941	-	-	-	-	-	-	-	-	-	-	-	-
1940	-	-	-	-	-	-	-	-	-	-	-	-
1939	-	-	-	-	-	-	-	-	-	-	-	-
1938	-	-	-	-	-	-	-	-	-	-	-	-
1937	-	-	1	-	-	-	-	-	-	-	-	-
1936	-	-	-	-	-	-	-	-	-	-	-	-
1935	-	-	-	-	-	-	-	-	-	-	-	-
1934	-	-	-	-	-	-	-	-	-	-	-	-
1933	-	-	-	-	-	-	-	-	-	-	-	-
1932	-	-	3	-	-	-	-	-	-	-	-	-
1931	-	-	-	-	-	-	-	-	-	-	-	-
1930	-	-	-	-	-	-	-	-	-	-	-	-
1929	-	-	-	-	-	-	-	-	-	-	-	-
1928	-	-	-	-	-	-	-	-	-	-	-	-
1927	-	-	-	-	-	-	-	-	-	-	-	-
1926	-	-	-	1	-	-	-	-	-	-	-	-
1925	4	46	81	97	-	-	-	-	-	-	-	7
1924	-	-	-	-	-	-	-	-	-	-	-	-
1923	-	-	-	-	-	-	-	-	-	-	-	-
1922	1	168	184	19	-	-	-	-	-	-	-	10
1921	-	-	-	2	-	-	-	-	-	-	-	-
1920	-	-	3	3	-	221	-	-	-	-	-	-

See footnotes at end of table.

TABLE 31.- 1/ CASES OF SELECTED COMMUNICABLE DISEASES BY PREFECTURE: JAPAN, 1920 - 1949 - Cont'd
CHOLERA - Cont'd

Year	Aichi	Mie	Shiga	Kyoto	Osaka	Ryogo	Nara	Wakayama	Tottori	Shimane	Okayama	Hiroshima
*1949	-	-	-	-	-	-	-	-	-	-	-	-
1948	-	-	-	-	-	-	-	-	-	-	-	-
1947	-	-	-	-	-	-	-	-	-	-	-	-
1946	10	2	-	17	70	15	-	6	6	17	34	169
1945	-	-	-	-	-	-	-	-	-	-	-	-
1944	-	-	-	-	-	-	-	-	-	-	-	-
1943	-	-	-	-	-	-	-	-	-	-	-	-
1942	-	-	-	-	-	-	-	-	-	-	-	-
1941	-	-	-	-	-	-	-	-	-	-	-	-
1940	-	-	-	-	-	-	-	-	-	-	-	-
1939	-	-	-	-	-	-	-	-	-	-	-	-
1938	-	-	-	-	-	-	-	-	-	-	9	6
1937	-	-	-	-	-	1	-	-	-	-	5	40
1936	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1935	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1934	-	-	-	-	-	-	-	-	-	-	-	-
1933	-	-	-	-	-	-	-	-	-	-	-	-
1932	-	-	-	-	-	-	-	-	-	-	-	-
1931	-	-	-	-	-	-	-	-	-	-	-	-
1930	-	-	-	-	-	-	-	-	-	-	-	-
1929	-	4	-	-	72	14	21	5	1	-	26	45
1928	-	-	-	-	1	-	-	-	-	-	-	-
1927	-	-	-	-	1	-	-	-	-	-	1	-
1926	-	-	-	-	2	11	-	-	-	-	3	1
1925	-	19	-	4	53	124	-	19	2	-	20	50
1924	-	-	-	-	-	-	-	-	-	-	-	-
1923	-	-	-	-	3	-	-	-	-	-	-	-
1922	1	-	-	-	3	2	-	-	-	3	-	-
1921	-	-	-	-	-	-	-	-	-	-	-	-
1920	1	8	-	60	752	1,109	189	96	-	37	61	264

See footnotes at end of table.

TABLE 31.- 1/ CASES OF SELECTED COMMUNICABLE DISEASES BY PREFECTURE: JAPAN, 1920 - 1949 - Cont'd

CHOLERA - Cont'd

Year	Yamaguchi	Tokushima	Kagawa	Ehime	Kochi	Fukuoka	Saga	Nagasaki	Kumamoto	Oita	Miyazaki	Kagoshima	2/ Unknown
*1949	-	-	-	-	-	-	-	-	-	-	-	-	-
1948	-	-	-	-	-	-	-	-	-	-	-	-	-
1947	-	-	-	-	-	-	-	-	-	-	-	-	-
1946	90	-	1	17	-	185	88	164	25	6	17	65	-
1945	-	-	-	-	-	-	-	-	-	-	-	-	-
1944	-	-	-	-	-	-	-	-	-	-	-	-	-
1943	-	-	-	-	-	-	-	-	-	-	-	-	-
1942	-	-	-	-	-	-	-	-	-	-	-	-	-
1941	-	-	-	-	-	-	-	-	-	-	-	-	-
1940	-	-	-	-	-	-	-	-	-	-	-	-	-
1939	-	-	-	-	-	-	-	-	-	-	-	-	-
1938	-	-	-	-	-	2	-	1	-	-	-	-	-
1937	10	-	-	-	-	-	-	-	-	-	-	-	-
1936	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1935	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1934	-	-	-	-	-	-	-	-	-	-	-	-	-
1933	-	-	-	-	-	-	-	-	-	-	-	-	-
1932	1	-	-	-	-	-	-	-	-	-	-	-	-
1931	-	-	-	-	-	-	-	-	-	-	-	-	-
1930	-	-	-	-	-	-	-	-	-	-	-	-	-
1929	2	11	1	-	-	2	-	-	-	1	-	-	-
1928	-	-	-	-	-	-	-	-	-	-	-	-	-
1927	-	-	-	-	-	-	-	-	-	-	-	-	-
1926	-	-	7	-	-	-	-	-	-	-	-	-	-
1925	2	10	3	48	15	1	-	13	-	-	1	4	-
1924	-	-	-	-	-	-	-	-	-	-	-	-	-
1923	1	-	-	-	-	-	-	-	-	-	-	-	-
1922	5	-	-	-	-	172	22	22	1	-	-	-	-
1921	1	-	-	-	-	6	-	20	-	-	-	-	-
1920	165	129	37	185	90	499	122	363	272	198	67	38	16

See footnotes at end of table.

TABLE 31.- 1/ CASES OF SELECTED COMMUNICABLE DISEASES BY PREFECTURE: JAPAN, 1920 - 1949 - Cont'd

EPIDEMIC MENINGITIS

Year	All Japan	Hokkaido	Aomori	Iwate	Miyagi	Akita	Yamagata	Fukushima	Ibaraki	Tochigi	Gumma
1949	1,467	154	36	27	62	38	34	54	33	11	22
1948	2,035	187	63	18	111	73	44	83	74	12	32
1947	3,371	370	103	60	136	89	83	145	197	31	38
1946	1,468	232	73	41	73	40	45	69	38	13	19
1945	4,384	1,452	124	47	212	13	15	113	159	14	54
1944	1,468	169	50	10	52	5	12	5	69	4	21
1943	1,113	68	70	9	49	28	8	19	25	3	23
1942	823	118	60	12	40	29	12	15	18	5	7
1941	1,160	174	125	43	64	54	12	15	28	9	7
1940	1,350	235	220	36	31	92	16	14	37	2	8
1939	1,632	175	333	17	20	19	14	26	37	1	6
1938	995	133	38	6	7	9	6	27	4	-	5
1937	837	26	34	3	6	5	8	6	22	2	5
1936	1,002	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1935	1,304	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1934	1,186	17	4	4	5	3	1	2	4	1	4
1933	359	40	7	3	12	2	10	6	5	1	19
1932	238	5	7	2	3	3	4	3	4	1	6
1931	280	5	5	1	3	13	-	8	6	-	6
1930	275	4	-	1	7	3	3	2	1	-	6
1929	352	2	1	1	7	1	1	10	3	-	17
1928	294	2	1	1	6	5	1	10	3	-	5
1927	330	1	2	-	11	2	1	7	6	3	3
1926	408	3	-	2	10	-	-	2	10	4	3
1925	445	5	3	2	14	-	6	3	3	4	4
1924	1,347	13	2	35	26	36	7	10	9	7	3
1923	708	1	1	2	37	8	1	4	58	7	4
1922	934	4	1	7	36	16	1	11	3	8	4
1921	772	5	3	6	44	6	6	7	4	1	-
1920	946	16	2	4	6	12	1	14	6	5	-

See footnotes at end of table.

TABLE 31.- 1/ CASES OF SELECTED COMMUNICABLE DISEASES BY PREFECTURE; JAPAN, 1920 - 1949 - Cont'd
EPIDEMIC MENINGITIS - Cont'd

Year	Saitama	Chiba	Tokyo	Kanagawa	Niigata	Toyama	Ishikawa	Fukui	Yamaguchi	Nagano	Gifu	Shizuoka
1949	31	23	256	75	24	17	4	15	11	23	10	31
1948	23	39	417	108	51	27	11	16	20	40	13	53
1947	72	62	663	77	67	20	45	12	27	41	18	93
1946	12	29	140	37	28	4	-	2	20	11	12	38
1945	11	12	90	142	21	19	3	9	-	16	10	28
1944	8	13	109	69	9	8	-	-	-	3	5	18
1943	5	10	104	58	1	1	1	-	-	3	2	6
1942	6	8	157	35	2	2	-	-	-	3	1	6
1941	7	8	127	37	5	1	1	1	3	4	5	8
1940	3	13	116	24	2	-	4	1	2	3	1	12
1939	2	10	107	13	3	-	-	2	4	6	2	20
1938	2	7	121	20	2	1	2	1	-	1	1	11
1937	7	4	127	12	3	2	2	1	2	6	3	6
1936	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1935	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1934	-	2	84	5	-	2	3	1	3	4	8	4
1933	2	3	58	8	4	-	3	2	1	-	2	-
1932	1	2	49	7	1	1	1	-	5	5	-	4
1931	2	1	81	14	3	-	-	-	2	2	3	2
1930	3	3	74	9	1	2	1	-	2	2	-	5
1929	1	1	80	9	-	1	-	-	1	-	-	5
1928	5	2	67	8	2	2	1	-	2	3	-	3
1927	1	-	71	5	1	1	1	-	2	1	1	6
1926	-	6	71	10	-	2	1	-	6	4	1	7
1925	2	11	81	6	2	-	1	1	14	2	3	45
1924	4	10	169	7	5	1	1	-	7	9	1	26
1923	3	5	134	11	2	-	2	-	11	2	4	65
1922	2	5	119	16	4	-	-	-	5	11	2	12
1921	-	2	100	21	3	-	2	-	9	2	1	8
1920	-	5	93	17	2	-	-	-	9	10	3	1

See footnotes at end of table.

TABLE 31.- 1/ CASES OF SELECTED COMMUNICABLE DISEASES BY PREFECTURE; JAPAN, 1920 - 1949 - Cont'd
EPIDEMIC MENINGITIS - Cont'd

Year	Aichi	Mie	Shiga	Kyoto	Osaka	Hyogo	Nara	Wakayama	Tottori	Shimane	Okayama	Hiroshima
•1949	17	14	12	57	99	33	7	4	28	11	8	26
1948	24	11	8	63	90	40	9	7	24	16	9	22
1947	43	28	29	71	151	69	6	10	45	17	12	66
1946	16	22	9	44	47	22	3	2	15	8	7	18
1945	22	22	15	108	43	67	7	15	19	9	12	92
1944	14	1	5	16	64	31	2	4	10	1	2	46
1943	14	3	6	40	91	44	2	5	4	1	4	11
1942	11	4	2	32	68	26	2	3	5	1	-	10
1941	17	4	4	46	96	33	7	2	10	2	4	2
1940	13	4	8	36	116	25	18	4	6	-	6	6
1939	12	5	10	62	156	42	36	12	29	2	14	10
1938	6	3	7	52	122	44	5	15	3	1	7	11
1937	7	2	6	29	104	26	2	12	7	2	1	5
1936	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1935	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1934	7	2	9	35	135	20	-	9	-	-	3	20
1933	6	-	7	26	52	21	-	6	7	1	1	4
1932	3	1	3	11	64	7	-	4	1	1	2	7
1931	6	-	3	9	54	8	1	8	-	2	2	7
1930	7	2	3	18	61	9	1	4	-	3	-	3
1929	10	1	2	14	68	18	-	4	3	4	8	5
1928	6	2	2	15	64	10	2	4	-	5	3	8
1927	4	4	2	37	82	17	2	3	6	3	3	7
1926	3	3	2	19	109	22	2	-	5	3	6	11
1925	10	3	1	26	87	18	2	2	9	-	4	3
1924	17	4	-	33	93	34	6	9	44	24	381	140
1923	14	7	-	28	102	32	2	3	7	29	7	11
1922	10	5	-	56	105	50	8	3	21	27	95	76
1921	7	-	-	61	164	50	5	4	24	34	5	18
1920	8	6	-	54	262	43	3	8	16	9	7	18

See footnotes at end of table.

TABLE 31.- 1/ CASES OF SELECTED COMMUNICABLE DISEASES BY PREFECTURE: JAPAN, 1920 - 1949 - Cont'd
EPIDEMIC MENINGITIS - Cont'd

Year	Yamaguchi	Tokushima	Kagawa	Ehime	Kochi	Fukuoka	Saga	Nagasaki	Kumamoto	Oita	Miyazaki	Kagoshima	Unknown
1949	12	1	9	13	11	59	10	8	7	14	8	8	-
1948	19	5	9	20	5	44	7	27	19	10	7	25	-
1947	35	9	20	37	25	86	17	33	33	15	27	38	-
1946	32	17	9	6	7	95	11	39	14	5	25	19	-
1945	164	1	20	27	32	738	14	322	33	7	21	10	-
1944	17	2	1	6	4	129	5	429	21	-	5	14	-
1943	7	1	4	7	4	95	5	248	9	-	5	10	-
1942	10	-	3	7	10	72	3	13	5	-	-	1	-
1941	6	3	12	4	10	124	10	11	2	-	4	9	-
1940	6	-	2	9	6	174	4	17	3	4	6	5	-
1939	12	5	9	4	22	302	11	21	11	8	4	16	-
1938	6	8	8	6	10	246	1	16	8	1	1	5	-
1937	13	1	1	1	12	278	2	12	9	8	2	3	-
1936	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1935	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1934	3	1	5	1	3	744	-	10	8	-	3	7	-
1933	2	-	4	-	7	8	-	12	1	-	-	5	-
1932	2	-	-	-	1	6	-	5	3	-	1	2	-
1931	1	1	1	-	3	12	-	3	-	1	-	1	-
1930	1	2	4	-	6	6	-	5	9	-	2	-	-
1929	13	1	11	1	18	17	-	8	4	-	-	1	-
1928	2	2	2	2	8	16	-	3	3	-	3	-	-
1927	2	5	1	1	3	16	-	2	1	-	3	-	-
1926	16	4	-	-	7	22	-	3	10	-	2	17	-
1925	4	8	7	2	4	23	-	10	6	-	2	1	-
1924	25	5	4	7	31	55	1	26	8	1	-	9	-
1923	6	5	4	3	12	38	4	14	13	3	-	2	-
1922	7	46	25	14	25	59	5	14	7	-	-	7	-
1921	12	1	15	13	14	66	10	14	17	-	1	7	-
1920	13	9	7	15	17	176	-	38	12	-	-	2	-

See footnotes at end of table.

TABLE 31.- 1/ CASES OF SELECTED COMMUNICABLE DISEASES BY PREFECTURE, JAPAN, 1920 - 1949 - Cont'd
SCARLET FEVER

Year	All Japan	Hokkaido	Aomori	Iwate	Miyagi	Akita	Yamagata	Fukushima	Ibaraki	Tochigi	Gumma
*1949	4,667	413	36	58	68	47	63	66	62	58	81
1948	2,924	482	24	19	93	24	27	39	62	51	92
1947	2,635	366	28	28	104	31	42	47	64	42	81
1946	2,209	395	23	14	55	50	55	58	35	22	42
1945	2,405	359	26	25	34	39	34	28	30	19	59
1944	6,354	720	82	54	94	42	87	84	56	50	133
1943	9,891	869	88	115	173	77	78	88	82	38	133
1942	12,688	1,164	150	199	216	86	65	138	150	69	156
1941	14,997	1,201	138	148	296	287	117	179	151	81	190
1940	19,325	1,077	216	146	387	161	106	314	164	97	311
1939	19,907	1,065	267	179	520	174	128	361	173	184	331
1938	19,002	1,171	127	181	381	144	119	167	158	132	258
1937	17,603	1,081	148	296	323	195	66	61	127	88	251
1936	16,707	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1935	16,509	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1934	16,688	1,073	115	197	159	95	30	33	73	68	166
1933	12,631	790	143	181	83	108	39	43	78	39	107
1932	8,257	506	63	94	35	77	17	28	42	23	92
1931	6,480	351	33	94	58	77	26	19	48	23	75
1930	6,025	431	48	65	83	112	33	17	33	23	84
1929	5,663	541	44	15	83	34	25	22	39	21	81
1928	5,727	418	37	35	74	27	14	29	14	17	59
1927	4,148	222	23	10	66	24	5	24	22	21	38
1926	3,319	196	22	11	25	15	5	14	12	6	34
1925	2,572	105	10	5	11	8	4	12	6	17	29
1924	1,843	64	10	6	17	18	8	7	8	29	18
1923	1,562	69	8	4	10	22	6	4	4	4	11
1922	1,657	63	13	8	15	4	3	4	-	7	33
1921	1,589	57	17	14	9	15	-	4	7	10	32
1920	1,368	76	4	28	23	17	5	7	2	8	13

See footnotes at end of table.

TABLE 31.- 1/ CASES OF SELECTED COMMUNICABLE DISEASES BY PREFECTURE: JAPAN, 1920 - 1949 - Cont'd
SCARLET FEVER - Cont'd

Year	Saitama	Chiba	Tokyo	Kanagawa	Niigata	Toyama	Ishikawa	Fukui	Yamanashi	Nagano	Gifu	Shizuoka
*1949	200	147	1,080	250	52	11	8	10	47	381	55	66
1948	127	21	632	101	35	9	4	6	33	132	56	58
1947	58	51	517	117	32	15	6	6	23	99	26	133
1946	66	41	410	96	26	5	5	5	11	76	23	63
1945	121	57	337	102	14	5	2	9	17	142	34	92
1944	195	120	1,813	417	31	11	-	17	39	202	51	159
1943	275	180	3,883	589	33	4	4	12	58	168	45	169
1942	336	167	5,046	711	36	4	4	24	31	143	75	181
1941	305	214	5,833	1,119	52	20	5	29	69	199	115	190
1940	566	296	8,340	1,222	62	16	11	14	69	242	129	237
1939	512	374	8,139	1,164	67	6	6	22	94	355	102	173
1938	350	317	7,788	985	98	7	2	15	144	320	142	188
1937	383	220	6,628	699	110	8	8	9	40	409	89	177
1936	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1935	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1934	218	105	5,889	825	29	20	11	13	46	189	98	219
1933	230	96	4,985	431	21	15	14	11	40	144	84	184
1932	133	47	3,828	337	13	8	12	10	20	112	70	106
1931	173	43	3,123	312	14	8	8	9	10	99	106	94
1930	168	20	2,667	231	16	2	6	5	26	89	64	126
1929	77	31	2,496	240	8	5	5	4	24	48	20	125
1928	54	33	2,592	184	18	7	6	6	31	25	10	98
1927	46	18	1,776	151	9	2	4	1	91	29	11	65
1926	21	20	1,077	90	4	1	4	2	3	22	9	30
1925	20	15	739	61	4	2	1	-	7	14	16	35
1924	15	22	734	57	6	2	-	4	4	14	10	17
1923	21	8	486	76	7	-	-	1	7	17	3	16
1922	18	10	503	69	2	2	2	3	30	17	6	21
1921	25	7	574	77	-	-	3	6	8	12	6	22
1920	23	9	457	52	11	-	4	1	3	4	4	11

See footnotes at end of table.

TABLE 31.-- 1/ CASES OF SELECTED COMMUNICABLE DISEASES BY PREFECTURE: JAPAN, 1920 - 1949 - Cont'd
SCARLET FEVER - Cont'd

Year	Aichi	Mie	Shiga	Kyoto	Osaka	Hyogo	Nara	Wakayama	Tottori	Shimane	Okayama	Hiroshima
1949	182	55	110	262	189	87	31	26	18	60	111	79
1948	131	46	64	119	97	55	7	6	9	19	38	23
1947	121	47	50	139	58	63	10	7	6	30	27	22
1946	84	16	43	127	49	49	11	24	8	9	25	14
1945	84	43	47	178	63	93	7	11	8	11	39	58
1944	280	65	82	331	291	258	7	27	8	16	75	81
1943	453	61	62	411	617	433	20	27	14	7	90	48
1942	536	94	101	685	800	543	26	29	17	14	76	61
1941	468	164	142	785	1,046	645	35	31	23	12	65	74
1940	427	158	101	920	1,585	784	43	45	15	30	104	81
1939	420	150	130	910	1,645	843	53	67	24	64	104	101
1938	469	103	144	963	1,742	853	41	83	40	38	121	149
1937	494	106	170	1,092	1,678	865	31	91	34	35	184	265
1936	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1935	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1934	551	90	117	1,671	2,150	770	60	61	12	13	238	175
1933	759	122	100	1,199	1,284	416	33	36	11	11	140	98
1932	983	73	81	598	614	216	13	37	6	6	77	33
1931	147	40	58	451	315	153	14	18	-	7	51	33
1930	141	64	44	427	330	162	9	32	-	5	123	14
1929	99	72	43	462	386	206	14	40	1	9	69	29
1928	95	63	48	715	431	206	20	22	-	12	53	35
1927	94	73	24	473	284	136	21	24	2	11	25	33
1926	65	50	32	351	242	115	9	16	5	6	36	52
1925	37	31	28	308	159	96	8	10	3	9	16	21
1924	60	28	15	263	113	97	11	8	2	6	37	28
1923	30	8	22	377	111	78	8	15	1	5	48	6
1922	24	18	37	383	121	75	5	11	1	2	93	5
1921	24	19	33	389	105	32	2	9	1	7	4	5
1920	14	8	21	324	120	28	1	4	-	-	12	6

See footnotes at end of table.

TABLE 31.- 1/CASES OF SELECTED COMMUNICABLE DISEASES BY PREFECTURE: JAPAN, 1920 - 1949 - Cont'd
SCARLET FEVER - Cont'd

Year	Yamaguchi	Tokushima	Kagawa	Ehime	Kochi	Fukuoka	Saga	Nagasaki	Kumamoto	Oita	Miyazaki	Kagoshima	Unknown
•1949	50	11	22	24	11	37	5	17	3	8	6	4	-
1948	20	4	18	26	12	56	8	17	2	5	7	8	-
1947	15	3	14	24	9	22	2	27	6	3	11	3	-
1946	25	2	21	22	24	40	2	17	3	3	10	5	-
1945	37	1	20	9	10	55	2	31	6	4	1	2	-
1944	44	22	30	37	27	107	9	58	18	2	10	12	-
1943	47	42	61	44	35	151	8	53	8	6	18	14	-
1942	61	17	80	74	44	156	5	64	24	2	10	18	-
1941	92	23	51	81	53	232	8	91	42	7	7	9	-
1940	119	23	39	74	47	335	22	119	32	11	12	16	-
1939	137	41	48	73	65	410	12	139	36	11	12	16	-
1938	123	28	72	102	89	313	21	199	65	11	9	24	-
1937	108	55	176	74	120	265	12	132	29	9	19	143	-
1936	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1935	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1934	238	25	89	77	40	451	2	117	40	4	14	12	-
1933	95	26	38	33	37	218	1	69	24	1	6	8	-
1932	80	6	36	32	5	114	7	83	39	2	14	29	-
1931	37	6	19	28	4	93	3	86	89	3	10	12	-
1930	41	2	26	30	1	73	3	61	67	-	11	22	-18
1929	58	4	44	26	8	42	1	36	2	6	3	15	-
1928	36	-	41	25	15	67	-	21	14	5	10	5	-
1927	30	3	22	22	10	44	-	25	3	4	124	1	-
1926	50	2	20	17	5	323	26	153	8	9	90	14	-
1925	36	4	101	14	5	537	1	19	5	4	-	19	-
1924	17	1	4	12	3	32	-	5	20	8	2	1	-
1923	17	-	10	10	11	11	-	3	2	3	1	1	-
1922	6	1	6	5	7	8	-	9	1	2	1	2	1
1921	11	-	10	10	5	4	-	2	5	4	1	2	-
1920	15	-	17	16	6	6	-	3	2	2	-	3	-

See footnotes at end of table.

TABLE 31.- 1/ CASES OF SELECTED COMMUNICABLE DISEASES BY PREFECTURE: JAPAN, 1920 - 1949 - Cont'd
3/ DIPHTEHERIA

Year	All Japan	Hokkaido	Aomori	Iwate	Miyagi	Akita	Yamagata	Fukushima	Ibaraki	Tochigi	Gumma
1949	14,825	871	263	334	435	383	210	278	265	225	213
1948	16,198	1,289	231	290	459	566	289	220	313	329	255
1947	28,346	2,369	501	421	595	697	668	416	538	710	335
1946	49,166	4,149	772	838	942	1,012	1,260	1,299	599	921	258
1945	85,833	5,792	1,373	1,188	1,681	1,089	1,595	2,494	971	2,146	688
1944	94,274	5,672	1,534	1,348	1,907	1,050	1,524	3,713	1,391	2,196	566
1943	63,761	3,786	1,019	1,451	1,329	893	872	2,569	607	988	566
1942	44,431	2,502	579	819	880	814	563	1,922	571	574	603
1941	40,442	2,317	674	1,074	684	1,147	543	1,958	548	370	430
1940	38,303	2,342	888	1,089	805	1,030	644	1,926	526	276	470
1939	35,801	2,251	867	987	810	903	518	1,523	486	291	631
1938	28,323	2,345	715	693	595	900	394	1,115	306	323	414
1937	28,001	2,863	862	581	642	991	478	987	412	445	568
1936	28,234	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1935	28,200	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1934	29,992	3,337	710	554	728	1,052	434	1,313	547	559	564
1933	28,500	3,266	516	477	729	1,127	445	1,159	323	595	544
1932	21,811	2,130	367	425	570	758	494	1,042	357	494	484
1931	21,048	2,304	389	462	489	737	319	1,044	415	399	522
1930	18,522	2,220	546	409	694	762	432	898	606	431	380
1929	19,674	2,065	398	251	561	468	374	752	462	444	364
1928	17,526	2,037	322	234	472	388	294	742	358	352	513
1927	15,172	1,806	246	237	442	371	290	801	266	267	251
1926	13,621	1,985	239	255	520	389	240	761	224	232	302
1925	13,775	1,692	308	225	478	335	308	805	274	260	367
1924	13,081	1,590	250	271	521	374	315	671	230	325	338
1923	12,729	1,722	324	212	455	454	386	678	271	325	403
1922	13,696	1,330	343	255	572	625	488	896	385	357	465
1921	14,449	1,878	476	306	458	643	367	552	330	376	436
1920	15,113	2,161	445	377	560	525	307	664	397	439	448

See footnotes at end of table.

TABLE 31.- 1/ CASES OF SELECTED COMMUNICABLE DISEASES BY PREFECTURE: JAPAN, 1920 - 1949 - Cont'd
3/ DIPHTHERIA - Cont'd

Year	Saitama	Chiba	Tokyo	Kanagawa	Niigata	Toyama	Ishikawa	Fukui	Yamanashi	Nagano	Gifu	Shizuoka
1949	339	223	1,048	395	595	251	250	169	67	300	167	285
1948	370	181	1,034	447	665	173	281	130	65	398	130	275
1947	649	420	1,603	564	843	227	626	226	109	626	204	555
1946	956	864	2,252	1,056	1,269	490	960	366	220	826	463	1,304
1945	2,819	1,664	4,503	2,830	2,187	1,043	1,737	680	522	2,213	1,077	2,895
1944	3,298	2,412	15,011	4,031	1,881	977	1,347	543	753	1,682	713	2,366
1943	2,586	1,453	13,565	2,881	1,395	367	310	337	414	1,350	470	1,202
1942	1,742	893	10,355	3,060	1,021	56	186	204	299	982	284	803
1941	1,104	639	7,754	2,546	798	52	162	191	262	1,017	347	700
1940	828	750	7,396	2,287	580	38	137	155	296	842	296	620
1939	569	543	6,712	2,125	570	26	158	127	238	928	294	622
1938	453	462	4,405	1,339	661	36	125	82	150	832	229	406
1937	664	448	4,058	1,437	647	34	105	61	179	912	214	365
1936	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1935	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1934	721	387	5,339	1,379	475	35	53	88	147	887	238	477
1933	629	427	6,066	1,404	536	47	58	90	131	735	204	495
1932	454	282	4,128	1,044	415	44	54	120	127	591	194	526
1931	506	325	3,871	1,060	448	44	36	120	103	571	130	437
1930	512	296	2,561	640	518	38	70	108	106	518	142	327
1929	454	297	3,437	810	466	43	74	73	117	607	122	363
1928	342	225	2,999	745	463	51	64	47	140	503	145	294
1927	274	243	2,272	524	395	22	59	66	124	419	131	236
1926	235	210	1,456	354	439	34	51	59	151	545	163	246
1925	264	266	1,599	416	454	46	49	56	164	512	147	297
1924	278	293	1,508	436	489	46	62	65	173	482	119	374
1923	339	308	1,003	292	438	36	92	119	137	592	160	358
1922	430	330	1,074	338	469	33	123	103	180	655	148	326
1921	415	339	1,269	403	550	46	138	74	134	537	171	412
1920	468	332	1,086	478	573	39	160	95	113	519	155	366

See footnotes at end of table.

TABLE 31.- 1/ CASES OF SELECTED COMMUNICABLE DISEASES BY PREFECTURE; JAPAN, 1920 - 1949 - Cont'd
3/ DIPHTHERIA - Cont'd

Year	Aichi	Mie	Shiga	Kyoto	Osaka	Hyogo	Nara	Wakayama	Tottori	Shimane	Okayama	Hiroshima
*1949	374	204	106	202	361	458	99	101	86	276	189	420
1948	458	264	107	259	304	486	141	119	91	304	266	428
1947	1,591	679	215	505	427	877	173	225	176	515	386	659
1946	1,989	1,161	679	963	1,003	1,632	428	579	376	879	946	1,244
1945	5,363	1,860	1,167	3,161	3,407	4,025	917	631	507	990	2,003	2,705
1944	7,269	1,599	938	2,505	3,983	4,280	231	339	451	594	1,670	2,087
1943	4,098	992	546	2,290	3,546	2,832	177	339	311	295	726	1,892
1942	2,048	568	399	1,742	2,744	2,050	191	215	203	215	436	686
1941	1,145	445	311	2,086	3,023	1,985	168	198	258	366	340	721
1940	1,073	380	250	1,650	3,101	2,019	177	255	200	393	273	560
1939	1,079	386	254	1,700	3,812	1,508	157	276	210	195	283	402
1938	915	367	215	1,222	3,211	1,076	138	290	150	165	246	280
1937	707	380	149	884	2,665	1,183	154	185	148	227	290	244
1936	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1935	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1934	484	429	257	1,180	2,763	906	92	362	146	170	263	262
1933	364	422	170	748	2,813	760	87	282	130	94	143	178
1932	355	316	142	661	1,828	590	86	210	71	138	144	207
1931	415	360	168	740	1,622	540	120	170	66	107	171	131
1930	376	303	84	640	1,163	515	87	138	72	86	163	200
1929	375	415	84	629	1,864	613	108	161	73	141	266	199
1928	298	335	86	499	1,519	657	73	136	44	100	181	190
1927	306	339	121	450	1,327	469	98	193	39	78	142	185
1926	294	402	135	414	1,080	484	56	169	110	89	160	281
1925	321	275	158	319	1,007	375	63	162	66	63	88	235
1924	377	290	93	359	680	309	46	122	51	80	70	143
1923	347	320	134	289	633	267	65	134	72	92	76	123
1922	383	337	100	361	548	316	52	128	98	129	87	132
1921	383	339	159	471	635	182	85	132	95	141	125	183
1920	368	349	182	387	825	214	76	222	60	124	110	77

See footnotes at end of table.

Table 31.- 1/ CASES OF SELECTED COMMUNICABLE DISEASES BY PREFECTURE, JAPAN, 1920 - 1949 Cont'd

3/ DIPHTHERIA - Cont'd

Year	Yamaguchi	Tokushima	Kagawa	Ehime	Kochi	Fukuoka	Saga	Nagasaki	Kumamoto	Oita	Miyazaki	Kagoshima	Unknown
1949	372	124	120	217	136	944	428	441	253	383	526	439	-
1948	326	127	117	350	146	891	553	460	179	550	419	463	-
1947	712	291	272	885	326	1,777	840	673	230	775	580	655	-
1946	2,023	618	803	1,260	1,246	3,263	897	999	370	1,130	812	790	-
1945	2,453	934	1,446	1,328	1,835	3,825	663	900	377	1,275	340	514	-
1944	2,302	528	677	1,072	1,148	1,871	533	1,426	212	1,441	352	851	-
1943	1,105	270	396	709	213	1,110	263	609	104	166	153	204	-
1942	651	127	172	520	133	691	159	402	89	33	148	69	-
1941	902	101	170	407	173	900	260	736	113	47	192	78	-
1940	842	86	82	330	173	828	291	678	129	54	158	100	-
1939	755	118	137	312	159	800	208	518	102	39	138	74	-
1938	572	120	119	334	138	752	269	416	72	46	134	86	-
1937	627	134	105	265	225	877	157	298	53	66	122	103	-
1936	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1935	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1934	686	103	158	330	123	569	37	299	48	57	125	119	-
1933	589	60	137	243	208	475	21	266	44	41	103	99	-
1932	551	210	86	210	195	982	19	251	55	27	128	77	-
1931	345	45	99	203	119	431	10	188	45	27	119	76	-
1930	184	52	109	201	80	292	11	193	44	20	216	77	-
1929	310	48	128	215	78	299	13	225	43	29	230	126	-
1928	314	79	94	198	63	592	17	226	65	34	119	87	-
1927	333	64	84	221	71	362	13	250	63	38	75	109	-
1926	249	59	88	196	70	330	18	203	59	49	58	78	-
1925	234	61	70	216	51	283	31	185	41	42	46	61	-
1924	222	47	62	218	38	253	12	176	48	41	63	71	-
1923	160	41	88	163	44	214	17	148	67	23	56	52	-
1922	162	50	152	80	80	169	13	178	52	24	43	88	9
1921	218	50	101	169	71	178	15	151	60	42	66	88	-
1920	265	56	115	190	113	185	14	183	83	52	64	92	-

See footnotes at end of table.

TABLE 31.-- 1/ CASES OF SELECTED COMMUNICABLE DISEASES BY PREFECTURE: JAPAN, 1920 - 1949 - Cont'd

4/ DYSENTERY

Year	All Japan	Hokkaido	Aomori	Iwate	Miyagi	Akita	Yamagata	Fukushima	Ibaraki	Tochigi	Gumma
1949	24,001	483	136	621	305	232	434	454	1,067	608	1,499
1948	14,628	932	120	528	228	195	237	471	746	279	472
1947	39,249	1,368	302	1,093	766	453	1,640	2,207	1,201	1,375	1,375
1946	87,737	3,713	1,466	6,786	3,928	2,476	4,531	8,036	3,077	2,527	3,453
1945	96,462	1,031	1,225	675	241	1,036	974	1,591	973	969	2,130
1944	55,196	1,543	966	348	148	503	422	659	1,234	561	795
1943	50,188	376	792	973	181	255	267	345	617	309	1,168
1942	55,785	1,06	314	684	161	322	152	577	1,091	495	518
1941	58,803	150	369	448	200	232	154	562	1,346	692	1,077
1940	83,689	200	128	660	275	127	128	687	2,162	1,010	1,720
1939	97,259	599	209	362	342	63	166	704	1,942	1,035	954
1938	80,221	165	146	349	299	73	110	426	1,589	756	1,040
1937	78,286	117	134	347	173	49	129	308	1,447	454	920
1936	52,075	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1935	48,968	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1934	42,939	76	41	301	218	61	42	208	850	332	831
1933	38,040	53	62	196	114	45	22	137	826	241	421
1932	32,249	47	40	169	101	47	27	96	709	284	587
1931	29,655	69	42	126	139	26	13	103	730	284	561
1930	29,672	51	62	251	110	40	58	127	450	180	556
1929	30,230	98	54	92	136	31	99	136	651	301	538
1928	25,193	63	73	136	147	31	92	105	439	193	514
1927	21,379	147	51	66	84	18	9	40	327	82	388
1926	17,132	70	81	60	39	15	16	39	344	90	283
1925	14,719	131	180	213	76	117	25	47	245	62	187
1924	18,647	33	372	582	97	395	21	130	227	75	234
1923	20,266	44	554	595	129	87	26	120	298	127	188
1922	15,102	48	1,232	735	97	172	7	215	257	28	87
1921	12,443	37	861	665	124	267	23	262	174	44	61
1920	12,733	20	685	783	315	450	29	206	208	102	61

See footnotes at end of table.

TABLE 31.- 1/ CASES OF SELECTED COMMUNICABLE DISEASES BY PREFECTURE, JAPAN, 1920 - 1949 - Cont'd
4/ DYSENTERY - Cont'd

Year	Saitama	Chiba	Tokyo	Kanagawa	Niigata	Toyama	Ishikawa	Fukui	Yamanashi	Nagano	Gifu	Shizuoka
*1949	1,793	862	3,372	983	1,657	140	175	114	154	528	474	1,072
1948	406	284	1,537	296	602	61	36	202	76	331	399	331
1947	1,726	1,005	2,965	708	1,751	189	208	369	667	1,596	633	1,202
1946	1,999	1,258	1,844	1,581	2,580	533	205	336	1,313	2,106	1,005	2,879
1945	1,784	943	2,651	1,675	978	25	26	162	1,453	3,316	1,647	8,810
1944	1,475	534	4,791	2,979	552	20	51	130	509	996	961	2,499
1943	1,022	685	7,716	1,995	676	35	24	89	234	486	611	2,614
1942	1,074	872	14,582	3,754	200	12	31	166	204	144	895	1,860
1941	1,529	1,127	15,703	3,925	259	28	25	121	485	165	1,095	2,042
1940	1,985	1,613	24,659	6,083	338	15	47	73	567	217	1,264	2,388
1939	1,589	1,582	27,412	5,387	339	10	38	67	337	193	1,002	2,653
1938	1,603	1,686	22,874	3,868	248	34	34	65	146	149	1,039	2,811
1937	1,210	1,457	20,830	2,813	183	11	28	42	90	117	707	2,207
1936	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1935	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1934	797	612	12,621	2,016	168	12	30	52	44	83	639	1,820
1933	633	645	13,083	1,412	202	21	39	48	58	63	526	1,733
1932	620	473	10,102	1,088	76	10	23	49	55	38	664	1,614
1931	631	418	9,439	841	61	8	43	18	41	66	599	1,649
1930	414	338	9,554	963	42	18	22	16	63	62	441	2,100
1929	505	698	8,345	888	28	5	19	20	65	100	490	1,979
1928	538	423	7,236	958	37	38	34	19	107	80	450	1,896
1927	456	280	5,469	581	172	3	25	18	62	64	403	1,444
1926	396	241	2,957	333	23	13	31	6	95	37	371	1,447
1925	328	159	2,643	349	27	-	29	19	93	44	350	1,126
1924	300	306	2,970	277	273	6	16	8	68	103	271	1,314
1923	346	506	3,605	546	60	9	25	7	166	64	320	1,644
1922	147	207	1,152	310	79	8	73	10	78	47	160	591
1921	122	417	773	234	55	3	55	23	84	28	284	418
1920	218	114	675	311	79	7	127	34	181	23	277	444

See footnotes at end of table.

TABLE 31.- 1/ CASES OF SELECTED COMMUNICABLE DISEASES BY PREFECTURE: JAPAN, 1920 - 1949 - Cont'd
4/ DYSENTERY - Cont'd

Year	Aichi	Mie	Shiga	Kyoto	Osaka	Hyogo	Nara	Yekeyama	Tottori	Shimane	Okeyama	Hiroshima
*1949	1,163	293	76	388	596	331	48	112	77	312	173	423
1948	698	181	94	293	520	457	46	57	67	219	321	321
1947	1,889	491	298	857	915	1,389	174	141	183	463	425	597
1946	1,935	462	451	979	814	1,685	255	301	662	747	1,024	1,521
1945	4,029	979	680	1,356	1,613	1,612	553	1,609	1,012	1,683	1,444	5,148
1944	1,390	411	198	1,277	2,189	2,968	328	214	469	844	823	2,743
1943	1,796	472	160	1,710	5,709	2,321	73	175	313	349	722	2,746
1942	2,686	629	226	2,071	6,469	3,480	112	224	154	325	519	1,483
1941	2,337	757	234	2,082	6,026	2,825	156	313	182	414	734	1,557
1940	2,923	686	244	2,602	9,122	4,182	234	401	239	555	910	1,956
1939	2,704	745	240	2,891	13,219	5,452	285	434	360	1,071	1,536	3,365
1938	2,713	768	216	2,346	10,648	4,581	178	343	220	489	925	1,816
1937	2,057	785	139	2,225	7,934	3,458	90	214	194	388	628	1,572
1936	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1935	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1934	1,248	466	122	1,754	5,300	2,420	96	122	137	201	578	1,305
1933	767	432	135	1,717	4,492	2,461	61	182	115	233	496	624
1932	893	275	84	1,609	3,822	1,587	61	87	103	248	277	606
1931	802	273	115	1,422	3,133	1,312	32	67	68	197	258	622
1930	844	284	121	1,232	2,662	1,360	41	63	28	243	273	647
1929	1,037	290	175	1,658	2,845	1,188	57	75	40	233	445	635
1928	706	408	83	1,540	1,919	867	13	63	14	198	316	608
1927	645	373	58	1,184	1,612	1,168	19	80	18	177	414	696
1926	524	347	82	1,101	1,365	809	9	104	58	208	427	505
1925	572	336	81	862	816	664	20	29	28	160	330	380
1924	665	217	43	770	719	669	8	82	20	206	413	643
1923	704	324	36	833	727	849	27	49	24	251	338	595
1922	264	74	16	434	656	1,017	4	43	32	64	498	1,100
1921	316	165	25	414	433	496	77	37	13	78	380	1,080
1920	286	147	58	375	469	423	47	76	12	85	188	633

See footnotes at end of table.

TABLE 31.- 1/ CASES OF SELECTED COMMUNICABLE DISEASES BY PREFECTURE: JAPAN, 1920 - 1949 - Cont'd
4/ DYSENTERY - Cont'd

Year	Yamaguchi	Tokushima	Kagawa	Ehime	Kochi	Fukuoka	Saga	Nagasaki	Kumamoto	Oita	Miyazaki	Kagoshima	Unknown
1949	254	171	243	311	122	445	117	214	209	188	396	166	-
1948	242	155	128	359	111	433	182	250	218	191	262	155	-
1947	287	842	523	983	307	636	212	542	352	321	536	708	-
1946	1,134	1,455	1,985	2,068	1,219	2,312	1,147	1,742	1,868	976	2,084	1,279	-
1945	3,055	2,080	3,898	3,714	1,167	11,278	1,589	3,567	2,823	2,258	1,458	3,522	-
1944	1,580	1,021	3,365	1,399	965	4,574	792	1,809	973	645	1,157	446	-
1943	1,743	2,512	1,552	1,136	509	2,007	124	660	930	293	460	246	-
1942	1,610	821	655	424	463	2,721	206	579	825	215	414	230	-
1941	1,260	895	829	477	510	2,274	295	678	1,165	488	284	307	-
1940	1,895	1,067	1,456	538	708	3,521	257	1,099	1,197	686	514	351	-
1939	2,408	1,360	1,462	751	768	6,022	237	1,286	1,618	652	448	960	-
1938	1,998	1,082	869	540	1,075	5,664	379	794	1,789	388	413	477	-
1937	1,700	892	679	398	800	16,213	194	837	1,701	395	354	666	-
1936	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1935	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1934	1,127	805	653	319	437	1,424	42	416	900	204	338	671	-
1933	1,045	617	506	224	397	1,034	46	464	816	131	230	233	-
1932	833	841	405	226	425	895	63	433	953	213	247	144	-
1931	877	702	460	239	264	817	48	476	897	198	161	308	-
1930	1,049	534	324	283	319	1,039	56	660	973	379	196	144	-
1929	1,396	452	447	289	261	1,114	51	633	973	174	302	182	-
1928	830	390	443	289	298	808	28	464	880	125	211	81	-
1927	628	408	408	236	329	777	45	359	735	193	151	52	-
1926	747	525	452	243	226	750	81	415	683	267	106	111	-
1925	667	509	413	186	157	595	51	369	683	153	89	119	-
1924	561	1,059	649	175	209	1,458	215	424	989	156	155	64	-
1923	659	1,407	378	327	165	1,049	196	524	774	384	127	56	-3
1922	300	578	740	208	177	653	170	558	736	820	101	123	-4
1921	240	205	373	116	81	359	41	846	1,030	410	94	123	-3
1920	335	402	702	197	80	506	99	667	975	399	88	125	410

See footnotes at end of table.

TABLE 31.- 1/ CASES OF SELECTED COMMUNICABLE DISEASES BY PREFECTURE: JAPAN, 1920 - 1949 - Cont'd
SMALLPOX

Year	All Japan	Hokkaido	Aomori	Iwate	Miyagi	Akita	Yamagata	Fukushima	Ibaraki	Tochigi	Gumma
*1949	124	-	-	-	-	1	-	-	-	-	-
1948	29	9	-	-	-	-	-	-	1	-	-
1947	391	48	-	1	1	12	8	1	21	23	3
1946	17,800	2,328	75	49	113	71	120	230	183	74	91
1945	1,614	390	1	9	32	4	-	155	16	-	6
1944	311	40	-	-	8	-	-	29	-	2	3
1943	546	3	1	16	10	1	1	-	41	4	3
1942	381	2	-	1	-	2	-	20	2	-	-
1941	654	202	58	2	15	15	1	11	3	-	-
1940	575	17	11	24	36	1	13	1	-	-	4
1939	287	1	-	-	2	-	-	-	10	8	-
1938	60	-	-	-	-	-	-	-	-	-	-
1937	90	-	-	-	-	-	-	-	-	-	-
1936	478	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1935	113	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1934	320	12	5	-	-	-	1	-	-	-	-
1933	375	-	-	-	-	36	-	2	-	23	2
1932	305	4	-	2	-	-	-	-	-	2	-
1931	23	-	-	-	-	-	-	-	-	-	-
1930	7	-	-	-	-	-	-	-	-	-	-
1929	114	1	-	-	-	-	-	-	-	-	-
1928	723	2	-	1	-	-	-	20	-	2	-
1927	352	4	-	-	-	-	-	1	-	-	-
1926	1,256	11	-	1	-	-	-	-	-	-	-
1925	430	1	-	2	-	-	-	-	-	1	111
1924	1,702	555	12	39	56	61	79	41	8	32	2
1923	1,922	249	7	-	3	19	-	-	1	-	-
1922	679	27	-	-	3	9	-	7	14	4	-
1921	899	-	-	1	-	-	1	6	12	7	20
1920	9,167	33	2	5	-	111	33	37	9	1	2

See footnotes at end of table.

TABLE 31.- 1/ CASES OF SELECTED COMMUNICABLE DISEASES BY PREFECTURE; JAPAN, 1920 - 1949 - Cont'd

SMALLPOX - Cont'd

Year	Saitama	Chiba	Tokyo	Kanagawa	Niigata	Toyama	Ishikawa	Fukui	Yamanashi	Nagano	Gifu	Shizuoka
*1949	-	-	3	2	1	-	-	-	-	-	-	-
1948	-	2	-	-	-	-	-	-	-	-	1	1
1947	3	13	18	4	4	1	1	-	-	-	-	4
1946	207	218	1,821	427	134	187	120	144	87	189	250	251
1945	36	1	23	48	16	8	6	2	-	37	-	61
1944	-	2	13	-	17	12	66	2	-	-	-	-
1943	-	-	36	5	2	-	8	3	2	1	2	5
1942	-	2	36	3	-	-	-	25	-	14	-	-
1941	1	20	19	7	65	10	1	2	1	1	1	3
1940	11	1	37	5	37	3	7	1	4	13	14	19
1939	-	1	5	2	-	-	-	-	-	-	-	-
1938	-	-	1	-	-	-	-	-	-	-	-	-
1937	-	2	-	1	-	2	-	-	-	-	-	-
1936	NA	NA	NA	NA	NA	NA	NA	NA	NA	25	NA	5
1935	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1934	2	-	2	31	-	-	-	1	2	-	-	9
1933	1	2	63	1	-	1	1	1	-	-	-	-
1932	1	-	-	3	-	-	2	-	-	-	-	18
1931	-	1	-	1	-	-	-	-	-	-	-	-
1930	-	-	1	-	-	-	4	-	-	-	-	-
1929	-	-	2	1	14	2	2	-	-	-	-	1
1928	-	3	67	12	1	1	5	-	10	-	-	1
1927	-	6	-	4	-	-	-	-	-	-	-	-
1926	-	1	5	71	-	7	3	-	6	-	-	1
1925	-	1	1	7	-	-	-	-	-	-	-	-
1924	5	1	181	6	1	21	15	10	16	3	16	6
1923	-	3	19	6	2	-	1	1	-	2	5	1
1922	-	-	16	13	-	40	-	15	1	-	-	-
1921	1	-	4	6	-	-	-	-	-	-	-	2
1920	2	4	118	1	20	-	13	15	-	2	3	22

See footnotes at end of table.

TABLE 31.- 1/ CASES OF SELECTED COMMUNICABLE DISEASES BY PREFECTURE: JAPAN, 1920 - 1949 - Cont'd
SMALLPOX - Cont'd

Year	Aichi	Mie	Shiga	Kyoto	Osaka	Hyogo	Nara	Wakayama	Tottori	Shimane	Okayama	Hiroshima
* 1949	-	2	-	-	62	-	1	9	3	-	-	-
1948	-	-	-	2	2	-	-	1	-	1	2	1
1947	9	5	-	1	11	42	1	34	1	7	11	3
1946	1,176	232	162	697	2,432	2,160	406	227	55	127	230	397
1945	48	-	13	4	70	332	2	-	3	28	7	15
1944	12	10	-	-	4	3	-	7	-	1	-	-
1943	-	-	-	2	17	225	2	7	1	-	7	-
1942	4	-	-	2	1	7	-	5	-	-	87	6
1941	1	11	1	-	49	7	2	-	62	-	1	10
1940	8	-	-	3	44	8	13	1	2	-	9	131
1939	4	-	-	23	14	106	-	-	-	-	2	5
1938	1	-	-	3	-	4	-	-	-	-	7	-
1937	-	-	-	-	2	1	-	-	-	-	-	-
1936	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1935	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1934	-	-	8	-	11	4	-	-	1	-	9	29
1933	17	1	2	12	16	3	-	-	15	-	-	-
1932	-	1	-	-	201	11	1	1	-	-	1	-
1931	4	-	-	-	-	2	-	-	-	-	-	-
1930	-	-	-	-	-	-	-	-	-	-	-	-
1929	-	-	-	-	2	1	-	-	-	67	1	-
1928	12	3	1	3	48	17	2	1	-	-	4	3
1927	-	-	-	-	3	4	-	-	3	-	-	-
1926	2	35	-	6	29	31	3	1	-	1	-	27
1925	-	-	-	3	56	10	-	-	1	1	3	9
1924	44	24	21	11	21	36	-	3	-	-	-	-
1923	2	-	4	5	236	107	3	-	13	2	7	371
1922	-	-	4	9	5	17	3	-	7	1	6	14
1921	13	1	-	3	90	96	4	-	-	-	6	2
1920	10	23	62	56	1,209	86	137	133	6	22	27	74

See footnotes at end of table.

TABLE 31.-- 1/ CASES OF SELECTED COMMUNICABLE DISEASES BY PREFECTURE: JAPAN, 1920 - 1949 - Cont'd

Year	SMALLPOX - Cont'd										2/	
	Yamaguchi	Tokushima	Kagawa	Ehime	Kochi	Fukuoka	Saga	Nagasaki	Kumamoto	Oita	Miyazaki	Kagoshima
1949	12	-	-	4	-	21	-	1	-	2	-	-
1948	-	-	-	-	-	1	5	-	-	-	-	-
1947	7	1	4	13	1	40	5	2	3	2	1	18
1946	127	173	290	161	185	427	54	324	84	91	41	173
1945	76	-	8	1	-	32	4	111	2	3	1	3
1944	2	-	2	2	-	9	1	15	18	3	-	28
1943	35	4	6	10	7	23	-	5	17	22	1	11
1942	3	-	2	-	1	9	3	30	2	2	56	56
1941	8	8	1	8	13	26	3	4	-	-	-	1
1940	32	-	2	13	1	13	2	7	13	8	3	3
1939	44	50	2	-	3	3	-	7	-	1	1	-
1938	-	-	-	-	-	3	28	-	1	-	2	3
1937	17	-	-	-	-	25	6	1	1	3	-	-
1936	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1935	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1934	19	-	-	22	-	5	2	1	102	25	-	20
1933	3	-	-	20	2	12	-	1	7	12	5	112
1932	1	1	8	-	2	9	-	14	6	3	10	3
1931	-	-	-	-	-	11	-	-	3	1	-	-
1930	-	-	1	-	-	-	-	1	-	-	-	-
1929	-	-	-	-	1	3	-	5	-	-	-	-
1928	98	2	-	5	40	103	1	5	90	155	5	11
1927	4	-	-	-	1	252	2	52	1	15	-	-
1926	42	36	169	2	7	390	37	93	54	128	5	52
1925	69	-	-	1	-	40	-	105	-	4	4	-
1924	37	-	7	1	-	70	-	-	31	176	1	53
1923	95	73	31	29	-	235	49	58	105	151	29	4
1922	83	-	1	24	-	239	14	68	15	16	-	2
1921	34	8	86	5	11	120	4	154	166	23	1	2
1920	73	82	16	33	132	252	1	19	22	220	12	29

See footnotes at end of table.

TABLE 31.- 1/ CASES OF SELECTED COMMUNICABLE DISEASES BY PREFECTURE: JAPAN, 1920 - 1949 - Cont'd
TYPHUS FEVER

Year	All Japan	Hokkaido	Aomori	Iwate	Miyagi	Akita	Yamagata	Fukushima	Ibakeki	Tochigi	Gumma
*1949	121	1	-	-	13	-	1	11	-	-	-
1948	474	5	18	7	4	5	4	4	5	-	-
1947	1,114	56	8	-	20	2	42	4	36	11	4
1946	31,141	2,452	846	121	467	219	529	284	281	160	108
1945	2,461	1,669	15	31	2	-	41	8	25	-	18
1944	3,941	2,067	105	17	162	64	88	14	87	11	-
1943	1,374	1,088	12	5	32	69	57	-	-	-	-
1942	100	-	1	-	62	-	6	-	-	-	1
1941	87	-	-	-	31	-	20	-	-	1	-
1940	3	-	-	-	-	-	-	-	-	-	-
1939	5	-	-	-	-	-	-	-	-	-	-
1938	-	-	-	-	-	-	-	-	-	-	-
1937	17	-	16	-	-	-	-	-	-	-	-
1936	1	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1935	18	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1934	26	-	25	-	-	-	-	-	-	-	-
1933	4	-	-	-	-	-	-	-	-	-	-
1932	3	-	-	-	-	-	-	-	-	-	-
1931	3	-	1	-	-	-	1	-	-	-	-
1930	1	-	-	-	-	-	-	-	-	-	-
1929	15	-	-	-	8	-	5	-	-	-	-
1928	23	-	-	-	5	-	17	1	-	-	-
1927	7	-	-	-	2	-	4	-	-	-	-
1926	36	-	-	-	-	-	34	-	-	-	-
1925	28	-	-	-	18	-	10	-	-	-	-
1924	14	-	9	-	-	2	-	-	-	-	1
1923	14	8	-	-	4	-	-	-	-	-	-
1922	23	-	6	-	-	1	14	1	-	-	-
1921	171	-	3	-	82	14	70	-	-	-	-
1920	66	2	31	-	-	7	25	1	-	-	-

See footnotes at end of table.

TABLE 31.- 1/ CASES OF SELECTED COMMUNICABLE DISEASES BY PREFECTURE: JAPAN, 1920 - 1949 - Cont'd

Year	Saitama	Chiba	Tokyo	Kanagawa	Niigata	Toyama	Ishikawa	Fukui	Yamanashi	Nagano	Gifu	Shizuoka
*1949	2	1	23	11	-	-	-	-	-	-	-	-
1948	10	6	55	47	2	6	1	-	-	1	10	-
1947	32	26	233	48	12	8	10	6	7	9	27	30
1946	704	341	9,587	906	126	40	79	57	140	242	86	85
1945	-	2	32	77	5	-	-	-	-	-	-	22
1944	1	3	840	52	-	-	-	2	-	1	-	-
1943	1	2	59	-	-	-	-	-	1	-	-	-
1942	-	-	23	-	-	-	-	2	-	-	-	-
1941	9	-	20	-	-	-	-	-	-	-	-	-
1940	-	-	1	-	1	-	1	-	-	-	-	-
1939	-	-	-	-	-	-	-	-	-	-	-	-
1938	-	-	-	-	-	-	-	-	-	-	-	-
1937	-	-	-	-	-	-	-	-	-	-	-	-
1936	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1935	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1934	-	-	-	-	-	-	-	-	-	-	-	-
1933	-	-	-	-	-	-	-	-	-	-	-	-
1932	-	-	-	-	-	1	-	-	-	-	-	-
1931	-	-	-	-	1	-	-	-	-	-	-	-
1930	-	-	-	-	-	1	-	-	-	-	-	-
1929	-	-	-	-	-	-	-	-	-	-	-	-
1928	-	-	-	-	-	-	-	-	-	-	-	-
1927	-	-	-	-	-	-	-	-	-	-	-	-
1926	-	-	-	-	-	2	-	-	-	-	-	-
1925	-	-	-	-	-	-	-	-	-	-	-	-
1924	-	-	-	-	-	-	-	-	-	-	-	2
1923	-	-	-	-	-	-	-	-	-	-	-	-
1922	-	-	-	-	-	-	-	-	-	-	-	-
1921	-	-	-	-	-	-	-	-	-	-	-	-
1920	-	-	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

TABLE 31.- 1/ CASES OF SELECTED COMMUNICABLE DISEASES BY PREFECTURE: JAPAN, 1920 - 1949 - Cont'd
TYPHUS FEVER - Cont'd

Year	Aichi	Mie	Shiga	Kyoto	Osaka	Hyogo	Nara	Wakayama	Tottori	Shimane	Okayama	Hiroshima
*1949	-	2	-	-	27	1	-	4	-	10	-	1
1948	4	1	1	38	150	3	21	-	-	12	-	12
1947	227	5	-	7	89	6	2	18	7	12	5	2
1946	1,124	85	63	1,065	6,388	2,664	189	95	45	26	130	61
1945	5	-	-	8	-	49	-	-	-	-	12	-
1944	1	-	-	1	60	10	-	-	-	-	-	-
1943	1	-	-	2	2	28	-	-	2	1	-	-
1942	3	-	-	-	-	1	-	-	-	-	-	-
1941	-	-	-	-	-	2	-	-	-	-	3	-
1940	-	-	-	-	-	-	-	-	-	-	-	-
1939	-	-	-	-	-	-	-	-	-	-	-	-
1938	-	-	-	-	-	-	-	-	-	-	-	-
1937	-	-	-	-	-	-	-	-	-	-	-	-
1936	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1935	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1934	-	-	-	-	-	1	-	-	-	-	-	-
1933	-	-	-	-	1	3	-	-	-	-	-	-
1932	-	-	-	-	-	-	-	-	-	-	-	1
1931	-	-	-	-	-	-	-	-	-	-	-	-
1930	-	-	-	-	-	-	-	-	-	-	-	-
1929	-	-	-	-	-	-	-	2	-	-	-	-
1928	-	-	-	-	-	-	-	-	-	-	-	-
1927	-	-	-	-	-	-	-	-	-	-	-	1
1926	-	-	-	-	-	-	-	-	-	-	-	-
1925	-	-	-	-	-	-	-	-	-	-	-	-
1924	-	-	-	-	-	-	-	-	-	-	-	-
1923	-	1	-	-	-	1	-	-	-	-	-	-
1922	-	-	-	-	-	-	-	-	1	-	-	-
1921	-	-	-	-	-	-	-	-	-	-	1	1
1920	-	-	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

TABLE 31.- 1/ CASES OF SELECTED COMMUNICABLE DISEASES BY PREFECTURE; JAPAN, 1920 - 1949 - Cont'd
TYPHUS FEVER - Cont'd

Year	Yamaguchi	Tokushima	Kagawa	Ehime	Kochi	Fukuoka	Saga	Nagasaki	Kumamoto	Oita	Miyazaki	Kagoshima	2/ Unknown
*1949	-	-	-	-	-	1	-	11	1	-	-	-	-
1948	1	-	8	-	-	4	-	26	3	-	-	-	-
1947	16	2	54	6	2	3	2	7	3	1	-	-	-
1946	92	22	76	35	43	481	41	439	10	34	7	51	-
1945	3	-	7	4	-	401	22	3	-	-	22	-	-
1944	1	-	-	-	-	88	59	-	-	-	2	205	-
1943	2	-	-	1	-	4	2	2	-	-	-	1	-
1942	-	-	-	-	-	-	-	1	-	-	-	-	-
1941	-	-	-	-	-	-	-	-	1	-	-	-	-
1940	-	-	-	-	-	-	-	-	-	-	-	-	-
1939	-	-	-	-	-	5	-	-	-	-	-	-	-
1938	-	-	-	-	-	-	-	-	-	-	-	-	-
1937	-	-	1	-	-	-	-	-	-	-	-	-	-
1936	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1935	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1934	-	-	-	-	-	-	-	-	-	-	-	-	-
1933	-	-	-	-	-	-	-	-	-	-	-	-	-
1932	-	-	-	1	-	-	-	-	-	-	-	-	-
1931	-	-	-	-	-	-	-	-	-	-	-	-	-
1930	-	-	-	-	-	-	-	-	-	-	-	-	-
1929	-	-	-	-	-	-	-	-	-	-	-	-	-
1928	-	-	-	-	-	-	-	-	-	-	-	-	-
1927	-	-	-	-	-	-	-	-	-	-	-	-	-
1926	-	-	-	-	-	-	-	-	-	-	-	-	-
1925	-	-	-	-	-	-	-	-	-	-	-	-	-
1924	-	-	-	-	-	-	-	-	-	-	-	-	-
1923	-	-	-	-	-	-	-	-	-	-	-	-	-
1922	-	-	-	-	-	-	-	-	-	-	-	-	-
1921	-	-	-	-	-	-	-	-	-	-	-	-	-
1920	-	-	-	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

TABLE 31.- 1/ CASES OF SELECTED COMMUNICABLE DISEASES BY PREFECTURE: JAPAN, 1920 - 1949 - Cont'd

5/ MALARIA

Year	All Japan	Hokkaido	Aomori	Iwate	Miyagi	Akita	Yamagata	Fukushima	Ibaraki	Tochigi	Gumma
*1949	3,732	48	54	17	15	16	31	25	38	20	11
1948	4,940	110	40	33	33	26	24	43	49	38	15
1947	11,841	296	181	183	25	186	118	249	316	115	89
1920 - 1946	Not Available.										

6/ JAPANESE "B" ENCEPHALITIS

Year	All Japan	Hokkaido	Aomori	Iwate	Miyagi	Akita	Yamagata	Fukushima	Ibaraki	Tochigi	Gumma
*1949	1,284	1	-	-	3	-	1	-	31	2	22
1948	7,208	7	46	140	165	139	152	66	335	88	143
1947	259	-	2	-	1	2	1	-	-	1	1
1920 - 1946	Not Available.										

See footnotes at end of table.

TABLE 31.- 1/ CASES OF SELECTED COMMUNICABLE DISEASES BY PREFECTURE: JAPAN, 1920 - 1949 - Cont'd

5/ MALARIA - Cont'd

Year	Saitama	Chiba	Tokyo	Kanagawa	Niigata	Toyama	Ishikawa	Fukui	Yamanashi	Nagano	Gifu	Shizuoka
*1949	34	41	131	39	46	23	18	32	14	24	26	24
1948	26	33	312	82	103	50	32	25	23	16	42	43
1947	66	116	765	452	304	167	60	73	68	186	38	208
1920 - 1946	Not Available.											

6/ JAPANESE 'B' ENCEPHALITIS - Cont'd

Year	Saitama	Chiba	Tokyo	Kanagawa	Niigata	Toyama	Ishikawa	Fukui	Yamanashi	Nagano	Gifu	Shizuoka
*1949	15	6	216	115	9	24	21	-	2	29	7	49
1948	299	337	1,969	589	239	105	81	65	111	191	107	377
1947	-	-	5	1	1	1	-	1	-	-	1	-
1920 - 1946	Not Available.											

See footnotes at end of table.

TABLE 31.-- 1/ CASES OF SELECTED COMMUNICABLE DISEASES BY PREFECTURE: JAPAN, 1920 - 1949 - Cont'd

5/ MALARIA - Cont'd

Year	Aichi	Mie	Shiga	Kyoto	Osaka	Hyogo	Nara	Wakayama	Tottori	Shimane	Okayama	Hiroshima
*1949	68	32	2,200	132	25	56	14	11	26	15	30	47
1948	49	51	2,258	86	55	66	31	20	47	34	51	110
1947	264	226	1,881	174	160	316	74	82	157	131	79	236
1920 - 1946	Not Available.											

6/ JAPANESE "B" ENCEPHALITIS

Year	Aichi	Mie	Shiga	Kyoto	Osaka	Hyogo	Nara	Wakayama	Tottori	Shimane	Okayama	Hiroshima
*1949	75	36	6	25	105	12	23	-	2	-	42	84
1948	389	84	62	34	100	136	15	18	27	18	69	34
1947	-	6	-	5	46	12	-	1	22	7	69	6
1920 - 1946	Not Available.											

See footnotes at end of table.

TABLE 31.- 1/ CASES OF SELECTED COMMUNICABLE DISEASES BY PREFECTURE: JAPAN, 1920 - 1949 - Cont'd

5/ MALARIA - Cont'd

Year	Yamaguchi	Tokushima	Kagawa	Ehime	Kochi	Fukuoka	Saga	Nagasaki	Kumamoto	Oita	Miyazaki	Kagoshima ^{2/}	Unknown
*1949	31	10	27	33	16	68	16	40	34	20	24	30	-
1948	47	22	24	107	21	246	41	68	63	54	23	158	-
1947	281	220	146	486	111	996	277	216	208	373	200	286	-
1920 - 1946	Not Available.												

6/ JAPANESE *B* ENCEPHALITIS

Year	Yamaguchi	Tokushima	Kagawa	Ehime	Kochi	Fukuoka	Saga	Nagasaki	Kumamoto	Oita	Miyazaki	Kagoshima ^{2/}	Unknown
*1949	76	9	12	17	10	15	19	4	86	11	26	37	-
1948	30	26	18	64	34	45	26	32	47	11	66	72	-
1947	-	1	31	16	13	1	-	1	2	1	1	-	-
1920 - 1946	Not Available.												

See footnotes next page.

TABLE 31.-- 1/ CASES OF SELECTED COMMUNICABLE DISEASES BY PREFECTURE: JAPAN, 1920 - 1949 - Cont'd

FOOTNOTES:

- Data for 1949 are provisional
- 1/ Date refer to cases of communicable disease among the civilian population of Japan Proper, 1900-1934 and 1937-1949. Totals for 1935 and 1936 include Okinawa, and prefectural data for those years are not available. Numbers for 1946-1949 represent compilations of weekly periods, while those for 1920-1945 inclusive are totals for calendar years.
- 2/ The unknown column contains corrections that cannot be allocated by prefecture.
- 3/ Diphtheria includes croup, 1920-1923.
- 4/ Dysentery includes ekiri, 1922-1949.
- 5/ Annual data for malaria are not available prior to 1947.
- 6/ Annual data for Japanese "B" encephalitis are not available prior to 1947.
- "NA" indicates that data are not available.
- A dash (-) indicates that no cases were reported.

Sources:

- 1920-1923: Statistics of Notifiable Infectious Diseases, Sanitary Bureau, Home Department, 1924.
- 1924-1934: Statistics of Notifiable Infectious Diseases, Sanitary Bureau, Home Department, 1937.
- 1935-1942: Annual Reports, Ministry of Welfare.
- 1943-1945: Unpublished Annual Reports, Ministry of Welfare.
- 1946-1949: Weekly Reports, Ministry of Welfare

TABLE 32. - 1/CASE RATES FOR SELECTED COMMUNICABLE DISEASES BY PREFECTURE: JAPAN, 1920-1949
TYPHOID FEVER

Year	All Japan	Hokkaido	Aomori	Iwate	Miyagi	Akita	Yamagata	Fukushima	Ibaraki	Tochigi	Gumma
* 1949	7.9	4.7	7.0	6.3	12.8	4.2	4.3	5.5	4.1	4.2	3.5
1948	11.8	9.8	11.1	6.8	12.8	7.0	5.6	9.0	7.7	8.8	7.9
1947	22.9	19.7	20.2	18.3	25.3	12.5	25.3	22.3	27.5	27.5	17.7
1946	59.2	73.4	72.4	58.1	61.8	36.8	73.3	93.7	52.1	100.5	44.5
1945	82.9	141.9	92.8	97.9	64.3	73.7	85.6	99.9	52.6	189.2	58.4
1944	79.1	104.6	66.2	45.9	77.9	54.1	77.4	62.7	34.3	84.7	29.6
1943	72.2	66.4	58.0	77.5	108.3	25.2	47.1	55.3	28.3	75.7	29.6
1942	49.3	41.7	60.2	64.4	92.4	37.4	33.7	45.1	22.0	42.0	22.1
1941	56.6	39.0	51.8	97.7	80.5	101.1	54.9	57.9	35.0	42.2	33.1
1940	57.4	45.8	69.6	107.2	100.1	110.4	99.7	51.2	42.6	51.5	25.8
1939	53.7	48.5	56.1	162.4	96.3	100.8	81.3	70.6	32.4	41.9	26.6
1938	60.0	43.8	54.7	130.0	164.8	71.4	79.3	78.8	38.1	59.5	58.7
1937	54.5	40.6	50.1	96.7	95.1	92.4	53.1	62.6	36.3	42.6	38.7
1936	53.5	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1935	55.3	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1934	62.7	61.7	67.2	164.5	166.0	91.0	62.7	69.7	42.5	36.6	27.4
1933	57.5	73.5	60.7	72.7	133.1	65.6	68.2	51.1	58.4	34.1	34.6
1932	53.9	60.2	49.7	78.7	123.0	69.5	67.2	60.5	42.0	49.9	48.2
1931	59.0	63.0	52.2	65.8	115.4	52.1	55.9	53.1	31.3	66.6	52.5
1930	64.8	90.4	107.5	86.1	126.0	78.6	82.1	74.6	32.7	90.7	62.7
1929	59.2	70.3	82.2	54.6	101.8	42.7	63.7	78.8	31.4	60.5	43.6
1928	67.6	94.9	68.7	91.5	92.9	49.1	73.0	72.6	50.7	73.2	51.4
1927	61.4	86.5	106.5	82.6	66.1	62.2	88.3	79.6	56.3	66.4	67.2
1926	72.9	70.8	129.1	53.2	82.7	86.8	80.1	76.0	51.7	45.7	47.6
1925	77.3	136.8	92.7	77.9	137.9	105.6	51.4	102.6	54.6	57.0	56.1
1924	109.1	86.2	75.3	85.8	156.8	69.3	63.6	123.4	72.8	77.8	53.7
1923	91.3	78.0	94.0	96.7	141.6	104.7	100.7	136.0	85.8	94.8	63.1
1922	82.1	101.1	167.0	148.1	114.7	196.9	132.4	173.1	77.1	77.1	74.3
1921	89.3	94.6	94.6	145.7	154.0	106.1	128.5	165.2	92.3	85.9	85.8
1920	97.1	136.4	98.8	145.1	220.4	153.4	137.1	162.3	69.8	91.9	75.1

See footnotes at end of table.

TABLE 32. - 1/CASE RATES FOR SELECTED COMMUNICABLE DISEASES BY PREFECTURE: JAPAN, 1920-1949 Cont'd
TYPHOID FEVER - Cont'd

Year	Saitama	Chiba	Tokyo	Kanagawa	Miicata	Toyama	Ishikawa	Fukui	Yamanashi	Nagano	Gifu	Shizuoka
*1949	9.1	7.1	16.1	11.5	12.4	8.9	4.8	18.1	1.3	5.4	13.4	10.1
1948	11.6	11.4	23.3	19.7	11.4	15.3	7.8	17.1	2.7	21.6	25.1	13.6
1947	23.8	19.3	27.8	30.1	25.7	37.8	21.0	22.4	16.8	16.2	40.3	27.6
1946	59.6	63.9	52.2	77.7	58.8	92.1	53.3	43.2	40.1	41.5	74.0	65.4
1945	109.6	90.4	35.5	135.0	131.3	130.8	60.1	56.6	145.6	102.0	181.2	129.5
1944	59.1	69.5	89.9	195.6	98.1	112.0	62.2	56.2	56.4	46.9	72.3	78.8
1943	65.4	60.5	92.8	182.7	65.5	101.3	83.2	70.5	30.0	50.9	79.4	74.3
1942	36.1	37.5	42.4	110.4	86.2	65.7	41.7	57.5	44.1	29.6	32.2	56.0
1941	47.8	38.0	46.6	93.4	64.8	62.8	53.8	71.2	82.1	29.5	38.0	61.2
1940	58.2	45.5	42.8	73.3	85.0	77.4	68.8	51.6	42.3	35.8	34.1	48.6
1939	27.6	42.0	32.7	62.2	72.6	61.4	54.2	20.2	41.4	32.3	33.7	78.1
1938	38.2	48.1	36.7	46.9	76.2	92.9	65.8	23.2	36.6	47.2	46.8	66.7
1937	39.1	41.9	32.2	46.5	48.8	55.1	39.8	39.0	70.9	31.9	68.4	48.9
1936	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1935	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1934	41.8	27.7	44.1	46.9	73.4	45.7	48.0	26.5	39.5	26.3	49.0	52.5
1933	34.2	23.8	54.8	55.9	62.3	31.4	36.6	35.3	54.7	33.0	38.2	48.7
1932	31.2	26.4	55.4	49.7	68.2	19.6	38.9	32.1	51.9	33.8	31.6	39.9
1931	39.3	22.4	75.0	67.7	60.7	32.6	55.7	38.0	42.5	38.9	36.0	48.2
1930	42.1	41.4	71.0	52.9	85.7	34.0	61.4	18.8	66.7	46.9	39.7	52.1
1929	27.4	27.7	75.9	44.1	38.3	40.0	25.4	18.8	50.3	60.1	40.2	57.4
1928	36.0	37.5	82.8	53.7	44.8	44.3	31.2	20.3	129.5	45.5	42.6	50.4
1927	29.6	23.4	82.3	75.6	51.8	20.4	30.1	53.4	65.4	48.3	40.5	58.7
1926	39.6	27.8	112.7	60.0	52.1	27.6	38.7	42.0	40.2	61.4	55.6	91.2
1925	36.3	41.4	126.7	70.6	52.7	49.1	38.0	19.2	82.2	50.0	44.9	74.7
1924	59.6	54.9	243.2	254.1	77.5	33.4	32.0	25.9	105.8	56.2	45.4	144.0
1923	64.2	59.5	192.1	235.9	58.8	50.8	48.4	50.9	65.6	72.8	61.9	109.6
1922	74.4	59.6	138.9	101.6	115.3	68.1	72.8	26.8	136.6	77.2	57.5	180.2
1921	86.2	33.9	134.5	167.1	83.5	27.9	52.8	38.1	138.1	81.2	66.5	162.9
1920	130.6	49.1	141.0	156.0	55.0	19.1	36.8	31.5	243.7	97.9	55.1	148.1

See footnotes at end of table.

TABLE 32. - 1/CASE RATES FOR SELECTED COMMUNICABLE DISEASES BY PREFECTURE: JAPAN, 1920-1949 Cont'd
TYPHOID FEVER - Cont'd

Year	Aichi	Mie	Shiga	Kyoto	Osaka	Hyogo	Nara	Wakayama	Tottori	Shimane	Okayama	Hiroshima
*1949	10.0	19.0	3.5	9.0	7.0	9.2	10.9	9.1	10.9	8.5	6.9	13.7
1948	13.7	20.5	6.1	15.3	10.1	12.2	15.4	12.6	15.8	13.6	10.9	14.0
1947	31.3	55.0	14.9	23.3	17.4	32.6	17.6	48.6	26.3	31.3	21.9	36.0
1946	42.6	80.5	38.8	82.0	43.3	87.5	66.4	84.6	84.3	84.2	77.2	86.0
1945	65.8	79.8	66.2	112.3	23.7	201.8	124.4	100.2	78.5	84.2	80.4	114.9
1944	65.2	77.9	47.8	84.4	83.7	232.7	80.0	74.0	52.4	58.6	52.4	130.4
1943	86.2	54.2	50.7	73.3	92.0	137.6	74.4	73.9	60.6	50.2	58.2	121.4
1942	48.8	35.3	40.7	41.1	52.1	199.5	53.5	76.0	47.1	51.5	61.5	85.3
1941	47.9	48.3	51.7	54.3	67.6	116.1	56.7	71.3	97.9	66.2	66.2	96.3
1940	47.5	50.3	53.6	51.6	71.0	101.1	44.8	60.9	62.4	37.9	62.4	105.5
1939	45.9	43.6	46.7	82.6	73.6	82.3	85.4	64.3	36.5	44.3	46.4	84.5
1938	50.4	46.7	44.4	75.2	87.4	100.7	67.8	76.9	73.4	31.4	57.1	99.7
1937	50.8	43.9	54.8	61.3	73.9	109.6	56.6	91.2	53.8	39.8	46.7	138.2
1936	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1935	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1934	80.2	49.1	61.5	97.9	118.7	93.6	47.4	52.7	64.3	31.2	51.7	89.3
1933	76.3	53.5	69.2	79.1	96.5	93.8	64.8	82.3	34.2	29.7	42.2	66.5
1932	71.1	38.0	46.6	61.6	82.8	73.5	38.4	55.0	29.4	49.2	58.8	60.6
1931	77.9	41.1	40.4	83.7	81.2	68.1	44.9	52.9	52.6	29.9	29.6	92.5
1930	67.8	43.9	43.8	63.2	72.9	75.3	38.7	49.7	42.3	107.2	45.9	80.4
1929	94.0	38.6	40.6	79.7	65.5	75.1	36.7	54.5	47.7	86.0	54.2	94.5
1928	91.8	50.6	54.6	92.9	94.9	95.9	64.5	61.7	76.1	50.3	47.3	79.2
1927	70.8	30.7	30.7	80.0	73.6	73.7	42.0	57.6	73.4	42.5	44.6	67.0
1926	74.3	47.0	42.7	109.8	75.6	75.2	39.1	55.7	89.8	58.1	65.4	50.2
1925	84.3	64.7	49.4	128.9	78.6	100.8	44.7	67.2	89.8	57.3	42.3	97.4
1924	94.3	56.2	37.2	198.1	118.1	99.0	59.1	53.8	118.9	63.7	45.2	184.0
1923	90.5	51.4	47.8	197.5	77.7	87.0	56.4	40.0	99.4	57.7	58.1	116.2
1922	76.1	25.3	25.3	159.7	103.6	86.2	45.5	38.6	104.6	57.5	78.3	72.2
1921	89.8	48.7	32.5	160.6	74.4	74.2	84.2	51.2	58.7	68.8	86.6	80.0
1920	89.1	51.3	31.8	138.3	76.1	103.0	92.5	41.6	79.8	103.8	108.6	152.5

See footnotes at end of table.

TABLE 32. - 1/CASE RATES FOR SELECTED COMMUNICABLE DISEASES BY PREFECTURE: JAPAN, 1920-1949 - Cont'd
TYPHOID FEVER - Cont'd

Year	Yamaguchi	Tokushima	Kagawa	Ehime	Kochi	Tokuoka	Sara	Nazasaki	Kusumoto	Oita	Miyazaki	Kagoshima
1949	3.3	7.4	4.4	3.8	11.3	5.1	4.6	3.7	1.4	2.6	3.2	0.7
1948	5.9	14.7	5.7	11.2	18.5	5.9	5.9	4.4	1.4	9.4	8.4	1.1
1947	7.9	30.0	21.7	15.0	49.2	11.2	9.0	7.2	1.4	8.7	16.2	1.7
1946	26.8	93.0	67.0	36.5	94.2	43.5	31.2	30.5	13.7	20.0	50.0	11.7
1945	35.4	80.7	89.5	28.2	109.2	50.7	49.2	18.6	10.0	27.0	18.4	1.5
1944	36.0	107.6	83.7	29.8	132.2	45.2	77.5	23.2	7.4	31.5	20.2	5.9
1943	74.6	65.0	62.9	43.1	90.7	46.0	33.4	32.9	6.5	75.3	27.4	18.4
1942	23.5	61.5	65.6	43.6	77.9	35.2	34.4	12.0	6.7	33.6	17.1	10.4
1941	29.0	65.1	114.4	46.9	78.7	41.9	77.7	38.2	12.3	24.8	20.8	7.4
1940	30.8	69.8	92.2	81.2	81.3	42.0	47.0	43.4	14.4	24.8	26.8	5.8
1939	20.5	63.1	68.9	62.8	91.1	45.5	28.4	44.7	10.2	24.5	7.5	7.3
1938	29.8	65.0	63.7	75.0	117.1	65.1	43.6	30.1	19.1	22.3	12.3	7.1
1937	34.1	49.5	74.6	70.3	74.5	54.3	53.7	32.5	22.3	25.9	15.7	9.9
1936	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1935	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1934	27.9	50.9	92.7	79.6	51.4	73.2	36.5	35.0	28.7	56.8	27.8	23.1
1933	29.9	25.9	56.8	76.9	56.6	65.8	47.1	32.6	25.4	40.1	20.5	14.2
1932	31.2	42.3	56.3	49.7	54.2	61.1	78.7	57.3	27.0	40.5	11.5	17.7
1931	39.1	40.6	77.9	50.3	55.1	60.5	21.2	164.1	52.5	37.9	14.7	18.1
1930	35.5	52.3	79.8	49.5	61.0	85.6	39.5	65.3	66.7	37.7	24.3	37.3
1929	39.1	46.1	100.4	52.6	55.1	86.5	26.0	47.6	63.3	35.4	25.4	22.0
1928	40.2	36.7	92.3	62.2	76.4	83.3	35.0	77.3	116.3	37.8	17.3	17.9
1927	44.7	22.5	66.1	40.8	53.3	99.8	30.4	61.8	86.0	34.6	14.6	11.6
1926	63.8	21.4	64.6	42.6	113.5	158.3	15.1	96.6	202.6	38.2	17.4	22.7
1925	75.0	44.4	74.7	58.5	63.3	111.5	31.2	81.4	77.5	43.6	21.8	28.9
1924	148.4	62.3	110.4	50.4	47.6	105.6	39.2	63.9	86.2	55.0	19.8	44.2
1923	158.6	28.3	72.9	40.0	68.7	86.1	27.4	62.9	87.1	144.1	33.7	21.1
1922	74.5	19.6	111.0	42.4	67.7	95.0	15.9	79.1	57.6	37.9	27.8	19.0
1921	57.6	17.0	119.4	49.3	81.3	122.3	10.8	83.7	92.3	40.6	21.8	19.5
1920	59.0	25.7	95.2	44.5	72.4	123.1	11.3	47.2	73.9	41.5	21.2	23.9

See footnotes at end of table.

TABLE 32. - 1/CASE RATES FOR SELECTED COMMUNICABLE DISEASES BY PREFECTURE: JAPAN, 1920-1949 - Cont'd
PARATYPHOID FEVER

Year	All Japan	Hokkaido	Aomori	Iwate	Miyagi	Akita	Yamagata	Fukushima	Ibaraki	Tochigi	Gumma
*1949	2.7	2.0	3.2	2.5	7.1	1.2	1.5	6.5	2.3	1.4	1.9
1948	3.6	3.1	4.3	4.2	8.6	0.8	2.1	3.9	4.1	3.3	3.5
1947	6.1	6.0	4.6	5.2	16.7	3.7	8.0	5.1	9.6	6.7	7.7
1946	12.1	25.5	15.4	9.1	18.1	4.0	32.6	15.8	13.2	18.5	7.1
1945	14.7	50.0	6.7	14.0	31.4	14.4	16.0	17.0	15.2	27.7	23.1
1944	20.4	23.7	13.1	12.0	43.6	24.4	9.7	23.9	18.3	21.7	7.1
1943	17.0	13.9	9.4	16.4	70.9	10.3	13.6	17.8	14.8	18.9	7.9
1942	8.6	10.7	14.9	10.2	31.3	9.5	9.3	9.0	7.5	30.7	7.3
1941	8.7	7.9	7.0	9.8	16.1	7.8	15.8	13.7	37.9	8.2	10.1
1940	8.8	9.0	16.3	18.8	7.3	14.3	22.5	11.4	10.8	8.6	7.2
1939	7.4	7.6	8.2	12.9	5.2	8.3	26.6	8.1	9.3	7.5	3.8
1938	8.7	9.3	8.6	9.3	4.7	8.2	15.8	10.3	12.4	9.4	12.1
1937	6.4	7.5	9.5	15.9	8.0	9.2	23.3	7.0	3.3	8.1	4.2
1936	6.9	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1935	6.2	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1934	6.6	9.6	4.9	9.9	11.4	15.0	13.2	17.7	6.7	10.6	5.5
1933	7.9	26.5	4.6	7.5	3.2	5.2	14.0	8.3	28.2	12.3	10.3
1932	7.1	27.2	9.6	10.9	1.9	4.5	29.7	6.2	3.7	11.9	13.4
1931	6.2	13.2	5.4	2.5	4.4	2.8	25.6	5.6	4.4	11.5	10.7
1930	7.0	15.5	8.3	5.5	3.7	4.4	38.8	5.6	5.7	13.2	20.7
1929	6.7	11.6	9.0	5.5	8.5	3.1	18.5	13.3	8.6	13.5	13.3
1928	7.9	19.6	5.3	8.7	5.1	5.1	17.7	7.6	6.3	18.9	14.1
1927	7.7	22.6	6.2	6.6	5.1	4.6	6.6	9.3	13.1	10.7	21.7
1926	7.3	18.7	4.5	8.1	6.5	8.1	9.3	8.8	6.3	11.2	14.2
1925	8.5	29.1	6.0	12.7	7.3	6.1	5.8	9.4	11.6	14.6	17.5
1924	9.1	15.2	4.8	7.7	5.8	5.5	6.8	9.4	9.9	16.5	29.5
1923	9.2	14.6	12.3	17.8	14.0	9.1	8.8	8.1	8.3	14.8	30.0
1922	12.5	20.0	22.3	16.0	7.1	16.8	10.1	16.4	13.6	18.3	19.6
1921	11.2	14.8	11.9	13.4	10.1	12.8	20.0	14.0	6.4	12.4	25.7
1920	13.9	30.0	11.9	15.7	11.5	11.4	10.9	18.6	7.8	21.6	67.5

See footnotes at end of table.

TABLE 32. - 1/CASE RATES FOR SELECTED COMMUNICABLE DISEASES BY PREFECTURE: JAPAN, 1920-1949 - Cont'd
PARATYPHOID FEVER - Cont'd

Year	Saitama	Chiba	Tokyo	Kanagawa	Miigata	Toyama	Ishikawa	Fukui	Yamanashi	Nagano	Gifu	Shizuoka
1949	2.5	2.0	8.8	2.9	3.2	4.2	3.2	4.0	0.6	2.3	1.7	4.2
1948	2.1	2.8	12.6	5.1	4.5	2.8	1.8	3.0	1.5	3.4	5.3	6.9
1947	4.7	6.3	9.8	7.3	7.8	11.7	5.4	5.8	6.1	6.7	9.3	6.8
1946	8.9	8.7	16.0	14.7	13.9	10.4	7.4	8.2	12.2	16.4	13.4	20.1
1945	18.6	10.0	13.8	13.3	20.9	19.6	17.7	7.7	49.6	45.6	8.9	21.7
1944	14.3	9.0	74.1	28.6	14.1	41.7	8.6	11.3	8.9	16.9	14.5	17.8
1943	9.5	10.5	45.8	24.9	19.7	12.6	7.8	9.6	4.8	20.2	13.6	17.4
1942	11.9	2.8	12.1	10.0	16.8	15.2	5.6	18.8	3.8	10.3	2.7	5.0
1941	8.6	2.0	11.7	11.3	9.5	4.3	4.5	10.2	13.6	15.8	4.5	9.7
1940	19.1	4.5	9.3	11.5	15.6	0.9	5.4	6.8	15.3	19.7	6.1	8.3
1939	6.1	3.9	6.1	8.4	15.1	2.5	8.2	5.7	14.3	11.6	12.9	19.6
1938	9.2	4.0	6.7	5.1	12.0	0.6	19.5	9.4	14.2	21.8	5.2	10.5
1937	5.4	4.2	4.6	7.5	6.3	1.5	4.7	5.6	3.1	10.4	11.2	8.2
1936	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1935	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1934	8.3	1.6	9.2	6.4	12.6	2.4	1.4	1.3	5.0	7.2	2.4	6.4
1933	6.1	4.7	8.5	5.0	14.8	2.0	3.0	1.4	12.9	6.0	4.8	11.3
1932	5.1	2.2	10.5	4.3	9.4	0.9	2.4	2.4	3.9	5.6	1.8	7.0
1931	4.9	2.2	10.6	4.8	10.1	2.4	2.9	2.1	1.1	6.7	1.5	7.3
1930	9.0	2.7	8.8	7.4	9.8	2.8	2.4	1.3	4.1	8.5	2.6	8.8
1929	11.2	2.6	7.3	6.8	8.5	3.7	3.7	2.0	6.6	8.2	1.3	13.3
1928	11.7	4.5	8.3	18.1	9.2	3.9	3.2	2.8	3.6	8.9	3.3	9.0
1927	8.2	3.6	8.3	9.2	9.8	1.8	4.4	3.0	3.3	16.3	4.5	12.4
1926	7.3	3.4	11.0	6.5	8.0	1.5	19.9	2.7	5.1	10.6	3.0	8.7
1925	17.6	7.5	9.7	6.7	11.8	3.3	3.5	2.0	1.8	6.2	1.9	6.9
1924	9.5	7.2	13.5	11.1	13.1	4.7	2.1	5.5	5.1	9.9	3.2	15.5
1923	14.0	7.7	12.3	13.6	13.5	6.8	3.6	8.5	7.2	8.3	5.2	15.7
1922	26.6	9.8	10.0	14.7	32.8	17.7	8.1	11.9	16.7	8.6	7.1	16.1
1921	15.3	7.0	10.5	17.4	24.2	1.2	4.1	7.5	19.0	10.1	3.5	29.6
1920	27.8	7.3	11.6	20.6	19.5	7.7	6.8	2.8	19.7	9.0	9.8	39.0

See footnotes at end of table.

TABLE 32. - 1/CASE RATES FOR SELECTED COMMUNICABLE DISEASES BY PREFECTURES: JAPAN, 1920-1949 - Cont'd
PARATYPHOID FEVER - Cont'd

Year	Aichi	Mie	Shiga	Kyoto	Osaka	Hyogo	Nara	Wakayama	Tottori	Shimane	Okayama	Hiroshima
*1949	2.7	3.6	2.1	3.2	1.4	1.0	1.5	3.3	2.5	1.7	1.2	3.9
1948	2.9	4.4	1.7	3.8	1.9	1.1	1.8	1.6	2.2	7.3	1.0	3.1
1947	6.4	9.2	3.4	5.6	8.3	3.6	2.1	6.8	5.6	12.8	1.3	8.2
1946	8.2	9.6	3.6	10.2	6.7	9.5	7.4	9.3	14.1	16.6	2.4	12.4
1945	6.3	4.2	8.8	7.2	2.1	20.8	13.2	7.7	10.5	18.3	2.7	11.9
1944	10.9	5.4	6.7	4.9	11.5	25.2	5.5	6.6	9.9	-	2.8	24.9
1943	11.7	6.0	5.7	12.6	15.3	20.7	2.4	5.3	7.4	6.4	3.8	21.8
1942	8.2	3.9	4.0	6.1	4.2	5.6	1.8	10.8	11.0	6.6	5.6	12.5
1941	6.7	7.1	8.2	8.3	4.7	11.1	5.2	7.0	5.5	7.6	2.8	9.0
1940	6.3	5.0	4.1	11.8	6.4	7.3	5.3	5.7	4.5	6.1	5.3	8.0
1939	6.3	4.9	2.5	6.8	4.3	5.2	1.5	6.6	1.9	7.3	2.4	7.7
1938	7.0	6.2	3.1	8.9	4.5	9.2	1.2	5.9	5.6	2.7	1.2	9.8
1937	6.5	3.7	5.0	3.6	7.2	6.2	1.7	3.9	3.0	6.1	3.0	16.5
1936	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1935	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1934	5.3	5.7	3.9	5.1	6.9	6.2	1.3	4.2	4.8	8.4	1.8	2.1
1933	9.7	3.2	3.2	6.8	8.5	10.2	1.5	8.0	10.6	2.0	1.4	2.4
1932	5.5	2.6	6.2	8.2	6.8	5.5	0.7	7.4	2.0	5.4	1.7	4.0
1931	6.2	2.3	4.9	11.8	6.6	5.1	3.0	6.6	7.7	7.1	1.5	2.5
1930	4.8	3.9	5.8	14.0	5.8	6.7	1.7	10.6	2.7	5.9	2.3	3.2
1929	5.3	2.5	3.1	7.7	5.8	8.0	1.2	11.7	3.3	6.8	1.6	3.1
1928	5.7	3.7	3.7	8.8	8.4	13.2	1.5	5.5	7.2	11.3	3.6	2.8
1927	5.7	3.9	3.3	10.0	7.9	11.2	4.7	8.8	14.3	10.3	3.1	4.2
1926	5.8	6.2	3.0	16.1	9.2	17.3	5.8	5.5	9.6	4.1	5.4	5.7
1925	5.8	5.8	1.7	29.4	11.5	11.2	6.2	2.4	5.1	4.2	2.8	3.0
1924	4.9	2.9	3.7	19.7	15.7	10.8	5.4	8.5	5.2	8.0	3.9	2.3
1923	4.9	2.7	4.6	19.8	10.7	9.6	9.3	4.6	5.9	9.9	3.1	2.6
1922	6.6	8.1	4.9	34.0	16.5	21.9	16.2	6.7	5.7	15.0	2.8	3.1
1921	7.7	8.8	4.5	26.2	15.3	13.4	27.2	6.5	3.9	10.6	4.5	3.4
1920	12.1	9.1	7.7	24.2	14.8	17.8	22.0	9.7	2.2	7.8	7.1	6.9

See footnotes at end of table.

TABLE 35. - 1/CASE RATES FOR SELECTED COMMUNICABLE DISEASES BY PREFECTURE: JAPAN, 1920-1949 - Cont'd
PARATYPHOID FEVER - Cont'd

Year	Yamanuchi	Tokushima	Kagawa	Ehime	Kochi	Fukuoka	Saga	Nagasaki	Kumamoto	Oita	Miyazaki	Kagoshima
1949	0.6	1.5	0.5	1.1	0.9	1.4	0.6	0.4	1.8	0.8	1.4	0.3
1948	0.8	0.9	3.6	2.2	4.1	1.7	1.8	1.2	1.2	1.9	1.8	0.5
1947	2.1	4.5	7.1	2.3	5.1	2.0	2.6	2.6	1.4	1.0	4.7	1.0
1946	6.2	6.0	23.7	4.1	8.2	11.4	13.8	13.2	2.9	4.5	11.8	3.5
1945	3.9	11.7	6.6	9.2	7.6	10.6	3.0	2.4	-	3.3	13.9	0.9
1944	7.3	12.5	6.9	12.0	3.2	6.3	9.2	3.6	1.0	6.0	3.6	19.1
1943	4.8	18.1	7.7	3.9	5.1	11.1	2.4	3.6	1.8	8.2	6.1	13.6
1942	5.1	11.9	5.1	4.1	7.2	7.3	0.4	3.0	2.0	3.3	1.8	1.0
1941	2.3	5.5	3.8	4.5	12.3	6.4	1.9	2.7	0.9	7.6	5.6	2.1
1940	3.4	7.0	3.7	2.5	6.2	6.5	2.4	36.7	3.1	1.6	1.1	1.3
1939	2.1	4.6	3.3	2.8	6.4	14.8	1.8	10.9	1.8	7.1	1.4	0.5
1938	1.9	6.8	3.1	1.8	3.8	33.4	4.0	5.9	3.5	19.3	1.0	1.1
1937	3.4	3.9	3.0	1.8	5.9	7.1	3.6	1.6	5.5	4.5	2.1	1.2
1936	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1935	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1934	5.7	24.0	2.3	3.4	2.6	3.7	3.9	1.0	2.6	1.4	2.8	3.8
1933	3.9	7.1	3.1	4.2	2.2	6.1	1.6	7.9	2.9	1.1	0.5	0.6
1932	11.1	5.1	3.3	3.0	1.0	5.4	2.6	2.2	11.0	2.1	0.5	0.1
1931	3.7	2.3	2.9	2.3	1.8	9.1	0.6	2.2	5.7	1.4	0.9	1.3
1930	4.9	3.3	2.6	1.5	2.2	4.9	3.9	2.8	4.2	2.9	2.4	0.1
1929	6.5	5.2	6.8	1.4	1.3	5.0	3.0	3.1	3.7	4.8	1.8	0.3
1928	6.2	9.5	5.7	3.9	3.0	5.7	4.9	2.6	9.0	4.1	5.4	1.2
1927	2.7	6.1	9.0	4.5	3.3	2.9	2.9	3.2	3.5	1.6	1.4	0.2
1926	3.2	3.4	5.5	2.3	1.6	5.1	2.9	3.2	2.8	1.6	1.9	0.2
1925	3.7	9.3	4.4	1.1	2.9	5.3	6.1	5.4	2.3	3.2	0.7	0.5
1924	6.3	16.2	4.9	1.6	4.2	5.2	9.1	4.7	5.9	10.1	1.1	0.4
1923	6.5	11.3	3.6	1.6	2.4	4.3	4.8	5.5	7.1	4.2	1.0	0.7
1922	7.8	9.2	6.0	2.7	2.1	5.9	4.9	7.4	6.0	5.6	2.4	1.1
1921	4.1	10.3	3.3	1.9	2.7	9.4	0.7	4.7	10.8	2.6	4.4	1.1
1920	4.7	6.1	1.8	2.1	6.3	7.9	1.6	5.5	6.2	3.8	5.1	2.5

See footnotes at end of table.

TABLE 32. - 1/CASE RATES FOR SELECTED COMMUNICABLE DISEASES BY PREFECTURE: JAPAN, 1920-1949 - Cont'd
PLAGUE

Year	All Japan	Hokkaido	Aomori	Iwate	Miyagi	Akita	Yamagata	Fukushima	Ibaraki	Tochigi	Gumma
* 1949	-	-	-	-	-	-	-	-	-	-	-
1948	-	-	-	-	-	-	-	-	-	-	-
1947	-	-	-	-	-	-	-	-	-	-	-
1946	-	-	-	-	-	-	-	-	-	-	-
1945	-	-	-	-	-	-	-	-	-	-	-
1944	-	-	-	-	-	-	-	-	-	-	-
1943	-	-	-	-	-	-	-	-	-	-	-
1942	-	-	-	-	-	-	-	-	-	-	-
1941	-	-	-	-	-	-	-	-	-	-	-
1940	-	-	-	-	-	-	-	-	-	-	-
1939	-	-	-	-	-	-	-	-	-	-	-
1938	-	-	-	-	-	-	-	-	-	-	-
1937	-	-	-	-	-	-	-	-	-	-	-
1936	-	-	-	-	-	-	-	-	-	-	-
1935	-	-	-	-	-	-	-	-	-	-	-
1934	-	-	-	-	-	-	-	-	-	-	-
1933	-	-	-	-	-	-	-	-	-	-	-
1932	-	-	-	-	-	-	-	-	-	-	-
1931	-	-	-	-	-	-	-	-	-	-	-
1930	-	-	-	-	-	-	-	-	-	-	-
1929	-	-	-	-	-	-	-	-	-	-	-
1928	-	-	-	-	-	-	-	-	-	-	-
1927	-	-	-	-	-	-	-	-	-	-	-
1926	0.0	-	-	-	-	-	-	-	-	-	-
1925	-	-	-	-	-	-	-	-	-	-	-
1924	0.0	-	-	-	-	-	-	-	-	-	-
1923	0.0	-	-	-	-	-	-	-	-	-	-
1922	0.2	-	-	-	-	-	-	-	-	-	-
1921	0.0	-	-	-	-	-	-	-	-	-	-
1920	0.0	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

TABLE 32. 1/CASE RATES FOR SELECTED COMMUNICABLE DISEASES BY PREFECTURE: JAPAN, 1920-1949 - Cont'd
PLAGUE - Cont'd

Year	Saitama	Chiba	Tokyo	Kanagawa	Niigata	Toyama	Ishikawa	Fukui	Yamanashi	Nagano	Gifu	Shizuoka
*1949	-	-	-	-	-	-	-	-	-	-	-	-
1948	-	-	-	-	-	-	-	-	-	-	-	-
1947	-	-	-	-	-	-	-	-	-	-	-	-
1946	-	-	-	-	-	-	-	-	-	-	-	-
1945	-	-	-	-	-	-	-	-	-	-	-	-
1944	-	-	-	-	-	-	-	-	-	-	-	-
1943	-	-	-	-	-	-	-	-	-	-	-	-
1942	-	-	-	-	-	-	-	-	-	-	-	-
1941	-	-	-	-	-	-	-	-	-	-	-	-
1940	-	-	-	-	-	-	-	-	-	-	-	-
1939	-	-	-	-	-	-	-	-	-	-	-	-
1938	-	-	-	-	-	-	-	-	-	-	-	-
1937	-	-	-	-	-	-	-	-	-	-	-	-
1936	-	-	-	-	-	-	-	-	-	-	-	-
1935	-	-	-	-	-	-	-	-	-	-	-	-
1934	-	-	-	-	-	-	-	-	-	-	-	-
1933	-	-	-	-	-	-	-	-	-	-	-	-
1932	-	-	-	-	-	-	-	-	-	-	-	-
1931	-	-	-	-	-	-	-	-	-	-	-	-
1930	-	-	-	-	-	-	-	-	-	-	-	-
1929	-	-	-	-	-	-	-	-	-	-	-	-
1928	-	-	-	-	-	-	-	-	-	-	-	-
1927	-	-	-	-	-	-	-	-	-	-	-	-
1926	-	-	-	0.6	-	-	-	-	-	-	-	-
1925	-	-	-	-	-	-	-	-	-	-	-	-
1924	-	-	-	-	-	-	-	-	-	-	-	0.2
1923	-	-	-	-	-	-	-	-	-	-	-	-
1922	-	-	-	-	-	-	-	-	-	-	-	-
1921	-	-	-	-	-	-	-	-	-	-	-	-
1920	-	-	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

TABLE 32. - 1/CASE RATES FOR SELECTED COMMUNICABLE DISEASES BY PREFECTURE: JAPAN, 1920-1949 - Cont'd
PLAGUE - Cont'd

Year	Aichi	Mie	Shiga	Kyoto	Osaka	Hyogo	Nara	Wakayama	Tottori	Shimane	Okayama	Hiroshima
1949	-	-	-	-	-	-	-	-	-	-	-	-
1948	-	-	-	-	-	-	-	-	-	-	-	-
1947	-	-	-	-	-	-	-	-	-	-	-	-
1946	-	-	-	-	-	-	-	-	-	-	-	-
1945	-	-	-	-	-	-	-	-	-	-	-	-
1944	-	-	-	-	-	-	-	-	-	-	-	-
1943	-	-	-	-	-	-	-	-	-	-	-	-
1942	-	-	-	-	-	-	-	-	-	-	-	-
1941	-	-	-	-	-	-	-	-	-	-	-	-
1940	-	-	-	-	-	-	-	-	-	-	-	-
1939	-	-	-	-	-	-	-	-	-	-	-	-
1938	-	-	-	-	-	-	-	-	-	-	-	-
1937	-	-	-	-	-	-	-	-	-	-	-	-
1936	-	-	-	-	-	-	-	-	-	-	-	-
1935	-	-	-	-	-	-	-	-	-	-	-	-
1934	-	-	-	-	-	-	-	-	-	-	-	-
1933	-	-	-	-	-	-	-	-	-	-	-	-
1932	-	-	-	-	-	-	-	-	-	-	-	-
1931	-	-	-	-	-	-	-	-	-	-	-	-
1930	-	-	-	-	-	-	-	-	-	-	-	-
1929	-	-	-	-	-	-	-	-	-	-	-	-
1928	-	-	-	-	-	-	-	-	-	-	-	-
1927	-	-	-	-	-	-	-	-	-	-	-	-
1926	-	-	-	-	-	-	-	-	-	-	-	-
1925	-	-	-	-	-	-	-	-	-	-	-	-
1924	-	-	-	-	-	0.1	-	-	-	-	-	-
1923	-	-	-	-	-	-	-	-	-	-	-	-
1922	-	-	-	-	2.9	-	0.4	-	-	-	-	-
1921	-	-	-	-	-	0.1	-	-	-	-	-	-
1920	-	-	-	-	-	-	-	2.9	-	-	-	-

See footnotes at end of table.

TABLE 32. - 1/CASE RATES FOR SELECTED COMMUNICABLE DISEASES BY PREFECTURE: JAPAN, 1920-1949 - Cont'd
PLAGUE - Cont'd

Year	Yamaguchi	Tokushima	Kagawa	Ehime	Kochi	Fukuoka	Saga	Nagasaki	Kumamoto	Oita	Miyazaki	Kagoshima
*1949	-	-	-	-	-	-	-	-	-	-	-	-
1948	-	-	-	-	-	-	-	-	-	-	-	-
1947	-	-	-	-	-	-	-	-	-	-	-	-
1946	-	-	-	-	-	-	-	-	-	-	-	-
1945	-	-	-	-	-	-	-	-	-	-	-	-
1944	-	-	-	-	-	-	-	-	-	-	-	-
1943	-	-	-	-	-	-	-	-	-	-	-	-
1942	-	-	-	-	-	-	-	-	-	-	-	-
1941	-	-	-	-	-	-	-	-	-	-	-	-
1940	-	-	-	-	-	-	-	-	-	-	-	-
1939	-	-	-	-	-	-	-	-	-	-	-	-
1938	-	-	-	-	-	-	-	-	-	-	-	-
1937	-	-	-	-	-	-	-	-	-	-	-	-
1936	-	-	-	-	-	-	-	-	-	-	-	-
1935	-	-	-	-	-	-	-	-	-	-	-	-
1934	-	-	-	-	-	-	-	-	-	-	-	-
1933	-	-	-	-	-	-	-	-	-	-	-	-
1932	-	-	-	-	-	-	-	-	-	-	-	-
1931	-	-	-	-	-	-	-	-	-	-	-	-
1930	-	-	-	-	-	-	-	-	-	-	-	-
1929	-	-	-	-	-	-	-	-	-	-	-	-
1928	-	-	-	-	-	-	-	-	-	-	-	-
1927	-	-	-	-	-	-	-	-	-	-	-	-
1926	-	-	-	-	-	-	-	-	-	-	-	-
1925	-	-	-	-	-	-	-	-	-	-	-	-
1924	-	-	-	-	0.1	-	-	-	-	-	-	-
1923	-	-	-	-	5.2	-	-	-	-	-	-	-
1922	-	-	-	-	-	-	-	-	-	-	-	-
1921	-	-	-	-	-	-	-	-	-	-	-	-
1920	-	-	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

TABLE 32. - 1/CASE RATES FOR SELECTED COMMUNICABLE DISEASES BY PREFECTURE: JAPAN, 1920-1949 - Cont'd
CHOLERA

Year	All Japan	Hokkaido	Aomori	Iwate	Miyagi	Akita	Yamagata	Fukushima	Ibaraki	Tochigi	Gumma
* 1949	-	-	-	-	-	-	-	-	-	-	-
1948	-	-	-	-	-	-	-	-	-	-	-
1947	-	-	-	-	-	-	-	-	-	-	-
1946	1.6	-	-	-	-	-	-	0.3	-	-	-
1945	-	-	-	-	-	-	-	-	-	-	-
1944	-	-	-	-	-	-	-	-	-	-	-
1947	-	-	-	-	-	-	-	-	-	-	-
1942	-	-	-	-	-	-	-	-	-	-	-
1941	-	-	-	-	-	-	-	-	-	-	-
1940	-	-	-	-	-	-	-	-	-	-	-
1939	-	-	-	-	-	-	-	-	-	-	-
1938	0.0	-	-	-	-	-	-	-	-	-	-
1937	0.1	-	-	-	-	-	-	-	-	-	-
1936	-	-	-	-	-	-	-	-	-	-	-
1935	-	-	-	-	-	-	-	-	-	-	-
1934	-	-	-	-	-	-	-	-	-	-	-
1933	-	-	-	-	-	-	-	-	-	-	-
1932	0.0	-	-	-	-	-	-	-	-	-	-
1931	-	-	-	-	-	-	-	-	-	-	-
1930	-	-	-	-	-	-	-	-	-	-	-
1929	0.3	-	-	-	-	-	-	-	-	-	-
1928	0.0	-	-	-	-	-	-	-	-	-	-
1927	0.0	-	-	-	-	-	-	-	-	-	-
1926	0.0	-	-	-	-	-	-	-	-	-	-
1925	1.1	-	-	-	-	-	-	-	-	0.1	-
1924	-	-	-	-	-	-	-	-	-	-	-
1923	0.0	-	-	-	-	-	-	-	-	-	-
1922	1.3	-	-	-	-	-	-	-	-	-	-
1921	0.1	-	-	-	-	-	-	-	-	-	-
1920	9.0	-	-	-	-	-	-	-	-	-	-
1924	-	-	-	-	-	-	-	-	-	-	-
1923	0.0	-	-	-	-	-	-	-	-	-	-
1922	1.3	-	-	-	-	-	-	-	-	-	-
1921	0.1	-	-	-	-	-	-	-	-	-	-
1920	9.0	-	-	-	-	-	-	-	-	-	-
1924	-	-	-	-	-	-	-	-	-	-	-
1923	0.0	-	-	-	-	-	-	-	-	-	-
1922	1.3	-	-	-	-	-	-	-	-	-	-
1921	0.1	-	-	-	-	-	-	-	-	-	-
1920	9.0	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

TABLE 32. - 1/CASE RATES FOR SELECTED COMMUNICABLE DISEASES BY PREFECTURE: JAPAN, 1920-1949 - Cont'd
CHOLERA - Cont'd

Year	Saitama	Chiba	Tokyo	Kanagawa	Miura	Toyama	Ishikawa	Fukui	Yamanashi	Nagano	Gifu	Shizuoka
1949	-	-	-	-	-	-	-	-	-	-	-	-
1948	-	-	-	-	-	-	-	-	-	-	-	-
1947	-	-	-	-	-	-	-	-	-	-	-	-
1946	0.0	1.6	0.3	1.4	1.1	8.0	2.4	2.0	-	0.1	-	0.1
1945	-	-	-	-	-	-	-	-	-	-	-	-
1944	-	-	-	-	-	-	-	-	-	-	-	-
1943	-	-	-	-	-	-	-	-	-	-	-	-
1942	-	-	-	-	-	-	-	-	-	-	-	-
1941	-	-	-	-	-	-	-	-	-	-	-	-
1940	-	-	-	-	-	-	-	-	-	-	-	-
1939	-	-	-	-	-	-	-	-	-	-	-	-
1938	-	-	-	-	-	-	-	-	-	-	-	-
1937	-	-	0.0	-	-	-	-	-	-	-	-	-
1936	-	-	-	-	-	-	-	-	-	-	-	-
1935	-	-	-	-	-	-	-	-	-	-	-	-
1934	-	-	-	-	-	-	-	-	-	-	-	-
1933	-	-	-	-	-	-	-	-	-	-	-	-
1932	-	-	0.1	-	-	-	-	-	-	-	-	-
1931	-	-	-	-	-	-	-	-	-	-	-	-
1930	-	-	-	-	-	-	-	-	-	-	-	-
1929	-	-	-	-	-	-	-	-	-	-	-	-
1928	-	-	-	-	-	-	-	-	-	-	-	-
1927	-	-	-	-	-	-	-	-	-	-	-	-
1926	-	-	-	0.1	-	-	-	-	-	-	-	-
1925	0.3	3.3	1.8	6.8	-	-	-	-	-	-	-	0.4
1924	-	-	-	-	-	-	-	-	-	-	-	-
1923	-	-	-	-	-	-	-	-	-	-	-	-
1922	0.1	12.6	4.7	1.4	-	-	-	-	-	-	-	0.6
1921	-	-	-	0.1	-	-	-	-	-	-	-	-
1920	-	-	0.1	0.2	-	30.5	-	-	-	-	-	-

See footnotes at end of table.

TABLE 32. - 1/CASE RATES FOR SELECTED COMMUNICABLE DISEASES BY PREFECTURE: JAPAN, 1920-1949 - Cont'd
CHOLERA - Cont'd

Year	Aichi	Mie	Shiga	Kyoto	Osaka	Hyogo	Nara	Wakayama	Tottori	Shimane	Okayama	Firoshima
*1949	-	-	-	-	-	-	-	-	-	-	-	-
1948	-	-	-	-	-	-	-	-	-	-	-	-
1947	-	-	-	-	-	-	-	-	-	-	-	-
1946	0.3	0.1	-	1.0	2.3	0.5	-	0.6	1.0	1.9	2.1	8.6
1945	-	-	-	-	-	-	-	-	-	-	-	-
1944	-	-	-	-	-	-	-	-	-	-	-	-
1943	-	-	-	-	-	-	-	-	-	-	-	-
1942	-	-	-	-	-	-	-	-	-	-	-	-
1941	-	-	-	-	-	-	-	-	-	-	-	-
1940	-	-	-	-	-	-	-	-	-	-	-	-
1939	-	-	-	-	-	-	-	-	-	-	-	-
1938	-	-	-	-	-	-	-	-	-	-	0.7	0.3
1937	-	-	-	-	-	0.0	-	-	-	-	0.4	2.2
1936	-	-	-	-	-	-	-	-	-	-	-	-
1935	-	-	-	-	-	-	-	-	-	-	-	-
1934	-	-	-	-	-	-	-	-	-	-	-	-
1933	-	-	-	-	-	-	-	-	-	-	-	-
1932	-	-	-	-	-	-	-	-	-	-	-	-
1931	-	-	-	-	-	-	-	-	-	-	-	-
1930	-	-	-	-	-	-	-	-	-	-	-	-
1929	-	0.3	-	-	2.1	0.5	3.5	0.6	0.2	-	2.1	2.7
1928	-	-	-	-	0.0	-	-	-	-	-	-	-
1927	-	-	-	-	0.0	-	-	-	-	-	0.1	-
1926	-	-	-	-	0.1	0.4	-	-	-	-	0.2	0.1
1925	-	1.7	-	0.3	1.7	5.1	-	2.4	0.4	-	1.6	5.1
1924	-	-	-	-	-	-	-	-	-	-	-	-
1923	-	-	-	-	0.1	-	-	-	-	-	-	-
1922	0.0	-	-	-	0.1	0.1	-	-	-	0.4	-	-
1921	-	-	-	-	-	-	-	-	-	-	-	-
1920	0.0	0.7	-	4.7	29.1	48.2	33.5	12.8	-	5.2	5.0	17.1

See footnotes at end of table.

TABLE 22. - 1/CASE RATES FOR SELECTED COMMUNICABLE DISEASES BY PREFECTURE: JAPAN, 1920-1949 - Cont'd
CHOLERA - Cont'd

Year	Yamaguchi	Tokushima	Karawa	Ehime	Kochi	Fukuoka	Saga	Nagasaki	Yamamoto	Oita	Miyazaki	Kagoshima
1949	-	-	-	-	-	-	-	-	-	-	-	-
1948	-	-	-	-	-	-	-	-	-	-	-	-
1947	-	-	-	-	-	-	-	-	-	-	-	-
1946	6.4	-	0.1	1.2	-	6.2	10.0	11.2	1.5	0.5	1.7	3.9
1945	-	-	-	-	-	-	-	-	-	-	-	-
1944	-	-	-	-	-	-	-	-	-	-	-	-
1943	-	-	-	-	-	-	-	-	-	-	-	-
1942	-	-	-	-	-	-	-	-	-	-	-	-
1941	-	-	-	-	-	-	-	-	-	-	-	-
1940	-	-	-	-	-	-	-	-	-	-	-	-
1939	-	-	-	-	-	-	-	-	-	-	-	-
1938	-	-	-	-	-	0.1	-	0.1	-	-	-	-
1937	0.8	-	-	-	-	-	-	-	-	-	-	-
1936	-	-	-	-	-	-	-	-	-	-	-	-
1935	-	-	-	-	-	-	-	-	-	-	-	-
1934	-	-	-	-	-	-	-	-	-	-	-	-
1933	-	-	-	-	-	-	-	-	-	-	-	-
1932	0.1	-	-	-	-	-	-	-	-	-	-	-
1931	-	-	-	-	-	-	-	-	-	-	-	-
1930	-	-	-	-	-	-	-	-	-	-	-	-
1929	0.2	1.5	0.1	-	-	0.1	-	-	-	0.1	-	-
1928	-	-	-	-	-	-	-	-	-	-	-	-
1927	-	-	-	-	-	-	-	-	-	-	-	-
1926	-	-	1.0	-	-	-	-	-	-	-	-	-
1925	0.2	1.4	0.4	4.4	2.2	0.0	-	1.1	-	-	0.1	0.3
1924	-	-	-	-	-	-	-	-	-	-	-	-
1923	0.1	-	-	-	-	-	-	-	-	-	-	-
1922	0.5	-	-	-	-	7.5	3.3	1.9	0.1	-	-	-
1921	0.1	-	-	-	-	0.3	-	1.7	-	-	-	-
1920	15.8	10.2	5.5	17.7	13.4	22.8	18.1	31.9	22.1	23.0	10.3	2.7

See footnotes at end of table.

TABLE 32. - 1/CASE RATES FOR SELECTED COMMUNICABLE DISEASES BY PREFECTURE: JAPAN, 1920-1949 - Cont'd
EPIDEMIC MENINGITIS

Year	All Japan	Hokkaido	Aomori	Iwate	Miyagi	Akita	Yamagata	Fukushima	Ibaraki	Tochigi	Gumma
1949	1.8	3.7	2.9	2.0	3.8	2.9	2.5	2.6	1.6	0.7	1.3
1948	2.6	4.7	5.2	1.4	7.0	5.7	3.3	4.1	3.6	0.8	2.0
1947	4.3	9.6	8.8	4.8	8.7	7.1	6.2	7.3	9.8	2.0	2.4
1946	2.0	6.5	6.5	3.3	4.8	3.2	3.4	3.5	1.9	0.8	1.2
1945	6.3	45.8	12.8	4.4	17.4	1.3	1.4	7.2	10.2	1.2	4.3
1944	2.0	5.1	5.0	0.9	4.1	0.5	1.1	0.3	4.2	0.3	1.6
1943	1.5	2.1	7.0	0.8	3.9	2.7	0.7	1.2	1.5	0.2	1.8
1942	1.1	3.6	6.0	1.1	3.2	2.8	1.1	0.9	1.1	0.4	0.5
1941	1.6	5.4	12.6	4.0	5.1	5.2	1.1	0.9	1.7	0.8	0.5
1940	1.9	7.3	22.5	3.4	2.3	9.0	1.5	0.9	2.3	0.2	0.6
1939	2.3	5.5	34.2	1.6	1.6	1.0	1.3	1.7	2.3	0.1	0.5
1938	1.4	4.2	3.9	0.6	0.6	0.5	0.6	1.7	0.3	-	0.4
1937	1.2	0.8	3.5	0.3	0.5	0.5	0.7	0.4	1.4	0.2	0.4
1936	1.5	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1935	1.9	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1934	1.2	0.6	0.4	0.4	0.4	0.3	0.1	0.1	0.3	0.1	0.3
1933	0.5	1.3	0.8	0.3	1.0	0.2	0.9	0.4	0.3	0.1	1.5
1932	0.4	0.2	0.8	0.2	0.3	0.3	0.4	0.2	0.3	0.1	0.5
1931	0.4	0.6	0.6	0.1	0.3	1.3	-	0.5	0.4	-	0.5
1930	0.4	0.1	-	0.1	0.6	0.3	0.3	0.1	0.1	-	0.5
1929	0.6	0.1	0.1	0.1	0.6	0.1	0.1	0.7	0.2	-	1.4
1928	0.5	0.1	0.1	0.1	0.5	0.5	0.1	0.7	0.2	0.3	0.4
1927	0.5	0.0	0.2	-	1.0	0.2	0.1	0.5	0.4	0.3	0.3
1926	0.7	0.1	-	0.2	0.9	-	-	0.1	0.7	0.4	0.3
1925	0.8	0.2	0.4	0.2	1.3	0.1	0.6	0.2	0.2	0.4	0.4
1924	2.3	0.5	0.3	4.0	2.6	3.9	0.7	0.7	0.6	0.6	0.3
1923	1.2	0.0	0.1	0.2	3.3	0.9	0.1	0.3	4.2	0.6	0.4
1922	1.6	0.2	0.1	0.8	3.7	1.8	0.1	0.8	0.2	0.7	0.5
1921	1.4	2.2	0.4	0.7	4.6	0.7	0.6	0.5	0.3	0.1	-
1920	1.7	0.7	0.3	0.5	0.6	1.3	0.1	1.0	0.4	0.5	-

See footnotes at end of table.

TABLE 32. - 1/CASE RATES FOR SELECTED COMMUNICABLE DISEASES BY PREFECTURE: JAPAN, 1920-1949 - Cont'd
EPIDEMIC MENINGITIS - Cont'd

Year	Saitama	Chiba	Tokyo	Kanagawa	Miigata	Toyama	Ishikawa	Fukui	Yamanashi	Nagano	Gifu	Shizuoka
1949	1.4	1.0	4.6	3.2	1.0	1.7	0.4	2.0	1.3	1.1	0.6	1.3
1948	1.1	1.8	7.7	4.7	2.1	2.7	1.2	2.2	2.5	1.9	0.9	2.2
1947	3.4	2.9	13.3	3.5	2.8	2.1	4.9	1.7	3.4	2.0	1.2	4.0
1946	0.6	1.4	3.2	1.8	1.2	0.4	-	0.3	2.4	0.5	0.9	1.6
1945	0.7	0.8	1.3	6.7	1.1	2.4	0.4	1.4	-	1.0	0.8	1.4
1944	0.5	0.8	1.5	3.1	0.4	1.0	-	-	-	0.2	0.4	0.9
1943	0.3	0.6	1.4	2.6	0.0	0.1	0.1	-	-	0.2	0.2	0.3
1942	0.4	0.5	2.1	1.6	0.1	0.2	0.1	0.2	-	0.2	0.1	0.3
1941	0.4	0.5	1.7	1.7	0.2	0.2	0.1	0.2	0.5	0.2	0.4	0.4
1940	0.2	0.8	1.6	1.1	0.1	-	0.5	0.2	0.3	0.2	0.1	0.6
1939	0.1	0.6	1.5	0.6	0.2	-	-	0.3	0.6	0.4	0.2	1.0
1938	0.1	0.5	1.7	0.9	0.1	0.1	0.3	-	-	0.1	0.1	0.6
1937	0.5	0.3	1.8	0.6	0.2	0.3	0.3	0.2	0.3	0.4	0.2	0.3
1936	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1935	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1934	-	0.1	1.4	0.3	-	0.2	0.4	0.2	0.5	0.2	0.7	0.2
1933	0.1	0.2	1.0	0.5	0.2	-	0.4	0.3	0.2	-	0.2	-
1932	0.1	0.1	0.8	0.4	0.1	0.1	0.1	-	0.8	0.3	-	0.2
1931	0.1	0.1	1.4	0.8	0.2	-	-	-	0.3	0.1	0.3	0.1
1930	0.2	0.2	1.4	0.6	0.1	0.3	0.1	-	0.3	0.1	-	0.3
1929	0.1	0.1	1.5	0.6	-	0.1	-	-	0.2	-	-	0.3
1928	0.3	0.1	1.3	0.5	0.1	0.3	0.1	-	0.3	0.2	-	0.2
1927	0.1	-	1.5	0.3	0.1	0.1	0.1	-	0.3	0.1	0.1	0.3
1926	-	0.4	1.5	0.7	-	0.3	0.1	-	1.0	0.2	0.1	0.4
1925	0.1	0.8	1.8	0.4	0.1	-	0.1	0.2	2.3	0.1	0.3	2.7
1924	0.3	0.7	4.3	0.5	0.3	0.1	0.1	-	1.1	0.5	0.1	1.7
1923	0.2	0.4	3.3	0.8	0.1	-	0.3	-	1.8	0.1	0.4	4.0
1922	0.1	0.4	3.0	1.2	0.2	-	0.2	-	0.8	0.7	0.2	0.8
1921	-	0.1	2.6	1.6	0.2	-	0.3	-	1.5	0.1	0.1	0.5
1920	-	0.4	2.5	1.3	0.1	-	-	-	1.5	0.6	0.3	0.1

See footnotes at end of table.

TABLE 32. - 1/CASE RATES FOR SELECTED COMMUNICABLE DISEASES BY PREFECTURE: JAPAN, 1920-1949 - Cont'd
EPIDEMIC MENINGITIS - Cont'd

Year	Aichi	Mie	Shiga	Kyoto	Osaka	Hyogo	Nara	Wakayama	Tottori	Shimane	Okayama	Hiroshima
*1949	0.5	0.9	1.3	3.1	2.7	1.0	0.9	0.4	4.6	1.2	0.5	1.2
1948	0.7	0.2	0.9	3.6	2.6	1.3	1.2	0.7	4.1	1.8	0.5	1.1
1947	1.4	2.0	3.4	4.1	4.5	2.3	0.8	1.0	7.7	0.9	0.7	3.3
1946	0.5	1.6	1.1	2.6	1.5	0.8	0.4	0.2	2.6	1.9	0.4	0.9
1945	0.7	1.9	2.2	6.5	0.9	2.2	1.2	1.8	4.1	1.3	0.9	5.2
1944	0.4	0.1	0.7	0.9	1.3	1.0	0.3	0.5	2.1	0.1	0.2	2.5
1943	0.4	0.3	0.9	2.3	1.9	1.4	0.3	0.6	0.8	0.1	0.3	0.6
1942	0.3	0.3	0.6	1.9	1.4	0.8	0.3	0.4	1.0	-	-	0.5
1941	0.5	0.3	0.6	2.7	2.0	1.0	1.1	0.2	2.1	0.3	0.3	0.1
1940	0.4	0.3	1.2	2.1	2.5	0.8	3.0	0.5	1.3	-	0.5	0.3
1939	0.4	0.4	1.5	3.7	3.3	1.3	6.0	1.4	6.2	0.3	1.1	0.6
1938	0.2	0.3	1.0	3.1	2.6	1.4	0.8	1.3	0.6	0.1	0.5	0.6
1937	0.2	0.2	0.9	1.7	2.2	0.8	0.3	1.5	1.5	0.3	0.1	0.3
1936	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1935	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1934	0.3	0.2	1.3	2.1	3.4	0.7	-	1.0	-	-	0.2	1.1
1933	0.2	-	1.0	1.6	1.4	0.2	-	0.7	1.4	0.1	0.1	0.2
1932	0.1	0.1	0.4	0.7	1.7	0.3	-	1.0	0.2	0.1	0.2	0.4
1931	0.2	-	0.4	0.6	1.5	0.3	0.2	1.0	-	0.3	0.2	0.4
1930	0.3	0.2	0.4	1.2	1.7	0.3	0.2	0.5	-	0.4	-	0.2
1929	0.4	0.1	0.3	0.9	1.9	0.7	-	0.5	0.6	0.5	0.6	0.3
1928	0.2	0.2	0.3	1.0	1.9	0.4	0.3	0.5	-	0.7	0.2	0.5
1927	0.2	0.4	0.3	2.5	2.5	0.7	0.3	0.4	1.2	0.4	0.2	0.4
1926	0.1	0.3	0.3	1.3	3.4	0.9	0.3	0.4	1.0	0.4	0.5	0.7
1925	0.4	0.3	0.2	1.8	2.8	0.7	0.3	0.3	1.9	-	0.3	0.2
1924	0.8	0.4	-	2.4	3.1	1.4	1.0	1.2	9.5	3.4	30.7	8.9
1923	0.6	0.6	-	2.1	3.5	1.3	0.4	0.4	1.5	4.1	0.6	0.7
1922	0.5	0.5	-	4.2	3.8	2.3	1.4	0.4	4.6	3.8	7.7	4.9
1921	0.3	-	-	4.7	6.1	2.1	0.9	0.5	3.5	4.8	0.4	1.2
1920	0.4	0.6	-	4.2	10.9	1.9	0.5	1.1	3.5	1.3	0.6	1.2

See footnotes at end of table.

TABLE 32. - 1/CASE RATES FOR SELECTED COMMUNICABLE DISEASES BY PREFECTURE: JAPAN, 1920-1949 - Cont'd
EPIDEMIC MENINGITIS - Cont'd

Year	Yamaguchi	Tokushima	Kagawa	Ehime	Kochi	Fukuoka	Saga	Nagasaki	Kumamoto	Oita	Miyazaki	Kagoshima
1949	0.8	0.1	0.9	0.9	1.2	1.7	1.0	0.5	0.4	1.1	0.7	0.4
1948	1.3	0.6	1.0	1.4	0.6	1.3	0.8	1.7	1.1	0.8	0.7	1.4
1947	2.4	1.1	2.2	2.6	3.0	2.7	1.9	2.2	1.9	1.2	2.6	2.2
1946	2.3	2.0	1.0	0.4	0.9	3.2	1.2	2.7	0.8	0.4	2.5	1.1
1945	13.3	0.1	2.9	2.4	4.7	24.8	2.1	24.6	2.5	0.8	2.6	0.7
1944	1.3	0.3	0.1	0.5	0.6	4.2	0.7	31.5	1.5	-	0.6	0.9
1947	0.6	0.1	0.5	0.6	0.6	3.1	0.7	18.2	0.7	-	0.6	0.6
1942	0.8	-	1.0	0.6	1.4	2.3	0.4	1.0	0.4	-	0.1	0.1
1941	0.5	0.4	1.7	0.3	1.4	4.1	1.5	0.8	0.1	-	0.5	0.6
1940	0.5	-	0.3	0.8	0.9	5.6	0.6	1.3	0.2	0.4	0.7	0.3
1939	1.0	0.7	1.3	0.3	3.2	10.1	1.6	1.6	0.8	0.9	0.5	1.0
1938	0.5	1.2	1.1	0.5	1.5	8.2	0.1	1.2	0.6	0.1	0.1	0.3
1937	1.1	0.1	0.1	0.1	1.8	9.3	0.3	0.9	0.7	0.9	0.2	0.2
1936	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1935	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1934	0.3	0.1	0.7	0.1	0.4	27.5	-	0.8	0.6	-	0.4	0.4
1937	0.2	-	0.5	-	0.9	0.3	-	0.9	0.1	-	-	0.3
1932	0.2	-	-	-	0.1	0.2	-	0.4	0.2	-	0.1	0.1
1931	0.1	0.1	0.1	-	0.4	0.5	-	0.2	-	0.1	-	0.1
1930	0.1	0.3	0.5	-	0.8	0.2	-	0.4	0.7	-	0.3	-
1929	1.1	0.1	1.5	0.1	2.5	0.7	-	0.7	0.3	-	-	0.1
1928	0.2	0.3	0.3	0.2	1.1	0.7	-	0.3	0.2	-	0.4	-
1927	0.2	0.7	0.1	0.1	0.4	0.7	-	0.2	0.1	-	0.4	-
1926	1.4	0.6	-	-	1.0	0.9	-	0.3	0.3	-	0.3	1.1
1925	0.4	1.2	1.0	0.2	0.6	1.0	-	0.9	0.5	-	0.3	0.1
1924	2.4	0.7	0.6	0.7	4.5	2.3	0.1	2.2	0.6	0.1	-	0.6
1923	0.6	0.7	0.6	0.3	1.8	1.6	0.6	1.2	1.0	0.3	-	0.1
1922	0.7	6.8	3.7	1.3	2.6	2.6	0.7	1.2	0.6	-	-	0.5
1921	1.1	1.1	2.2	1.2	2.1	2.9	1.5	1.2	1.4	-	0.2	0.5
1920	1.2	1.3	1.0	1.4	2.5	8.0	-	3.3	1.0	-	-	0.1

See footnotes at end of table.

TABLE 32. - 1/CASE RATES FOR SELECTED COMMUNICABLE DISEASES BY PREFECTURE: JAPAN, 1920-1949 - Cont'd
SCARLET FEVER

Year	All Japan	Fukushima	Aomori	Iwate	Miyagi	Akita	Yamagata	Fukushima	Ibaraki	Tochigi	Gumma
*1949	5.7	10.0	2.9	4.4	4.2	3.6	4.6	3.2	3.0	3.6	4.9
1948	3.7	12.1	2.0	1.5	5.9	1.9	2.0	1.9	3.0	3.3	5.7
1947	3.4	9.5	2.4	2.2	6.7	2.5	2.4	2.4	3.2	2.7	3.2
1946	2.9	11.0	2.0	1.1	7.7	4.1	4.1	1.8	2.9	1.4	2.7
1945	3.4	11.3	2.7	2.4	2.2	3.8	3.2	1.8	1.9	1.5	4.7
1944	8.8	21.9	8.2	4.9	7.4	4.0	7.8	5.2	3.4	4.1	10.2
1943	13.6	25.4	8.8	15.5	17.6	7.3	7.0	5.4	5.0	3.1	10.3
1942	17.6	35.6	15.0	18.2	17.1	8.2	5.8	8.6	9.3	5.7	12.0
1941	20.9	37.0	13.9	13.7	23.7	19.9	10.6	11.2	9.4	6.8	14.8
1940	27.3	33.5	22.1	13.6	31.3	15.7	9.7	19.8	10.3	8.2	24.5
1939	28.3	33.4	27.5	16.8	42.3	17.0	11.8	22.9	11.0	15.7	26.2
1938	27.1	36.9	13.1	17.1	31.2	14.2	11.0	10.7	10.1	11.3	20.5
1937	25.2	34.1	15.3	28.0	26.5	19.2	6.1	3.9	8.1	7.6	20.0
1936	24.2	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1935	24.2	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1934	24.6	35.0	12.3	19.0	13.0	9.2	2.7	2.1	4.7	5.7	13.4
1933	18.9	26.3	15.5	17.7	6.9	10.6	3.5	2.8	5.1	3.3	8.7
1932	12.5	17.2	6.9	9.3	3.0	7.6	1.5	1.8	2.8	2.0	7.6
1931	10.0	12.2	3.7	9.5	5.0	7.7	2.4	1.2	3.2	3.0	6.3
1930	9.4	15.3	5.5	6.7	7.3	11.3	3.1	1.1	2.2	2.5	7.1
1929	9.0	20.5	5.1	1.6	7.4	3.5	2.3	1.5	2.7	1.8	6.8
1928	9.2	16.0	4.3	3.7	6.7	2.8	1.3	1.9	1.0	1.5	5.0
1927	5.8	8.6	2.7	1.1	6.1	2.5	0.3	1.6	1.3	1.9	3.3
1926	5.5	7.7	2.7	1.2	2.3	1.6	0.5	1.0	0.8	0.5	3.0
1925	4.3	4.2	1.2	0.6	1.1	0.9	0.4	0.8	0.4	1.6	2.6
1924	3.2	2.3	1.3	0.7	1.7	1.9	0.2	0.5	0.6	2.6	1.6
1923	2.7	2.6	1.0	0.5	1.0	2.4	0.6	0.3	0.5	0.4	1.0
1922	2.9	2.4	1.7	0.9	1.5	0.4	0.3	-	-	0.7	3.1
1921	2.3	2.3	2.2	1.6	0.9	1.7	-	0.3	0.5	0.9	3.0
1920	2.5	3.2	0.5	3.1	2.4	1.9	0.5	0.5	0.1	0.8	1.2

See footnotes at end of table.

TABLE 32. - 1/CASE RATES FOR SELECTED COMMUNICABLE DISEASES BY PREFECTURE: JAPAN, 1920-1949 - Cont'd
SCARLET FEVER - Cont'd

Year	Saitama	Chiba	Tokyo	Kanagawa	Niigata	Toyama	Ishikawa	Fukui	Yamanashi	Nagano	Gifu	Shizuoka
1949	9.2	6.7	19.5	10.5	2.1	1.1	0.8	1.3	5.6	17.9	3.5	2.7
1948	6.0	1.0	11.7	4.4	1.4	0.9	0.4	0.8	4.1	6.4	3.7	2.4
1947	2.8	2.4	10.4	5.3	1.3	1.5	0.6	0.8	2.9	4.8	1.7	5.7
1946	3.2	2.0	9.5	4.6	1.1	0.5	0.6	0.7	1.3	3.6	1.5	2.7
1945	7.8	3.7	4.7	4.8	0.7	0.6	0.3	1.4	2.7	8.6	2.8	4.7
1944	12.1	7.6	24.4	13.0	1.5	1.3	-	2.6	5.9	11.8	4.0	7.9
1943	17.0	11.3	52.2	26.8	1.6	0.5	0.5	1.9	8.8	9.8	3.6	8.4
1942	21.0	10.6	68.3	32.0	1.8	0.5	0.5	3.7	4.7	8.4	6.0	9.0
1941	10.2	13.6	79.5	51.7	2.6	2.5	0.7	3.4	4.4	11.8	9.2	9.5
1940	36.0	19.1	115.1	57.1	3.1	2.0	1.5	2.2	10.7	14.5	10.5	12.0
1939	32.8	24.3	113.0	54.7	3.4	0.7	0.8	3.5	14.6	21.4	8.3	8.8
1938	22.5	20.7	108.6	45.5	4.9	0.0	0.3	2.4	22.5	19.3	11.6	9.7
1937	24.7	14.4	92.7	33.1	5.6	1.0	1.1	1.4	6.3	24.8	7.3	9.1
1936	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1935	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1934	14.4	6.9	95.7	46.3	1.4	2.5	1.4	2.0	7.0	10.6	3.1	11.5
1933	15.3	6.3	83.5	24.7	1.1	1.9	1.8	1.7	6.1	8.1	7.0	9.8
1932	8.9	3.1	66.3	19.9	0.7	1.0	1.6	1.6	3.1	6.4	5.8	5.7
1931	11.8	2.9	55.9	18.8	0.7	1.0	1.1	1.4	1.6	5.7	8.9	5.2
1930	11.5	1.4	49.3	14.3	0.8	0.3	0.8	0.8	4.1	5.2	5.4	7.0
1929	5.2	2.1	48.1	15.9	0.4	0.6	0.7	0.7	3.9	2.8	1.7	7.0
1928	7.7	2.3	51.6	12.4	0.9	0.9	0.8	1.0	5.0	1.5	0.8	5.6
1927	3.2	1.3	36.7	10.3	0.5	0.3	0.5	0.2	14.9	1.7	0.9	3.8
1926	1.5	1.4	23.1	6.2	0.2	0.1	0.5	-	0.5	1.3	0.8	1.8
1925	1.4	1.1	16.5	4.3	0.2	0.3	0.1	-	1.2	0.9	1.4	2.1
1924	1.1	1.6	18.5	4.4	0.3	0.3	-	0.7	0.7	0.9	0.9	1.0
1923	1.5	0.6	12.1	5.8	0.4	-	-	0.2	1.2	1.1	0.3	1.0
1922	1.3	0.7	12.8	5.1	0.1	0.3	0.3	0.5	5.1	1.1	0.6	1.3
1921	1.9	0.5	15.1	5.7	-	0	0.4	1.0	1.4	0.8	0.6	1.4
1920	1.7	0.7	12.4	3.9	0.6	-	0.5	0.2	0.5	0.3	0.4	0.7

See footnotes at end of table.

TABLE 32. - 1/CASE RATES FOR SELECTED COMMUNICABLE DISEASES BY PREFECTURE: JAPAN, 1920-1949 - Cont'd
SCARLET FEVER - Cont'd

Year	Aichi	Mie	Shiga	Kyoto	Osaka	Hyogo	Nara	Wakayama	Tottori	Shimane	Okayama	Hiroshima
1949	5.5	3.7	12.3	14.3	5.2	2.7	3.9	2.6	3.0	6.5	6.6	3.8
1948	4.1	3.2	7.4	6.7	2.8	1.8	0.9	0.6	1.5	2.1	2.3	1.1
1947	3.9	3.3	5.8	8.0	1.7	1.3	1.3	0.7	1.0	1.1	1.7	1.1
1946	2.8	1.1	5.0	7.6	1.6	1.7	1.4	2.5	1.4	1.0	1.6	0.7
1945	2.7	3.7	6.9	10.6	1.4	3.0	1.2	1.3	1.7	1.6	3.0	3.3
1944	8.8	5.4	11.6	19.1	6.0	8.0	1.1	3.1	1.7	2.2	5.6	4.4
1943	14.2	5.1	8.8	23.6	12.8	13.4	3.2	3.1	2.9	0.9	6.8	2.6
1942	17.0	7.9	14.4	39.7	16.7	16.9	4.2	3.4	3.5	1.9	5.7	3.3
1941	14.9	13.9	20.4	45.8	21.9	20.3	5.7	3.7	4.8	1.7	4.9	4.1
1940	13.8	13.5	14.7	54.3	33.6	24.9	7.1	5.4	3.2	4.2	8.0	4.5
1939	13.6	12.9	19.0	54.0	35.1	27.0	8.8	8.0	5.1	9.0	8.1	5.6
1938	15.3	8.9	21.2	57.4	37.4	27.4	6.9	10.0	8.6	5.3	9.4	8.4
1937	16.2	9.2	25.1	65.3	36.1	27.9	5.2	11.0	7.3	4.9	14.4	14.9
1936	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1935	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1934	19.9	7.5	16.3	100.0	54.7	27.5	9.9	7.0	2.4	1.7	18.0	10.0
1933	27.9	10.3	14.1	73.0	33.5	15.0	5.5	4.2	2.2	1.5	10.7	5.6
1932	14.4	6.2	11.5	37.1	16.4	7.9	2.2	4.4	1.2	0.8	5.9	1.9
1931	5.6	3.4	8.3	28.5	8.7	5.7	2.3	2.1	-	0.9	3.9	1.9
1930	5.5	5.5	6.4	27.5	9.3	6.1	1.5	3.9	-	0.7	9.6	0.8
1929	3.9	6.3	6.3	30.4	11.1	7.9	2.3	4.8	0.2	1.2	5.4	1.7
1928	3.8	5.5	7.1	47.8	12.7	8.0	3.3	2.7	-	1.6	4.2	2.1
1927	3.9	6.5	3.6	32.3	8.7	5.4	3.5	3.0	0.4	1.5	2.0	2.0
1926	2.7	4.5	4.8	24.4	7.6	4.6	1.5	2.0	1.0	0.8	2.9	3.2
1925	1.6	2.8	4.2	21.9	4.5	3.9	1.4	1.3	0.6	1.2	1.3	1.3
1924	2.7	2.6	2.3	19.0	3.8	3.9	1.9	1.0	0.4	0.9	3.0	1.8
1923	1.4	0.7	3.4	27.8	3.9	3.2	1.4	2.0	0.2	0.7	3.9	0.4
1922	1.1	1.7	5.7	28.8	4.4	3.1	0.9	1.4	0.2	0.3	7.6	0.3
1921	1.1	1.8	5.1	29.7	3.9	1.4	0.4	1.2	0.2	1.0	0.3	0.3
1920	0.7	0.7	3.2	25.2	4.5	1.2	0.2	0.5	-	-	1.0	0.4

See footnotes at end of table.

TABLE 32. - 1/CASE RATES FOR SELECTED COMMUNICABLE DISEASES BY PREFECTURE: JAPAN 1920-1949 - Cont'd
SCARLET FEVER - Cont'd

Year	Yamanashi	Tokushima	Karawa	Ehime	Kochi	Fukuoka	Saga	Nagasaki	Kumamoto	Oita	Miyazaki	Kagoshima
• 1949	7.2	1.2	2.3	1.6	1.2	1.1	0.5	1.1	0.2	0.6	0.6	0.2
1948	1.3	0.5	1.9	1.8	1.4	1.7	0.9	1.1	0.1	0.4	0.7	0.5
1947	1.0	0.4	1.5	1.7	1.1	0.7	0.2	1.8	0.3	0.2	1.1	0.2
1946	1.8	0.2	0.3	1.5	0.3	1.3	0.2	1.2	0.2	0.3	1.0	0.3
1945	3.0	0.1	2.9	0.8	1.5	1.8	0.3	2.4	0.5	0.4	0.1	0.1
1944	7.4	3.1	4.1	7.1	3.8	7.5	1.3	4.3	1.3	0.2	1.2	0.8
1943	7.7	5.8	9.4	7.7	4.9	4.9	1.1	7.9	0.6	0.6	2.2	0.9
1942	4.8	2.4	11.1	8.3	3.2	5.1	0.7	1.7	1.8	0.2	1.2	1.1
1941	7.2	7.2	7.1	7.0	7.6	7.6	1.2	6.8	7.1	0.7	0.8	0.6
1940	9.5	3.3	5.5	5.4	6.6	11.1	3.2	9.0	1.2	1.2	1.5	1.0
1939	11.0	5.0	6.5	6.4	9.4	13.7	1.8	10.8	2.7	1.2	1.5	1.0
1938	10.4	4.0	10.2	9.0	13.0	13.5	3.1	15.1	5.0	1.2	1.1	1.6
1937	8.7	7.0	15.1	6.5	17.8	8.9	1.2	19.1	2.2	1.0	2.4	9.4
1936	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1935	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1934	20.3	7.4	11.7	6.5	5.4	16.6	0.3	9.1	2.9	0.4	1.7	0.7
1933	8.2	3.5	5.0	2.8	5.0	9.2	0.1	5.4	1.7	0.1	0.7	0.5
1932	6.9	0.8	4.6	2.8	6.7	4.4	1.0	6.6	2.8	3.2	1.8	1.8
1931	7.2	0.8	2.6	2.4	0.6	3.6	0.4	6.2	1.5	0.3	1.3	0.8
1930	3.6	0.3	3.5	2.6	0.1	2.9	0.4	4.9	4.9	-	1.4	1.4
1929	5.0	0.6	6.1	2.3	1.1	1.7	0.1	3.0	0.1	0.6	0.4	1.0
1928	7.2	-	5.7	2.2	2.1	2.8	-	1.8	1.0	0.5	1.4	0.3
1927	2.7	0.4	3.1	2.0	1.4	1.9	-	2.1	0.2	0.4	17.4	0.1
1926	4.5	0.3	2.8	1.5	0.7	13.8	7.8	13.0	0.6	1.0	12.8	0.9
1925	7.3	0.6	1.4	1.3	0.7	23.3	0.1	1.6	0.4	0.4	-	1.3
1924	1.6	0.1	0.6	1.1	0.4	1.3	-	0.4	1.6	0.9	0.3	0.1
1923	1.6	-	1.5	0.9	1.6	0.5	-	0.3	0.9	3.7	0.1	0.1
1922	0.6	0.1	0.9	0.5	1.0	0.3	-	0.8	0.1	0.2	0.1	0.1
1921	1.1	-	1.5	1.0	0.7	0.3	-	0.2	0.4	0.5	0.2	0.1
1920	1.4	-	0.5	1.5	0.0	0.3	-	0.3	0.2	0.2	-	0.2

See footnotes at end of table.

TABLE 32. - 1/CASE RATES FOR SELECTED COMMUNICABLE DISEASES BY PREFECTURE: JAPAN, 1920-1949 - Cont'd
2/DIPHTHERIA

Year	All Japan	Hokkaido	Aomori	Iwate	Miyagi	Akita	Yamagata	Fukushima	Ibaraki	Tochigi	Gumma
*1949	18.0	21.1	21.1	25.2	26.6	29.1	15.2	13.4	12.6	14.1	12.9
1948	20.3	32.2	19.1	22.5	28.9	44.4	21.6	10.9	15.4	21.2	15.9
1947	36.4	61.7	42.6	33.5	38.1	55.7	50.2	21.0	26.8	46.5	21.4
1946	65.5	115.5	68.8	66.8	62.5	82.1	94.5	65.7	30.0	59.5	16.4
1945	122.8	182.8	142.2	112.4	137.8	107.4	148.1	153.7	62.1	184.5	54.8
1944	129.9	172.4	153.0	122.8	150.6	99.7	136.2	228.9	85.7	181.8	43.4
1943	87.7	114.9	101.5	132.0	104.8	84.7	77.8	158.1	37.7	81.7	43.4
1942	61.5	76.5	58.1	78.0	69.9	77.7	50.6	119.1	35.4	47.8	46.5
1941	56.4	71.3	68.1	99.1	54.7	110.3	49.2	122.2	34.2	31.0	33.4
1940	54.1	72.9	90.7	101.7	65.1	100.2	59.0	121.6	33.2	23.4	37.0
1939	50.9	70.6	89.2	92.7	65.5	88.4	47.8	96.8	30.9	24.8	49.9
1938	40.4	73.8	73.8	65.4	48.7	88.5	36.5	71.2	19.5	27.7	32.9
1937	40.1	84.1	89.3	55.0	52.6	97.7	44.4	63.2	26.4	38.3	45.3
1936	40.9	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1935	41.4	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1934	44.3	108.8	76.0	53.4	59.5	102.1	38.6	83.8	35.3	47.2	45.4
1933	42.7	108.7	56.0	46.7	60.6	110.4	40.0	74.6	21.0	50.6	44.3
1932	33.1	72.5	40.4	42.2	48.2	75.1	44.8	67.8	23.5	37.2	39.9
1931	32.5	80.2	43.6	46.6	42.1	73.9	29.3	68.6	27.6	34.6	43.5
1930	29.0	78.9	62.1	41.9	60.7	77.1	40.0	59.5	40.8	37.7	32.0
1929	31.3	73.3	45.9	26.3	50.0	47.9	34.5	49.7	31.4	39.1	30.7
1928	28.2	79.1	37.6	24.8	42.7	40.1	27.4	49.5	24.5	31.2	43.8
1927	24.8	70.2	29.2	25.5	40.8	38.7	27.4	54.2	18.5	23.9	21.8
1926	22.6	54.6	28.9	27.8	48.8	41.0	23.0	52.2	15.7	21.0	26.6
1925	23.3	67.7	37.9	25.0	45.8	35.8	30.0	56.0	19.4	23.8	32.8
1924	22.4	56.4	31.7	30.8	52.7	40.3	31.7	46.9	16.2	29.4	30.4
1923	22.1	63.7	41.7	24.4	46.6	49.5	39.2	48.1	19.5	30.0	36.9
1922	24.1	51.5	29.6	29.6	50.9	68.6	49.9	64.4	28.0	33.4	43.2
1921	25.8	76.1	62.4	35.9	71.0	71.0	37.7	40.0	24.2	36.5	40.9
1920	27.3	91.6	58.2	44.6	58.2	58.4	31.7	48.7	29.4	42.0	42.6

See footnotes at end of table.

TABLE 32. 1/CASE RATES FOR SELECTED COMMUNICABLE DISEASES BY PREFECTURE: JAPAN, 1920-1949 - Cont'd
2/DIPHTHERIA - Cont'd

Year	Saitama	Chiba	Tokyo	Kanagawa	Niigata	Toyama	Ishikawa	Fukui	Yamanashi	Nagano	Gifu	Shizuoka
1949	15.5	10.2	18.9	16.6	23.8	24.5	25.9	22.5	8.0	14.1	10.7	11.6
1948	17.4	8.5	19.2	19.4	27.5	17.4	30.0	17.8	8.0	19.2	8.6	11.5
1947	31.0	20.0	32.2	25.5	35.0	23.3	67.8	31.2	13.6	30.5	13.7	23.7
1946	45.8	41.8	52.3	50.8	51.0	51.0	106.3	51.1	26.8	39.5	31.1	56.0
1945	181.7	108.8	62.0	134.1	110.6	131.3	237.7	109.1	81.9	134.2	88.5	149.1
1944	204.6	151.8	202.1	183.8	91.6	118.4	177.5	83.9	113.8	98.2	56.4	117.3
1943	180.2	91.3	182.4	131.2	67.8	44.4	40.8	52.0	62.5	78.7	37.1	59.5
1942	108.7	56.5	140.2	140.3	50.0	6.8	24.6	31.7	45.4	57.6	22.6	40.0
1941	69.4	40.7	105.7	117.6	39.3	6.4	21.6	29.9	40.1	60.1	27.8	35.1
1940	52.6	48.4	102.0	106.9	28.9	4.7	18.5	24.5	45.8	50.4	24.0	31.5
1939	36.4	35.3	93.2	99.9	28.6	3.2	21.5	20.2	37.1	55.9	24.0	31.8
1938	29.1	30.1	61.4	63.2	33.3	4.5	17.1	13.1	25.0	50.3	18.8	20.8
1937	42.8	29.3	56.7	68.1	32.7	4.3	14.4	9.8	28.1	55.3	17.6	18.8
1936	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1935	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1934	47.7	25.3	86.8	77.3	23.7	4.4	6.9	13.9	22.4	49.6	19.6	25.1
1933	41.9	28.2	101.6	80.5	27.0	5.9	7.6	14.3	20.1	41.5	16.9	26.4
1932	30.5	18.8	71.4	61.4	21.1	5.6	7.1	19.1	19.7	33.7	16.2	28.4
1931	34.4	21.9	69.3	63.9	23.0	5.6	4.7	19.3	16.2	32.9	10.9	24.0
1930	35.1	20.3	47.3	39.5	26.8	4.9	9.2	17.5	16.8	30.2	12.1	18.2
1929	30.9	20.3	66.2	53.7	24.2	5.5	9.7	12.1	18.9	35.7	10.2	20.3
1928	23.5	15.5	59.7	50.1	23.7	6.6	8.4	7.8	22.7	29.8	12.3	16.7
1927	19.1	17.0	46.9	35.8	20.9	2.9	7.8	11.0	20.3	25.1	11.2	13.6
1926	16.6	14.8	31.2	24.6	23.4	4.5	6.8	9.8	24.9	33.0	14.4	14.4
1925	18.9	19.0	35.7	29.4	24.5	6.1	6.5	9.4	27.3	31.4	13.0	17.8
1924	20.3	21.4	38.0	33.5	26.9	6.2	8.2	10.8	28.4	29.4	10.7	22.7
1923	25.2	23.0	24.9	21.3	24.4	4.9	12.2	19.9	22.9	36.7	14.6	22.1
1922	32.2	24.7	27.4	24.9	26.2	4.5	16.4	17.2	30.3	41.1	13.6	20.4
1921	31.2	25.3	37.3	30.1	30.8	6.3	18.4	12.4	22.8	34.0	15.9	26.2
1920	35.5	24.8	29.4	36.1	32.3	5.4	21.4	15.9	19.4	33.2	14.5	23.6

See footnotes at end of table.

TABLE 32. - 1/CASE RATES FOR SELECTED COMMUNICABLE DISEASES BY PREFECTURE: JAPAN, 1920-1949 - Cont'd
2/DIPHTHERIA - Cont'd

Year	Aichi	Mie	Shiga	Kyoto	Osaka	Hyogo	Nara	Wakayama	Tottori	Shimane	Okayama	Hiroshima
1949	11.3	13.7	11.9	11.0	10.0	14.2	12.4	10.1	14.2	29.8	11.2	20.0
1948	14.3	18.3	12.3	14.6	8.7	15.5	18.2	12.2	15.4	33.9	16.2	21.0
1947	31.2	48.1	29.2	29.2	12.9	28.8	22.3	23.5	30.1	57.8	23.9	32.9
1946	66.2	82.2	79.3	57.7	32.7	56.1	55.8	60.2	65.5	100.6	59.7	63.5
1945	175.4	161.3	172.0	190.2	73.3	129.7	153.7	76.2	109.1	139.7	156.3	152.1
1944	228.8	133.5	133.1	144.2	82.5	132.8	37.3	39.4	93.4	80.7	125.5	113.0
1943	128.8	82.7	77.4	131.7	73.3	87.7	28.5	39.4	64.3	40.0	54.5	102.3
1942	64.8	47.7	56.9	100.9	57.1	64.0	31.0	28.4	42.3	29.4	32.9	37.4
1941	36.5	37.6	44.7	121.6	63.4	62.4	27.4	23.3	54.1	50.3	25.9	39.5
1940	34.6	32.5	36.3	97.3	65.8	64.2	29.3	30.4	42.4	54.7	21.0	31.1
1939	35.0	33.2	37.2	100.9	81.4	48.3	26.1	33.1	44.9	27.3	21.9	22.4
1938	29.8	31.7	31.6	72.9	68.9	34.6	23.1	34.9	32.2	23.2	19.1	15.7
1937	23.1	32.9	22.0	52.9	57.3	38.1	25.8	22.4	31.8	32.0	22.6	13.7
1936	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1935	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1934	17.5	35.8	35.9	70.6	70.3	32.3	15.2	41.8	29.0	22.5	19.9	14.9
1933	13.4	35.5	23.9	45.5	73.4	27.5	14.4	32.9	26.0	12.5	10.9	10.2
1932	13.3	26.8	20.2	41.0	49.0	21.7	14.3	24.7	14.3	18.5	11.0	12.0
1931	15.9	30.8	24.1	46.8	44.6	20.1	20.0	20.3	13.4	13.2	13.0	7.7
1930	14.6	26.2	12.1	41.2	32.9	19.5	14.6	16.6	14.7	11.6	12.7	11.8
1929	14.8	36.1	12.4	41.4	53.4	23.5	17.9	19.5	14.9	19.2	21.0	11.7
1928	12.0	29.3	12.7	53.4	44.9	25.5	12.1	16.6	9.0	13.6	11.3	11.3
1927	12.6	30.0	18.0	30.7	40.5	18.5	16.5	23.9	8.1	10.7	11.3	11.2
1926	12.4	25.9	20.2	22.8	34.1	19.4	9.5	21.2	23.0	12.2	12.8	17.1
1925	13.8	24.8	23.9	22.7	32.9	15.3	10.8	20.6	14.0	8.7	7.1	14.5
1924	17.1	26.7	14.2	26.0	22.8	12.4	8.0	15.7	11.0	11.4	5.6	9.1
1923	16.0	25.7	20.5	21.3	22.9	11.0	11.4	17.4	15.7	13.1	6.2	7.9
1922	17.8	31.4	15.4	27.1	19.7	13.3	9.2	16.8	21.4	18.3	7.1	8.5
1921	15.1	31.6	24.4	36.0	23.7	7.8	15.0	17.5	20.8	19.8	10.2	11.8
1920	17.6	32.6	28.0	30.1	31.9	9.3	13.5	29.6	13.2	17.3	9.0	5.0

See footnotes at end of table.

TABLE 32. - 1/CASE RATES FOR SELECTED COMMUNICABLE DISEASES BY PREFECTURE: JAPAN, 1920-1949 - Cont'd
2/DIPHTHERIA - Cont'd

Year	Yamaguchi	Tokushima	Kagawa	Ehime	Kochi	Fukuoka	Saga	Nagasaki	Kumamoto	Oita	Miyazaki	Kagoshima
* 1949	24.1	13.9	12.5	14.3	15.3	27.8	44.8	27.5	13.8	30.0	48.8	24.3
1948	21.8	14.7	12.6	23.8	17.0	27.0	59.7	29.6	10.1	44.4	40.0	26.4
1947	48.3	34.2	29.8	61.1	38.6	56.1	91.9	44.1	13.1	63.1	56.8	37.7
1946	142.8	72.4	89.4	88.6	151.6	109.0	131.7	68.4	22.0	95.4	82.4	47.1
1945	198.2	134.0	206.4	116.9	267.5	128.5	98.9	68.7	28.8	136.9	42.2	33.8
1944	179.1	73.4	93.0	90.9	161.7	60.5	76.6	104.8	15.6	149.0	42.1	53.9
1943	85.9	37.5	54.3	60.0	35.8	32.7	37.7	44.7	7.6	17.1	18.3	13.2
1942	50.9	17.8	23.8	44.3	18.8	22.5	23.0	29.7	6.6	3.4	17.8	4.4
1941	71.1	14.2	23.7	34.9	24.7	29.5	37.8	54.8	8.4	4.9	23.3	5.0
1940	67.1	12.3	11.5	28.7	25.0	27.4	42.8	51.1	9.7	5.7	19.4	6.5
1939	60.6	16.9	19.4	27.3	23.1	26.7	30.8	39.3	7.7	4.2	17.0	4.8
1938	46.1	17.3	16.9	29.3	20.1	25.2	40.0	31.7	5.5	4.9	16.6	5.6
1937	50.7	19.4	15.0	23.3	32.9	29.5	23.4	22.7	4.1	7.1	15.2	6.8
1936	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1935	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1934	58.6	13.9	20.8	28.0	16.5	21.0	5.3	23.2	3.4	5.9	15.3	7.3
1933	50.7	10.9	18.2	20.7	28.2	17.8	3.0	20.8	3.2	4.2	12.8	6.1
1932	47.8	5.9	11.5	18.1	26.7	14.6	2.7	19.9	4.0	2.8	16.2	4.8
1931	30.2	6.2	13.4	17.6	16.4	16.8	1.4	15.1	3.3	2.8	15.4	4.8
1930	16.2	7.3	14.9	17.6	11.1	11.6	1.6	15.6	3.2	2.1	28.4	4.9
1929	27.0	6.7	17.6	18.7	11.0	12.4	1.9	18.8	3.2	3.0	31.5	8.2
1928	27.6	11.2	13.0	17.4	8.9	16.4	2.4	19.0	4.8	3.5	16.5	5.7
1927	29.7	9.1	11.8	19.7	10.2	15.3	1.9	21.1	4.7	4.0	10.5	7.2
1926	22.4	8.5	12.4	17.6	10.1	14.1	2.6	17.3	4.5	5.3	8.3	5.2
1925	21.4	8.8	10.0	19.7	7.4	12.3	4.5	15.9	3.2	4.6	6.7	4.1
1924	20.9	6.9	9.3	20.3	5.6	10.4	1.8	14.9	3.8	4.7	9.0	4.8
1923	15.2	6.1	13.0	15.3	6.5	9.0	2.5	12.7	5.4	2.7	8.2	3.5
1922	15.4	7.4	11.9	14.4	11.8	7.3	1.9	15.4	4.2	2.8	6.4	6.1
1921	20.9	7.4	15.0	16.1	10.5	7.9	2.2	13.2	4.8	4.9	10.0	6.1
1920	25.5	8.4	17.0	18.2	16.8	9.5	2.1	16.1	6.7	6.0	9.8	6.5

See footnotes at end of table.

TABLE 32. - 1/CASE RATES FOR SELECTED COMMUNICABLE DISEASES BY PREFECTURE: JAPAN, 1920-1949 - Cont'd
3/DYSENTERY

Year	All Japan	Hokkaido	Aomori	Iwate	Miyagi	Akita	Yamagata	Fukushima	Ibaraki	Tochigi	Gumma
*1949	29.2	11.7	10.9	47.6	18.6	17.6	31.5	21.9	50.9	38.1	90.9
1948	18.3	23.8	9.9	41.0	14.4	18.3	17.7	23.4	36.7	18.0	29.5
1947	50.5	35.7	25.7	86.9	49.1	36.2	123.3	111.2	85.7	80.6	87.8
1946	116.8	103.4	130.7	541.1	260.8	201.0	339.7	406.7	153.9	163.1	219.9
1945	138.0	33.2	126.9	63.9	19.8	102.2	90.4	101.9	62.2	83.3	169.8
1944	76.0	46.9	96.3	31.7	11.7	47.8	37.7	40.6	76.0	46.4	61.0
1943	69.0	11.4	78.9	88.5	14.3	24.2	23.8	21.2	37.9	25.5	39.5
1942	77.3	3.2	31.5	62.7	12.8	30.7	13.7	35.8	67.6	41.2	40.0
1941	82.0	4.6	37.3	41.3	16.0	22.3	13.9	35.1	83.9	58.0	83.7
1940	118.1	6.2	13.1	61.6	22.2	12.4	11.7	43.4	136.4	85.7	135.2
1939	136.2	18.8	21.5	34.0	27.8	6.2	15.3	44.8	123.3	88.4	75.5
1938	114.4	5.2	15.1	32.9	24.4	7.2	10.2	27.2	101.3	64.8	82.7
1937	112.0	3.7	13.9	32.8	14.2	4.8	12.0	19.7	92.5	39.0	73.3
1936	75.4	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1935	71.8	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1934	63.4	2.5	4.4	29.0	17.8	5.9	3.7	13.3	54.8	28.0	66.9
1933	56.9	1.8	6.7	19.4	9.5	4.4	2.0	8.8	53.7	20.5	34.3
1932	49.0	1.6	4.4	16.8	8.5	4.7	2.4	6.2	46.7	24.4	48.4
1931	45.8	2.4	4.7	12.7	12.0	2.6	1.2	6.8	48.6	24.7	46.8
1930	46.5	1.8	7.0	25.7	9.6	4.0	5.4	8.4	30.3	15.8	46.9
1929	48.1	3.7	6.2	9.6	12.1	3.2	9.1	9.0	44.3	26.5	45.4
1928	46.6	2.4	8.5	14.6	13.3	3.2	8.6	7.0	30.1	17.1	43.9
1927	35.0	5.7	6.1	7.1	7.7	1.9	0.9	2.7	22.7	7.4	33.7
1926	28.5	2.8	9.8	6.5	3.7	1.6	1.5	2.7	24.1	8.2	24.9
1925	24.9	5.2	22.1	23.6	7.3	12.5	2.4	3.3	17.4	5.7	16.7
1924	32.0	1.2	47.2	66.2	9.8	42.6	2.1	9.1	16.0	5.8	21.1
1923	35.2	1.6	71.2	68.5	13.2	9.5	2.6	8.5	21.4	11.7	17.2
1922	26.6	1.9	160.1	85.4	10.0	18.9	0.7	15.4	18.7	2.6	8.1
1921	22.2	1.5	112.8	77.9	12.8	29.5	2.4	19.0	12.8	4.2	5.7
1920	23.0	0.8	90.6	92.6	32.8	50.1	3.0	15.1	15.4	9.7	5.8

See footnotes at end of table.

TABLE 32. - 1/CASE RATES FOR SELECTED COMMUNICABLE DISEASES BY PREFECTURE: JAPAN, 1920-1949 - Cont'd
3/DYSENTERY - Cont'd

Year	Saitama	Chiba	Tokyo	Kanagawa	Niigata	Toyama	Ishikawa	Fukui	Yamanashi	Nagano	Gifu	Shizuoka
1949	82.1	39.3	60.7	41.4	66.4	13.7	18.1	15.2	18.4	24.8	30.3	43.5
1948	19.1	13.3	28.5	12.8	24.9	6.1	3.8	27.7	9.4	16.0	26.3	13.8
1947	82.5	47.8	59.5	32.0	72.7	19.4	22.5	51.0	83.0	77.8	42.9	51.3
1946	95.7	60.8	42.8	76.0	107.7	55.5	23.7	46.9	159.9	100.8	67.6	123.7
1945	115.0	61.7	37.1	79.3	49.4	3.1	3.6	26.0	228.0	201.0	135.3	453.7
1944	91.5	33.6	64.5	135.9	26.9	2.4	6.7	20.1	76.9	58.1	76.0	123.9
1943	63.3	43.1	103.7	90.9	32.9	4.2	3.2	13.7	35.3	28.3	48.3	123.4
1942	67.0	55.2	197.4	172.2	9.8	1.5	4.1	25.8	31.0	8.5	71.2	92.7
1941	96.1	71.8	214.1	181.3	12.8	3.4	3.3	18.9	74.2	9.8	87.7	102.5
1940	126.2	104.0	340.2	284.2	16.9	1.9	6.3	11.6	87.8	13.0	102.4	121.3
1939	101.7	102.7	380.6	253.4	17.0	1.2	5.2	10.7	52.5	11.6	81.7	135.7
1938	103.0	109.9	319.0	182.7	12.5	4.3	4.6	10.4	22.8	9.0	85.1	144.4
1937	78.0	95.3	291.3	133.2	9.3	1.4	3.8	6.7	14.1	7.1	58.1	113.7
1936	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1935	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1934	52.7	40.0	205.2	113.0	8.4	1.5	3.9	8.2	6.7	4.6	52.5	95.7
1933	42.2	42.6	219.2	81.0	10.2	2.6	5.1	7.6	8.9	3.6	43.5	92.3
1932	41.7	31.5	174.8	64.0	3.9	1.3	3.0	7.8	8.5	2.2	55.4	87.2
1931	42.9	28.2	168.9	50.7	3.1	1.0	2.9	2.9	6.4	3.8	50.4	90.5
1930	28.4	23.0	176.6	59.5	2.2	2.3	2.9	2.6	10.0	3.6	37.4	116.8
1929	34.4	47.7	160.7	58.9	1.5	0.6	2.5	3.3	10.5	5.9	41.0	110.7
1928	37.0	29.1	144.0	64.4	1.9	4.9	4.5	3.2	17.3	4.7	38.1	107.6
1927	31.8	19.5	113.0	39.7	9.1	0.4	3.3	3.0	10.1	3.8	34.6	83.4
1926	28.0	17.0	63.4	23.1	1.2	1.7	4.1	1.0	15.7	2.2	32.3	85.0
1925	23.5	11.4	58.9	24.6	1.5	-	3.9	3.2	15.5	2.7	30.9	67.4
1924	21.9	22.4	74.8	21.3	15.0	0.8	2.1	1.3	11.2	6.3	24.5	79.7
1923	25.7	37.7	89.4	35.8	3.3	1.2	3.3	1.2	27.7	4.0	29.2	101.6
1922	11.0	15.5	29.4	22.9	4.4	1.1	2.9	1.7	13.1	2.9	14.7	37.1
1921	9.2	31.2	20.3	17.5	3.1	0.4	7.3	3.8	14.3	1.8	26.3	26.6
1920	16.5	8.5	18.2	23.5	4.4	1.0	17.0	5.7	31.0	1.5	25.9	28.6

See footnotes at end of table.

TABLE 32. - 1/CASE RATES FOR SELECTED COMMUNICABLE DISEASES BY PREFECTURE: JAPAN, 1920-1949 - Cont'd
3/DYSENTERY - Cont'd

Year	Aichi	Mie	Shiga	Kyoto	Osaka	Hyogo	Nara	Wakayama	Tottori	Shimane	Okayama	Hiroshima
1949	35.2	19.7	8.5	21.2	16.5	10.2	6.0	11.2	12.7	33.7	10.2	20.2
1948	21.8	12.5	10.8	16.5	14.9	14.6	5.9	5.8	11.4	24.4	12.2	15.8
1947	60.7	34.8	34.9	49.5	27.6	45.6	22.4	14.7	31.3	52.0	26.3	29.8
1946	64.4	32.7	52.7	58.6	26.6	57.9	33.2	11.3	115.3	85.5	64.6	77.7
1945	131.7	84.9	100.2	81.1	34.7	51.9	92.7	194.4	217.7	237.5	112.7	289.5
1944	43.8	34.3	28.1	73.5	45.3	92.1	52.9	24.9	97.1	114.7	61.8	148.5
1943	56.5	39.3	22.7	98.3	118.1	71.9	11.8	20.3	64.7	47.3	54.2	148.5
1942	85.0	52.8	32.2	119.9	134.7	108.6	18.2	26.2	32.1	44.4	39.2	80.7
1941	74.5	64.0	33.6	121.4	126.4	88.8	25.5	36.9	38.2	56.9	55.8	85.4
1940	94.3	58.7	35.5	153.5	193.6	132.9	38.7	47.8	50.7	77.3	70.1	108.5
1939	87.8	64.1	35.1	171.7	282.3	174.4	47.4	52.1	76.9	150.0	119.0	187.9
1938	88.5	66.4	31.7	139.9	223.4	147.2	29.7	41.3	47.2	68.8	72.0	101.8
1937	67.3	68.1	20.5	133.1	170.7	111.4	15.1	25.9	41.7	54.7	49.0	88.4
1936	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1935	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1934	45.1	38.9	17.0	104.9	134.9	86.3	15.8	14.1	27.2	26.7	43.7	74.4
1933	28.2	36.3	19.0	104.5	117.2	89.0	10.1	21.2	23.0	31.0	37.8	35.9
1932	33.5	23.3	11.9	99.8	102.4	58.2	10.1	10.2	20.7	21.2	21.2	35.2
1931	30.7	23.4	16.5	89.9	86.2	48.9	5.3	8.0	13.8	26.5	20.0	36.4
1930	32.9	24.5	17.5	79.3	75.2	51.4	6.9	7.6	5.7	32.9	21.3	38.2
1929	40.9	25.2	25.8	109.1	81.6	45.6	9.4	9.1	8.1	31.7	35.1	37.4
1928	28.4	35.7	12.3	102.0	56.7	33.7	2.2	7.7	2.9	27.0	25.0	36.2
1927	26.6	33.0	8.6	80.9	49.2	46.1	3.2	9.9	3.7	24.3	33.0	42.0
1926	22.0	31.0	12.3	76.6	43.1	32.4	1.5	13.0	12.1	28.6	34.2	30.3
1925	24.7	30.3	12.2	61.3	26.7	27.1	3.4	3.7	5.9	22.1	26.6	23.5
1924	30.1	20.0	6.6	55.7	24.1	26.9	1.4	10.6	4.3	29.3	33.3	40.8
1923	32.4	30.1	5.5	61.5	25.3	34.9	4.7	6.4	5.2	35.7	27.4	38.0
1922	12.3	6.9	2.5	22.6	23.6	42.5	0.7	5.6	9.1	40.6	40.6	70.7
1921	14.9	15.4	3.8	31.6	16.2	21.2	13.6	4.9	2.8	11.0	31.1	69.7
1920	13.7	13.7	8.9	29.1	12.1	18.4	8.3	10.1	2.6	11.9	15.4	41.1

See footnotes at end of table.

TABLE 32. - 1/CASE RATES FOR SELECTED COMMUNICABLE DISEASES BY PREFECTURE: JAPAN, 1920-1949 - Cont'd
3/DISENTERY - Cont'd

Year	Yamaguchi	Tokushima	Kagawa	Ehime	Kochi	Fukuoka	Saga	Nagasaki	Kumamoto	Oita	Miyazaki	Kagoshima
*1949	16.5	19.2	25.4	20.5	13.7	13.1	12.3	13.3	11.4	14.7	36.7	9.2
1948	16.2	17.9	13.8	24.4	12.9	13.1	19.7	16.1	12.3	15.4	26.0	8.8
1947	19.5	98.9	57.2	67.9	36.3	20.1	23.2	35.5	20.0	26.1	52.5	40.7
1946	80.0	170.4	221.0	145.4	148.3	77.2	130.0	119.4	111.1	82.4	211.4	76.2
1945	246.9	300.5	556.3	327.0	170.8	378.8	237.1	272.3	215.9	242.5	181.1	231.9
1944	122.9	142.0	462.4	113.5	136.0	147.9	113.8	133.0	71.6	66.7	138.4	28.3
1943	135.4	348.9	213.0	96.2	71.6	64.8	17.8	48.4	68.4	30.3	54.9	15.6
1942	126.0	114.8	90.5	36.1	65.6	88.5	29.8	42.8	61.1	28.5	49.8	14.7
1941	99.3	124.6	115.3	40.9	72.8	74.5	42.9	50.5	86.8	51.1	34.4	18.7
1940	181.1	152.0	205.0	46.7	102.2	116.7	37.8	82.8	90.3	72.7	63.0	22.8
1939	193.2	195.1	207.2	65.6	111.6	200.8	35.1	97.5	122.8	69.5	55.3	62.3
1938	161.0	155.9	123.7	47.4	156.9	189.7	56.4	60.4	136.4	41.5	51.2	31.3
1937	137.4	123.9	96.9	35.0	117.1	544.5	28.9	63.9	130.1	42.4	44.0	43.9
1936	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1935	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1934	96.3	109.9	86.0	27.0	58.8	52.5	6.0	32.2	64.2	21.0	41.4	41.3
1933	89.9	84.1	67.1	19.1	53.8	32.8	6.6	36.3	58.6	13.6	28.6	14.5
1932	72.2	115.5	54.3	19.5	58.1	34.2	9.1	34.3	69.1	22.2	31.3	9.0
1931	76.7	97.2	62.2	20.8	36.5	31.8	6.9	38.2	65.7	20.8	20.8	19.6
1930	92.4	74.5	44.2	24.8	44.4	41.1	8.1	53.5	71.9	40.1	25.8	9.3
1929	121.5	63.5	61.6	25.2	36.9	46.1	7.3	52.9	71.5	17.9	41.3	11.9
1928	72.9	88.1	61.5	25.4	42.3	33.8	4.0	38.9	65.3	13.0	29.2	5.3
1927	74.2	89.5	57.2	21.0	32.9	32.9	6.5	30.4	55.3	20.5	21.2	3.5
1926	67.3	75.4	63.9	21.9	32.6	32.1	11.7	35.3	52.0	28.7	15.1	7.5
1925	60.9	73.8	59.0	17.0	22.8	25.9	7.4	31.7	52.7	16.7	12.9	8.1
1924	52.9	156.2	96.8	16.3	30.5	60.0	31.9	35.9	78.7	17.9	22.2	4.3
1923	62.6	208.5	56.0	30.7	24.3	44.2	29.2	44.8	62.0	44.3	18.5	3.8
1922	28.6	85.9	110.1	19.7	26.2	28.3	25.3	48.2	59.2	95.0	15.0	8.5
1921	23.0	30.5	55.2	11.0	12.0	16.0	6.1	73.7	83.2	47.5	14.2	8.6
1920	22.2	60.0	103.6	18.8	11.9	23.1	14.7	58.7	79.1	46.4	13.5	8.8

See footnotes at end of table.

TABLE 32. - 1/CASE RATES FOR SELECTED COMMUNICABLE DISEASES BY PREFECTURE: JAPAN, 1920-1949 - Cont'd
SMALLPOX

Year	All Japan	Hokkaido	Aomori	Iwate	Miyagi	Akita	Yamagata	Fukushima	Ibakeki	Tochigi	Gumma
*1949	0.2	-	-	-	-	0.1	-	-	-	-	-
1948	0.0	0.2	-	-	-	-	-	-	0.0	-	-
1947	0.5	1.3	-	0.1	0.1	1.0	0.6	0.1	1.0	1.5	0.2
1946	23.7	64.8	6.7	3.9	7.5	5.8	9.0	11.6	9.2	4.8	5.8
1945	2.3	12.3	0.1	0.9	2.6	0.4	-	9.9	1.0	-	0.5
1944	0.4	1.2	-	-	0.6	-	-	1.8	-	0.2	0.2
1943	0.8	0.1	0.1	1.5	0.8	0.1	0.1	-	2.5	0.3	0.2
1942	0.5	0.1	-	0.1	-	0.2	-	1.2	0.1	-	-
1941	0.9	6.2	5.9	0.2	1.2	1.4	0.1	0.7	0.2	-	-
1940	0.8	0.5	1.1	2.2	2.9	0.1	1.2	0.1	-	-	0.3
1939	0.4	0.0	-	-	0.2	-	-	-	0.6	0.7	-
1938	0.1	-	-	-	-	-	-	-	-	-	-
1937	0.1	-	-	-	-	-	-	-	-	-	-
1936	0.3	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1935	0.2	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1934	0.5	0.4	0.5	-	-	-	0.1	-	-	-	-
1933	0.6	-	-	-	-	3.5	-	0.1	-	2.0	0.2
1932	0.5	0.1	-	0.2	-	-	-	-	-	0.2	-
1931	0.0	-	-	-	-	-	-	-	-	-	-
1930	0.0	-	-	-	-	-	-	-	-	-	-
1929	0.2	0.0	-	-	-	-	-	-	-	-	-
1928	1.2	0.1	-	0.1	-	-	-	1.3	-	0.2	-
1927	0.6	0.2	-	-	-	-	-	0.1	-	-	-
1926	2.1	0.4	-	0.1	-	-	-	-	-	-	-
1925	0.7	0.0	-	0.2	-	-	-	-	-	0.1	9.9
1924	2.9	19.7	1.5	4.4	5.7	6.6	7.9	2.9	0.6	2.9	0.2
1923	3.3	9.2	0.9	-	0.3	2.1	-	-	0.1	-	-
1922	1.2	1.0	-	-	0.3	1.0	-	0.5	1.0	0.4	-
1921	1.6	-	-	0.1	-	-	0.1	0.4	0.9	0.7	1.9
1920	5.7	1.4	0.3	0.6	-	12.4	3.4	2.7	0.7	0.1	0.2

See footnotes at end of table.

TABLE 33. - 1/CASE RATES FOR SELECTED COMMUNICABLE DISEASES BY PREFECTURE: JAPAN, 1920-1949 - Cont'd
SMALLPOX - Cont'd

Year	Saitama	Chiba	Tokyo	Kanagawa	Miyagi	Toyama	Ishikawa	Fukui	Yamanashi	Nagano	Gifu	Shizuoka
*1949	-	-	0.1	0.1	0.0	-	-	-	-	-	-	-
1948	-	0.1	-	-	-	-	-	-	-	-	0.1	0.0
1947	0.1	0.6	0.4	0.2	0.2	0.1	0.1	-	-	0.1	-	0.2
1946	9.9	10.5	42.3	20.5	5.6	19.5	13.3	20.1	10.6	9.0	16.8	10.8
1945	2.3	0.1	0.3	2.3	0.8	1.0	0.8	0.3	-	2.2	-	3.1
1944	-	0.1	0.2	-	0.8	1.5	8.7	0.3	-	-	-	-
1943	-	-	0.5	0.2	0.1	-	1.1	0.5	0.3	0.1	0.2	0.2
1942	-	0.1	0.5	0.1	-	-	-	3.9	0.8	0.8	-	-
1941	0.1	1.3	0.3	0.3	3.2	1.2	0.1	0.3	0.2	0.1	0.1	0.2
1940	0.7	0.1	0.5	0.2	1.8	0.4	0.9	0.2	0.6	0.8	1.1	1.0
1939	-	0.1	0.1	0.1	-	-	-	-	-	-	-	-
1938	-	-	0.0	-	-	-	-	-	-	-	-	-
1937	-	0.1	-	0.0	-	0.3	-	-	-	1.5	-	0.3
1936	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1935	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1934	0.1	-	0.0	1.7	-	-	-	-	-	-	-	0.5
1933	0.1	0.1	1.1	0.1	-	0.1	0.1	0.2	0.3	-	-	-
1932	0.1	-	-	0.2	-	-	0.3	-	-	-	-	1.0
1931	-	0.1	-	0.1	-	-	-	-	-	-	-	-
1930	-	-	0.0	-	-	-	0.5	-	-	-	-	-
1929	-	-	0.0	0.1	0.7	0.3	0.3	-	-	-	-	0.1
1928	-	0.2	1.3	0.8	0.1	0.1	0.7	-	1.6	-	-	0.1
1927	-	0.4	-	0.5	-	-	-	-	-	-	-	-
1926	-	0.1	0.1	4.9	-	0.9	0.4	-	1.0	-	-	0.1
1925	-	0.1	0.0	0.5	-	-	-	-	-	-	-	-
1924	0.4	0.1	4.6	0.5	0.1	2.8	2.0	1.7	2.6	0.2	1.4	0.4
1923	-	0.2	0.5	0.4	0.1	-	0.1	0.2	-	0.1	0.5	0.1
1922	-	-	0.4	1.0	-	5.5	-	2.5	0.2	-	-	-
1921	0.1	-	0.1	0.4	-	-	-	-	-	-	-	0.1
1920	0.2	0.3	3.2	0.1	1.1	-	1.7	2.5	-	0.1	0.3	1.4

See footnotes at end of table.

TABLE 32. - 1/CASE RATES FOR SELECTED COMMUNICABLE DISEASES BY PREFECTURE: JAPAN, 1920-1949 - Cont'd
SMALLPOX - Cont'd

Year	Aichi	Mie	Shiga	Kyoto	Osaka	Iworo	Nara	Wakayama	Tottori	Shimane	Okayama	Hiroshima
*1949	-	0.1	-	-	1.7	-	0.1	0.9	0.5	-	0.1	-
1948	-	-	-	0.1	0.1	-	-	0.1	-	0.1	0.1	0.0
1947	0.3	0.4	-	0.1	0.3	1.4	0.1	3.6	0.2	0.8	0.7	0.1
1946	39.1	16.4	18.9	41.7	79.3	74.2	52.9	23.6	8.6	14.5	14.5	20.3
1945	1.6	-	1.9	0.2	1.5	10.7	0.3	-	0.6	4.0	0.5	0.8
1944	0.4	0.8	-	-	0.1	0.1	-	0.8	-	0.1	-	-
1943	-	-	-	0.1	0.4	7.0	0.3	0.8	0.2	-	0.5	-
1942	0.1	-	-	0.1	0.0	0.2	-	0.6	-	-	6.6	0.3
1941	0.0	0.9	0.1	-	1.0	0.2	0.3	-	13.0	-	0.1	0.5
1940	0.3	-	-	0.2	0.9	0.3	2.1	0.1	0.4	-	0.7	7.3
1939	0.1	-	-	1.4	0.3	3.4	-	-	-	-	0.2	0.3
1938	0.0	-	-	0.2	-	0.1	-	-	-	-	0.5	-
1937	-	-	-	-	0.0	0.0	-	-	-	-	-	-
1936	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1935	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1934	-	-	1.1	-	0.3	0.1	-	-	0.2	-	0.7	1.7
1933	0.6	0.1	0.3	0.7	0.4	0.1	-	-	3.0	-	-	-
1932	-	0.1	-	-	5.4	0.4	0.2	0.1	-	-	0.1	-
1931	0.2	-	-	-	-	0.1	-	-	-	-	-	-
1930	-	-	-	-	-	-	-	-	-	-	-	-
1929	-	-	-	-	0.1	0.0	-	-	-	9.1	0.1	-
1928	0.5	0.3	0.1	0.2	1.4	0.7	0.3	0.1	-	-	0.3	0.2
1927	-	-	-	-	0.1	0.2	-	-	0.6	-	-	-
1926	0.7	3.1	-	0.4	0.9	1.2	0.5	0.1	-	0.1	-	1.6
1925	-	-	-	0.2	1.8	0.4	-	-	0.2	0.1	0.2	0.6
1924	2.0	2.2	3.2	0.8	0.7	1.4	-	0.4	-	-	-	-
1923	0.1	-	0.6	0.4	8.2	4.4	0.5	-	2.8	0.3	0.6	23.7
1922	-	-	0.6	0.7	0.2	0.7	0.9	-	1.5	0.1	0.5	0.9
1921	0.6	0.1	-	0.2	3.4	4.1	0.7	-	-	-	0.5	0.1
1920	0.5	2.2	9.5	4.4	46.7	3.7	24.3	17.7	1.3	3.1	2.2	4.8

See footnotes at end of table.

TABLE 32. - 1/CASE RATES FOR SELECTED COMMUNICABLE DISEASES BY PREFECTURE: JAPAN, 1920-1949 - Cont'd
SMALLPOX - Cont'd

Year	Yamaguchi	Tokushima	Kagawa	Ehime	Kochi	Fukuoka	Saga	Nagasaki	Kumamoto	Oita	Miyazaki	Kagoshima
1949	0.8	-	-	0.3	-	0.6	-	0.1	-	0.2	-	-
1948	-	-	-	-	-	0.0	0.5	-	-	-	-	-
1947	6.5	0.1	0.4	0.9	0.1	1.3	0.5	0.1	0.2	0.2	0.1	1.0
1946	9.0	20.3	32.3	11.3	22.5	14.3	6.1	22.2	5.0	7.7	4.2	10.3
1945	6.1	-	1.1	0.1	-	1.1	0.6	8.5	0.2	0.3	0.1	0.2
1944	0.2	-	0.3	0.2	-	0.3	0.1	1.1	1.3	0.3	-	1.8
1943	2.7	0.6	0.8	0.8	1.0	0.7	-	0.4	1.2	2.3	0.1	0.7
1942	0.2	-	0.3	-	0.1	0.3	0.4	2.2	-	0.2	6.7	3.6
1941	0.6	1.1	0.1	0.7	1.9	0.3	0.4	0.3	-	-	0.1	0.1
1940	2.6	-	0.3	1.1	0.1	0.4	0.3	0.5	1.0	0.8	0.4	0.2
1939	3.5	7.2	0.3	-	0.4	0.1	-	-	-	0.1	0.1	-
1938	-	-	-	-	-	0.1	4.2	0.5	0.1	-	0.2	0.2
1937	1.4	-	-	-	-	0.8	0.9	0.1	-	0.3	-	-
1936	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1935	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1934	1.6	-	-	1.9	-	0.2	0.3	0.1	7.3	2.6	-	1.2
1933	0.3	-	-	1.7	0.3	0.5	-	0.1	0.5	1.2	0.6	7.0
1932	0.1	0.1	1.1	-	0.3	0.3	-	1.1	0.4	0.3	1.3	0.2
1931	-	-	-	-	-	0.4	-	-	0.2	0.1	-	-
1930	-	-	0.1	-	-	-	-	0.1	-	-	-	-
1929	-	-	-	-	0.1	0.1	-	0.4	-	-	-	0.7
1928	8.6	0.3	-	0.4	5.7	4.3	0.1	0.4	6.7	16.2	0.7	-
1927	0.4	-	-	-	0.1	10.7	0.3	4.4	0.1	1.6	-	-
1926	3.8	5.2	23.9	0.2	1.0	16.7	5.4	7.9	4.1	13.8	0.7	3.5
1925	6.3	-	-	0.1	-	1.7	-	9.0	-	0.4	0.6	-
1924	3.5	-	1.0	0.1	-	2.9	-	-	2.5	20.2	0.1	3.6
1923	9.0	10.8	4.6	2.7	-	9.9	7.3	5.0	8.4	17.4	3.4	0.3
1922	7.9	-	0.1	2.3	-	10.4	2.1	5.9	1.2	1.9	0.1	0.1
1921	3.3	1.2	12.7	0.5	1.6	5.3	0.6	1.4	1.4	2.7	0.2	0.1
1920	7.0	12.2	2.4	3.2	19.7	11.5	0.1	1.7	1.8	25.6	1.8	2.0

See footnotes at end of table.

TABLE 32. - 1/CASE RATES FOR SELECTED COMMUNICABLE DISEASES BY PREFECTURE: JAPAN, 1920-1949 - Cont'd
TYPHUS FEVER

Year	All Japan	Hokkaido	Aomori	Iwate	Miyagi	Akita	Yamagata	Fukushima	Ibaraki	Tochigi	Gumma
*1949	0.1	0.0	-	-	0.8	-	0.1	0.5	-	-	-
1948	0.6	0.1	1.5	0.5	0.3	0.4	0.3	0.2	0.2	-	-
1947	1.4	1.5	0.7	-	1.3	0.2	3.2	0.2	1.8	0.7	0.3
1946	41.5	68.3	75.4	9.6	31.0	17.8	39.7	14.4	14.1	10.3	6.9
1945	3.5	52.7	1.6	2.9	0.2	-	3.8	0.5	1.6	-	1.4
1944	5.4	62.8	10.5	1.5	12.8	6.1	7.9	0.9	5.4	0.9	-
1943	1.9	33.0	1.2	0.5	2.5	6.5	5.1	-	-	-	-
1942	0.1	-	0.1	-	4.9	-	0.5	-	-	-	0.1
1941	0.1	-	-	-	2.5	-	1.8	-	-	0.1	-
1940	0.0	-	-	-	-	-	-	-	-	-	-
1939	0.0	-	-	-	-	-	-	-	-	-	-
1938	-	-	-	-	-	-	-	-	-	-	-
1937	0.0	-	1.7	-	-	-	-	-	-	-	-
1936	0.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1935	0.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1934	0.0	-	2.7	-	-	-	-	-	-	-	-
1933	0.0	-	-	-	-	-	-	-	-	-	-
1932	0.0	-	-	-	-	-	-	-	-	-	-
1931	0.0	-	0.1	-	-	-	0.1	-	-	-	-
1930	0.0	-	-	-	-	-	-	-	-	-	-
1929	0.0	-	-	-	0.7	-	0.5	-	-	-	-
1928	0.0	-	-	-	0.5	-	1.6	0.1	-	-	-
1927	0.0	-	-	-	0.2	-	0.4	-	-	-	-
1926	0.1	-	-	-	-	-	3.3	-	-	-	-
1925	0.0	-	-	-	1.7	-	1.0	-	-	-	-
1924	0.0	-	1.1	-	-	0.2	-	-	-	-	0.1
1923	0.0	0.3	-	-	0.4	-	-	-	-	-	-
1922	0.0	-	0.8	-	-	0.1	1.4	0.1	-	-	-
1921	0.3	-	0.4	-	8.5	1.5	7.2	-	-	-	-
1920	0.1	0.1	4.1	-	-	0.8	2.5	0.1	-	-	-

See footnotes at end of table.

TABLE 32. - 1/CASE RATES FOR SELECTED COMMUNICABLE DISEASES BY PREFECTURE: JAPAN, 1920-1949 - Cont'd
TYPHUS FEVER - Cont'd

Year	Saitama	Chiba	Tokyo	Kanagawa	Niigata	Toyama	Ishikawa	Fukui	Yamanashi	Nagano	Gifu	Shizuoka
1949	0.1	0.0	0.4	0.5	-	-	-	-	-	-	-	-
1948	0.5	0.3	1.0	2.0.	0.1	0.6	0.1	-	-	0.0	0.7	-
1947	1.5	1.2	4.7	2.2	0.5	0.8	1.1	0.8	0.9	0.4	1.8	1.3
1946	33.7	16.5	222.5	43.6	5.3	4.2	8.7	8.0	17.1	11.6	5.8	3.7
1945	-	0.1	0.4	3.6	0.3	-	-	-	-	-	-	1.1
1944	0.1	0.2	11.3	2.4	-	-	-	0.3	-	0.1	-	-
1943	0.1	0.1	0.8	-	-	-	-	0.3	0.2	-	-	-
1942	-	-	0.3	-	-	-	-	-	-	-	-	-
1941	0.6	-	0.3	-	-	-	-	-	-	-	-	-
1940	-	-	0.0	-	0.0	-	0.1	-	-	-	-	-
1939	-	-	-	-	-	-	-	-	-	-	-	-
1938	-	-	-	-	-	-	-	-	-	-	-	-
1937	-	-	-	-	-	-	-	-	-	-	-	-
1936	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1935	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1934	-	-	-	-	-	-	-	-	-	-	-	-
1933	-	-	-	-	-	-	-	-	-	-	-	-
1932	-	-	-	-	0.1	0.1	-	-	-	-	-	-
1931	-	-	-	-	-	0.1	-	-	-	-	-	-
1930	-	-	-	-	-	0.1	-	-	-	-	-	-
1929	-	-	-	-	-	-	-	-	-	-	-	-
1928	-	-	-	-	-	-	-	-	-	-	-	-
1927	-	-	-	-	-	-	-	-	-	-	-	-
1926	-	-	-	-	-	0.3	-	-	-	-	-	-
1925	-	-	-	-	-	-	-	-	-	-	-	-
1924	-	-	-	-	-	-	-	-	-	-	-	0.1
1923	-	-	-	-	-	-	-	-	-	-	-	-
1922	-	-	-	-	-	-	-	-	-	-	-	-
1921	-	-	-	-	-	-	-	-	-	-	-	-
1920	-	-	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

TABLE 32. - 1/CASE RATES FOR SELECTED COMMUNICABLE DISEASES BY PREFECTURE: JAPAN, 1920-1949 - Cont'd
TYPHUS FEVER - Cont'd

Year	Aichi	Mie	Shizu	Yvoto	Osaka	Hyogo	Nara	Wakayama	Tottori	Shimane	Okayama	Hiroshima
*1949	-	0.1	-	-	0.7	0.0	-	0.4	-	1.1	-	0.0
1948	0.1	0.1	0.1	2.1	4.3	0.1	2.7	-	-	1.3	-	0.6
1947	7.3	0.4	-	0.4	2.7	0.2	0.3	1.9	1.2	1.3	0.3	0.1
1946	37.4	6.0	7.4	63.8	208.4	91.5	24.6	9.9	7.8	3.0	8.2	3.1
1945	0.2	-	-	0.5	-	1.6	-	-	-	-	0.9	-
1944	0.0	-	-	0.1	1.2	0.3	-	-	-	-	-	-
1943	0.0	-	-	0.1	0.0	0.9	-	-	0.4	0.1	-	-
1942	0.1	-	-	-	-	0.0	-	-	-	-	-	-
1941	-	-	-	-	-	0.1	-	-	-	-	0.2	-
1940	-	-	-	-	-	-	-	-	-	-	-	-
1939	-	-	-	-	-	-	-	-	-	-	-	-
1938	-	-	-	-	-	-	-	-	-	-	-	-
1937	-	-	-	-	-	-	-	-	-	-	-	-
1936	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1935	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1934	-	-	-	-	-	0.0	-	-	-	-	-	-
1933	-	-	-	-	0.0	0.1	-	-	-	-	-	-
1932	-	-	-	-	-	-	-	-	-	-	-	0.1
1931	-	-	-	-	-	-	-	-	-	-	-	-
1930	-	-	-	-	-	-	-	-	-	-	-	-
1929	-	-	-	-	-	-	-	0.2	-	-	-	-
1928	-	-	-	-	-	-	-	-	-	-	-	-
1927	-	-	-	-	-	-	-	-	-	-	-	0.1
1926	-	-	-	-	-	-	-	-	-	-	-	-
1925	-	-	-	-	-	-	-	-	-	-	-	-
1924	-	-	-	-	-	-	-	-	-	-	-	-
1923	-	-	-	-	-	0.0	-	-	-	-	-	-
1922	-	-	-	-	-	-	-	-	0.2	-	-	-
1921	-	-	-	-	-	-	-	-	-	-	0.1	0.1
1920	-	-	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

TABLE 32. - 1/CASE RATES FOR SELECTED COMMUNICABLE DISEASES BY PREFECTURE: JAPAN, 1920-1949 - Cont'd
TYPHUS FEVER - Cont'd

Year	Yamaguchi	Tokushima	Kagawa	Ehime	Kochi	Fukuoka	Saga	Nagasaki	Kumamoto	Oita	Miyazaki	Kagoshima
1949	-	-	-	-	-	0.0	-	0.7	0.1	-	-	-
1948	0.1	-	0.9	-	-	0.1	-	1.7	0.2	-	-	-
1947	1.1	0.2	5.9	0.4	0.2	0.1	0.2	0.5	0.2	0.1	0.7	-
1946	6.5	2.6	8.5	2.5	5.2	16.1	4.6	30.1	0.6	2.9	2.2	3.0
1945	0.2	-	1.0	0.4	-	13.5	3.3	0.2	-	-	-	-
1944	0.1	-	-	-	-	2.8	8.5	-	-	-	0.2	13.0
1943	0.2	-	-	0.1	-	0.1	0.3	0.1	-	-	-	0.1
1942	-	-	-	-	-	-	-	0.1	-	-	-	-
1941	-	-	-	-	-	-	-	-	0.1	-	-	-
1940	-	-	-	-	-	-	-	-	-	-	-	-
1939	-	-	-	-	-	0.2	-	-	-	-	-	-
1938	-	-	-	-	-	-	-	-	-	-	-	-
1937	-	-	0.1	-	-	-	-	-	-	-	-	-
1936	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1935	NA	NA	NA	NA	NA	N/	NA	NA	NA	NA	NA	NA
1934	-	-	-	-	-	-	-	-	-	-	-	-
1933	-	-	-	-	-	-	-	-	-	-	-	-
1932	-	-	-	0.1	-	-	-	-	-	-	-	-
1931	-	-	-	-	-	-	-	-	-	-	-	-
1930	-	-	-	-	-	-	-	-	-	-	-	-
1929	-	-	-	-	-	-	-	-	-	-	-	-
1928	-	-	-	-	-	-	-	-	-	-	-	-
1927	-	-	-	-	-	-	-	-	-	-	-	-
1926	-	-	-	-	-	-	-	-	-	-	-	-
1925	-	-	-	-	-	-	-	-	-	-	-	-
1924	-	-	-	-	-	-	-	-	-	-	-	-
1923	-	-	-	-	-	-	-	-	-	-	-	-
1922	-	-	-	-	-	-	-	-	-	-	-	-
1921	-	-	-	-	-	-	-	-	-	-	-	-
1920	-	-	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

TABLE 32. - 1/CASE RATES FOR SELECTED COMMUNICABLE DISEASES BY PREFECTURE: JAPAN, 1920-1949 - Cont'd

4/MALARIA

Year	All Japan	Hokkaido	Aomori	Iwate	Miyagi	Akita	Yamagata	Fukushima	Ibaraki	Tochigi	Gumma
1949	4.5	1.2	4.3	1.3	0.9	1.2	2.2	1.2	1.8	1.5	0.7
1948	6.2	2.8	3.3	2.6	2.1	2.0	1.8	2.1	2.4	2.5	0.9
1947	15.2	7.7	15.4	14.6	1.6	14.9	8.9	12.5	15.8	7.5	5.7
1920 - 1946	Not Available										

5/JAPANESE "B" ENCEPHALITIS

Year	All Japan	Hokkaido	Aomori	Iwate	Miyagi	Akita	Yamagata	Fukushima	Ibaraki	Tochigi	Gumma
1949	1.6	0.0	-	-	0.2	-	0.1	-	1.5	0.1	1.3
1948	9.0	0.2	3.8	10.9	10.4	10.9	11.4	3.3	16.5	5.7	8.9
1947	0.3	-	0.2	-	0.1	0.2	0.1	-	-	0.1	0.1
1920 - 1946	Not Available										

See footnotes at end of table.

TABLE 32. 1/CASE RATES FOR SELECTED COMMUNICABLE DISEASES BY PREFECTURE: JAPAN, 1920-1949 - Cont'd

4/MALARIA - Cont'd

Year	Saitama	Chiba	Tokyo	Kanagawa	Niigata	Toyama	Ishikawa	Fukui	Yamanashi	Nagano	Gifu	Shizuoka
1949	1.6	1.9	2.4	1.6	1.8	2.2	1.9	4.3	1.7	1.1	1.7	1.0
1948	1.2	1.6	5.8	3.6	4.3	5.0	3.4	3.4	2.8	0.8	2.8	1.8
1947	3.2	5.5	15.4	20.5	12.6	17.1	6.5	10.1	8.5	9.1	2.6	8.9
1920 - 1946	Not Available.											

5/JAPANESE "B" ENCEPHALITIS - Cont'd

Year	Saitama	Chiba	Tokyo	Kanagawa	Niigata	Toyama	Ishikawa	Fukui	Yamanashi	Nagano	Gifu	Shizuoka
1949	0.7	0.3	3.9	4.8	0.4	2.3	2.2	-	0.2	1.4	0.4	2.0
1948	14.1	15.8	36.5	25.6	9.9	10.6	8.6	8.9	13.7	9.2	7.1	15.7
1947	-	-	0.1	0.0	0.0	0.1	-	0.1	-	-	0.1	-
1920 - 1946	Not Available.											

See footnotes at end of table.

TABLE 32. - 1/CASE RATES FOR SELECTED COMMUNICABLE DISEASES BY PREFECTURE: JAPAN, 1920-1949 - Cont'd

4/MALARIA - Cont'd

Year	Aichi	Mie	Shiga	Kyoto	Osaka	Hyogo	Nara	Wakayama	Tottori	Shimane	Okayama	Hiroshima
*1949	2.1	2.2	246.0	7.2	0.7	1.7	1.8	1.1	4.3	1.6	1.8	2.2
1948	1.5	4.2	260.1	4.8	1.6	2.1	4.0	2.1	8.0	3.8	3.1	5.4
1947	8.5	16.0	220.0	10.0	4.8	10.4	9.5	8.6	26.8	14.7	4.9	11.8
1920 - 1946	Not Available.											

5/JAPANESE "B" ENCEPHALITIS - Cont'd

Year	Aichi	Mie	Shiga	Kyoto	Osaka	Hyogo	Nara	Wakayama	Tottori	Shimane	Okayama	Hiroshima
*1949	2.3	2.4	0.7	1.4	2.9	0.4	2.9	-	0.3	-	2.5	4.0
1948	12.1	5.8	7.1	1.9	2.9	4.3	1.9	1.8	4.6	2.0	4.2	1.7
1947	-	0.4	-	0.3	1.4	0.4	-	0.1	3.8	0.8	4.3	0.3
1920 - 1946	Not Available.											

See footnotes at end of table.

TABLE 32. - 1/CASE RATES FOR SELECTED COMMUNICABLE DISEASES BY PREFECTURE: JAPAN, 1920-1949 - Cont'd

4/MALARIA - Cont'd

Year	Yamaguchi	Tokushima	Kagawa	Ehime	Kochi	Fukuoka	Saga	Nagasaki	Kumamoto	Oita	Miyazaki	Kagoshima
1949	2.0	1.1	2.8	2.2	1.8	2.0	1.7	2.5	1.9	1.6	2.2	1.7
1948	3.1	2.5	2.6	7.3	2.4	7.5	4.4	4.4	3.5	4.4	2.2	9.0
1947	19.1	25.8	16.0	33.6	13.1	31.5	30.3	14.2	11.8	30.4	19.6	16.4
1920 - 1946	Not Available.											

5/JAPANESE "B" ENCEPHALITIS - Cont'd

Year	Yamaguchi	Tokushima	Kagawa	Ehime	Kochi	Fukuoka	Saga	Nagasaki	Kumamoto	Oita	Miyazaki	Kagoshima
1949	4.9	1.0	1.3	1.1	1.1	0.4	2.0	0.2	4.7	0.9	2.4	2.0
1948	2.0	3.0	1.9	4.3	3.9	1.4	2.8	2.1	2.6	0.9	6.3	4.1
1947	-	0.1	3.4	1.1	1.5	0.0	-	0.1	0.1	0.1	0.1	-
1920 - 1946	Not Available.											

See footnotes at end of table.

TABLE 32. - 1/CASE RATES FOR SELECTED COMMUNICABLE DISEASES BY PREFECTURE: JAPAN, 1920-1949 - Cont'd

FOOTNOTES:

* Data for 1949 are provisional.

1/ Rates are the number of cases per 100,000 population and are computed on an annual basis. Case data refer to the civilian population in Japan Proper, 1900-1934 and 1937-1949. The totals for 1935 and 1936 include reports from Okinawa, but case data for Okinawa are very incomplete. Prefectural case data for 1935 and 1936 are not available. Rates for all the years 1920-1949 are based upon the estimated civilian population in Japan Proper.

2/ Diphtheria includes croup, 1920-1923.

3/ Dysentery includes ekiri, 1922-1949.

4/ Annual data for malaria are not available prior to 1947.

5/ Annual data for Japanese "B" encephalitis are not available prior to 1947.

"NA" indicates that data are not available.

A dash (-) indicates that no cases were reported.

A rate of 0.0 indicates that some cases were reported but the rate was less than 0.1.

Sources:

Rates were computed by Public Health and Welfare Section, GHQ, SCAP. Sources of case data are as follows:

1920-1923: Statistics of Notifiable Infectious Diseases, Sanitary Bureau, Home Department, 1924.

1924-1934: Statistics of Notifiable Infectious Diseases, Sanitary Bureau, Home Department, 1937.

1935-1942: Annual Reports, Ministry of Welfare.

1943-1945: Unpublished Annual Reports, Ministry of Welfare.

1946-1949: Weekly Reports, Ministry of Welfare.

TABLE 33. - NATURAL INCREASE IN POPULATION: JAPAN, 1947-1949

Year	Births	Deaths	Net Natural Increase	Rates of Net Natural Increase per 1,000 Population
*1949	2,723,873	950,719	1,773,154	21.6
*1948	2,709,871	950,610	1,759,261	21.9
1947	2,676,792	1,136,238	1,540,554	19.8

Footnotes:

- * Birth data for 1948 and 1949 are provisional. Death data for 1949 are provisional. Rates for 1948 and 1949 are provisional. All data refer to events occurring to Japanese nationals in Japan. Data for 1947 and death data for 1948 are from Final Annual Report. Ministry of Welfare. Data for 1949 and birth data for 1948 are from Monthly Schedule Reports, Ministry of Welfare. Rates were computed by Public Health and Welfare Section, GHQ, SCAP.

This Table corresponds to Table 27, Statistical Annex, Public Health and Welfare Section, GHQ, SCAP.

TABLE 34. - INFANT DEATHS BY MONTH: JAPAN PROPER, 1947-1949

Year	Total	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Unknown
*1949	168,989	19,706	17,277	19,040	15,994	13,794	12,145	12,049	10,473	9,060	9,764	12,918	16,819	
1948	185,406	20,845	19,556	18,744	13,385	11,305	11,397	12,158	10,714	9,555	9,976	12,306	15,858	7
1947	205,360	21,088	24,154	23,936	17,965	15,096	13,816	16,416	15,406	11,081	11,396	14,664	20,341	1

Footnotes: * Data for 1949 are provisional. Infant deaths refer to deaths under 1 year of age.
Data for 1947 and 1948 from Final Annual Report, Ministry of Welfare, and for 1949 from Monthly Schedule Reports, Ministry of Welfare.

This Table corresponds to Table 28, Statistical Annex, Public Health and Welfare in Japan, 1948.

TABLE 35. - INFANT DEATH RATES BY MONTH: JAPAN PROPER, 1947-1949
(Rates per 1,000 live births each month)

Year	Annual	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
*1949	62.0	60.4	70.7	76.3	72.2	67.7	63.9	56.9	47.7	40.8	44.4	61.2	82.1
*1948	61.0	66.2	75.1	71.3	59.7	56.1	60.7	59.1	49.8	44.4	45.8	56.1	83.3
1947	76.7	71.4	106.9	101.5	85.9	77.2	71.0	72.5	65.1	47.8	49.8	69.6	108.8

Footnotes: * Data for 1948 and 1949 are provisional. Rates are the number of infant deaths per 1,000 live births in the corresponding period and were computed by Public Health and Welfare Section, GHQ, SCAP. Base data for 1947 and infant death data for 1948 are from Monthly Schedule Reports, Ministry of Welfare.

This Table corresponds to Table 29, Statistical Annex, Public Health and Welfare in Japan, 1948.

TABLE 36. - INFANT DEATHS BY PREFECTURE: JAPAN, 1947-1949

Year	All Japan	Aichi	Akita	Aomori	Chiba	Ehime	Fukui	Fukuoka	Fukushima	Gifu	Gumma	
1949	165,989	6,986	4,059	4,643	4,278	3,111	1,848	7,362	4,655	3,523	2,935	
1948	165,406	6,335	4,050	4,528	4,340	3,009	2,390	6,014	4,628	3,419	2,859	
1947	205,360	7,715	4,640	4,883	5,722	3,990	2,185	8,748	5,335	3,828	3,609	
Year	Hiroshima	Hokkaido	Hyogo	Ibaraki	Ishikawa	Iwate	Kagawa	Kagoshima	Kanagawa	Kochi	Kumamoto	Kyoto
1949	3,422	10,467	5,710	4,436	2,654	4,443	1,970	3,374	3,297	1,548	3,327	2,954
1948	3,520	10,352	5,689	4,341	3,018	4,323	2,251	3,610	3,265	1,717	3,294	2,866
1947	4,603	11,716	7,326	5,615	3,241	4,524	2,793	4,357	4,503	2,177	4,284	3,705
Mie	Miyagi	Miyazaki	Nagano	Narasaki	Nara	Niigata	Oita	Okayama	Osaka	Saga	Saitama	
1949	3,134	3,587	2,272	3,099	3,631	1,674	2,856	3,046	7,136	2,392	4,514	
1948	3,120	3,846	2,235	3,138	3,282	1,866	2,734	3,448	5,830	2,313	4,266	
1947	4,099	4,184	2,843	3,804	4,275	2,186	3,705	4,283	8,287	3,063	5,622	
Year	Shiga	Shimane	Shizuoka	Tochigi	Tokushima	Tokyo	Tottori	Toyama	Wakayama	Yamagata	Yamaguchi	Yamanashi
1949	1,903	1,970	4,549	2,931	1,878	7,959	1,159	3,203	1,734	3,615	2,897	1,367
1948	1,782	2,053	4,437	2,715	2,115	7,680	1,302	3,120	1,735	3,437	2,867	1,304
1947	2,247	2,471	5,365	3,797	2,702	9,813	1,507	3,956	2,180	4,042	3,531	1,662

Footnotes: * Data for 1949 are provisional.

Infant deaths are deaths under one year of Japanese nationals in Japan. Data for 1947 and 1948 are from Final Annual Report, Ministry of Welfare, and for 1949 from Monthly Schedule Reports, Ministry of Welfare.

This Table corresponds to Table 30. Statistical Annex, Public Health and Welfare in Japan, 1948.

TABLE 37. - INFANT DEATH RATES BY PREFECTURE: JAPAN, 1947-1949
(Rates per 1,000 live births each year)

Year	Annual	Aichi	Akita	Aomori	Chiba	Ehime	Fukui	Fukuoka	Fukushima	Gifu	Gumma
*1949	62.0	65.2	85.0	89.8	62.8	58.9	74.1	58.7	63.2	70.5	56.0
*1948	61.0	57.1	92.0	96.2	64.5	56.2	91.5	50.3	64.2	64.1	54.5
1947	76.7	73.9	97.0	99.7	77.4	75.6	85.9	80.8	74.9	74.4	66.1

Year	Hiroshima	Hokkaido	Hyogo	Ibaraki	Ishikawa	Iwate	Kanagawa	Kagoshima	Kochi	Kumamoto	Kyoto
*1949	54.3	63.0	56.9	65.9	82.3	89.3	63.3	52.0	43.6	58.2	52.4
*1948	56.4	66.9	55.0	64.2	87.3	91.4	66.3	56.6	44.3	62.5	53.8
1947	67.9	82.8	75.2	81.2	86.9	98.2	81.5	74.1	60.3	76.0	70.0

Year	Mie	Miyagi	Miyazaki	Nagano	Nagasaki	Nara	Niigata	Oita	Okayama	Osaka	Saga	Saitama
*1949	71.9	62.5	56.1	52.2	58.8	76.3	64.8	67.2	60.6	63.0	69.6	62.8
*1948	65.7	37.8	54.2	51.8	56.7	66.5	67.4	62.2	65.2	51.5	70.0	60.2
1947	88.7	75.4	73.6	61.4	80.6	91.1	72.4	87.5	80.0	79.9	95.8	72.8

Year	Shiga	Shimane	Shizuoka	Tochigi	Tokushima	Tokyo	Tottori	Toyama	Wakayama	Yamagata	Yamaguchi	Yamanashi
*1949	73.8	66.4	55.9	54.2	62.9	46.5	61.6	93.1	59.5	81.5	57.1	53.7
*1948	64.9	67.0	53.1	49.9	65.6	46.6	65.1	86.8	57.6	81.3	53.2	51.9
1947	86.9	76.0	65.8	69.2	85.4	62.4	77.9	95.5	70.8	92.0	71.7	63.2

Footnotes: * All data for 1949 are provisional, birth data for 1948 are provisional. Rates for 1949 and 1948 are provisional. Rates are number of deaths per 1,000 live births in the corresponding period and were computed by Public Health and Welfare Section, GHQ, SCAP. Data refer to events occurring to Japanese nationals in Japan. Data for 1947 and death data for 1948 are from Final Annual Reports, Ministry of Welfare, and birth data for 1948 and all data for 1949 from Monthly Schedule Reports, Ministry of Welfare.

This Table corresponds to Table 31, Statistical Annex, Public Health and Welfare in Japan, 1948.

TABLE 36. - 1/INFANT DEATHS FROM SELECTED CAUSES: JAPAN, 1947-1949

Year	All Causes	2/Typhoid fever (1)	Paratyphoid fever (2)	Cholera (4)	Cerebrospinal meningitis (6)	Anthrax (7)	4/Scarlet fever (8)	Whooping cough (9)	5/Diphtheria (10)	Erysipelas (11)
*1949	168,989	-	-	-	81	-	5	5,021	212	770
1948	165,406	3	-	-	87	-	5	2,568	341	1,060
1947	205,360	11	2	-	117	1	9	8,532	524	1,674

Year	Tetanus (12)	7/Tuberculosis (all forms) (13-22)	Leprosy (23)	Septicemia and purulent infection (nonpuerperal) (24)	Glanders (26a)	Dysentery (27)	Malaria (28)	8/Syphilis (30)	Influenza (33)
*1949	868	1,332	-	898	-	113	-	1,148	165
1948	1,000	1,197	-	1,029	-	100	3	1,142	135
1947	972	1,022	-	1,407	-	182	5	1,016	416

Year	Measles (35)	Japanese "B" encephalitis (37b)	Yellow fever (38a)	Rabies (38b)	Typhus fever & Typhus-like diseases (39)	Beriberi (68)	Meningitis due to meningococcus (81)	10/Convulsions (86)	Bronchitis (106)
*1949	4,509	8	-	-	2	3,514	2,543	1,409	9,574
1948	1,914	43	-	-	-	3,759	2,769	1,310	8,778
1947	5,094	8	-	2	-	4,843	3,360	1,444	11,548

Year	11/Pneumonia (all forms) (107-109)	Diarrhea, enteritis and ulceration of intestines (119)	Conenital malformations (157)	Conenital debility (158)	Premature birth (159)	Injury at birth (160)	Other diseases peculiar to first year of life (161)
*1949	27,724	26,815	5,255	37,285	13,667	1,082	8,967
1948	22,735	31,254	4,560	38,204	13,720	1,030	8,840
1947	35,262	35,679	3,929	41,751	14,597	885	9,868

TABLE 38. - 1/INFANT DEATHS FROM SELECTED CAUSES: JAPAN, 1947-1949, Cont'd

Year	Accidental deaths (169-195)	Ill-defined and unknown causes (199-200)	All other causes
*1949	1,931	2,587	11,507
1948	2,105	2,745	12,970
1947	1,846	3,103	15,251

Footnotes: * Data for 1949 are provisional.

1/ Data refer to deaths under one year of Japanese nationals in Japan. Data for 1947 and 1948 are from Final Annual Reports, Ministry of Welfare, and for 1949 from Monthly Schedule Reports, Ministry of Welfare.

2/ Excludes typhoid pneumonia.

3/ Not used.

4/ Excludes puerperal scarlet fever.

5/ Includes diphtheric endocarditis, excludes croupous bronchitis and laryngeal croup.

6/ Includes spondylitis, excludes pleuro pneumonia and nontuberculous adenopathy.

7/ Includes general paralysis of insane and aneurysm of the aorta.

8/ Includes croupous bronchitis.

9/ Includes typhoid pneumonia.

10/ "NA" indicates data not available.

A dash (-) indicates that no infant deaths were reported.

This Table corresponds to table 32, Statistical Annex, Public Health and Welfare in Japan, 1948.

TABLE 39. 1/INFANT DEATH RATES FROM SELECTED CAUSES: JAPAN, 1947-1949
(Rates per 1,000 live births each year)

Year	All Causes	2/Typhoid fever (1)	Paratyphoid fever (2)	Cholera (4)	Cerebrospinal (meningococcus) meningitis (6)	Anthrax (7)	4/Scarlet fever (8)	Whooping cough (9)	5/Diphtheria (10)	Erysipelas (11)	Tetanus (12)
*1949	62.0	-	-	-	0.0	-	0.0	1.8	0.1	0.3	0.3
*1948	61.0	0.0	-	-	0.0	-	0.0	0.9	0.1	0.4	0.4
*1947	76.7	0.0	0.0	-	0.0	0.0	0.0	3.2	0.2	0.6	0.4

Year	7/Tuberculosis (all forms) (13-22)	Septicemia and purulent infection (23)	Glanders (24)	Dysentery (25)	Malaria (26)	8/Syphilis (30)	Influenza (31)	Measles (32)	Japanese encephalitis (37b)
*1949	0.5	-	0.3	-	-	0.4	0.1	1.7	0.0
*1948	0.4	-	0.4	-	0.0	0.4	0.0	0.7	0.0
*1947	0.4	-	0.5	-	0.1	0.4	0.2	2.3	0.0

Year	Yellow fever (38a)	Rabies (38b)	Typhus fever & typhus-like diseases (39)	Beriberi (40)	Meningitis (not due to meningococcus) (41)	10/Convulsions (42)	Bronchitis (43)	11/Pneumonia (all forms) (107-109)	Diarrhea, enteritis and ulceration of intestines (119)
*1949	-	-	0.0	1.3	0.9	0.5	3.5	10.2	9.8
*1948	-	-	-	1.4	1.0	0.5	3.2	8.4	11.5
*1947	-	0.0	-	1.8	1.3	0.5	4.3	13.2	13.3

Year	12/Congenital malformations (157)	Congenital debility (158)	Premature birth (159)	Injury at birth (160)	Other diseases peculiar to first year of life (161)	Accidental deaths (169-195)	Ill-defined and unknown causes (199-200)	All other causes
*1949	1.9	13.7	5.0	0.4	3.3	0.7	0.9	4.2
*1948	1.7	14.1	5.1	0.4	3.3	0.8	1.0	4.8
*1947	1.5	15.6	5.4	0.3	3.7	0.7	1.2	5.7

TABLE 39. - 1/INFANT DEATH RATES FROM SELECTED CAUSES: JAPAN, 1947-1949, Cont'd

Footnotes:

* All data for 1949 are provisional, birth data for 1948 are provisional. Rates for 1948 and 1949 are provisional.
 1/ Rates are the number of infant deaths per 1,000 live births in the corresponding period and were computed by Public Health and Welfare Section, GHQ, SCAP. Data refer to deaths occurring to Japanese nationals in Japan. All data for 1947 and death data for 1948 are from Final Annual Report, Ministry of Welfare, birth data for 1948 and all data for 1949 are from Monthly Schedule Reports, Ministry of Welfare.

2/ Excludes typhoid pneumonia.

4/ Excludes puerperal scarlet fever.

5/ Includes diphtheric endocarditis, excludes croupous bronchitis and laryngeal croup.

7/ Includes spondylitis, excludes chronic pleuro pneumonia and nontuberculous adenopathy.

8/ Includes general paralysis of the insane and aneurysm of the aorta.

10/ Includes croupous bronchitis

11/ Includes typhoid pneumonia

A dash (-) indicates that there were no infant deaths reported.

A rate of 0.0 indicates that there were some infant deaths but the rate was less than 0.1.

This Table corresponds to Table 33, Statistical Annex, Public Health and Welfare in Japan, 1948.

TABLE 40. - 1/INFANT DEATHS FROM THE TEN LEADING CAUSES OF INFANT DEATHS IN 1949:
JAPAN, 1947-1949

List No.	Cause	*1949	1948	1947
158	Congenital debility	37,285	38,204	41,751
107-109	2/ Pneumonia (all forms)	27,724	22,735	35,262
119	Diarrhea, enteritis & ulceration of intestines	26,815	31,254	35,679
159	Premature birth	13,667	13,720	14,597
106	3/ Bronchitis	9,574	8,778	11,548
161	Other diseases peculiar to first year of life	8,967	8,840	9,868
157	Congenital malformations	5,255	4,560	3,929
9	Whooping cough	5,021	2,568	8,532
35	Measles	4,508	1,914	6,094
68	Beriberi	3,514	3,759	4,843

TABLE 40. - 1/INFANT DEATHS FROM THE TEN LEADING CAUSES OF INFANT DEATHS IN 1949:
JAPAN, 1947-1949, Cont'd

Footnotes:

* Data for 1949 are provisional.

1/ Data refer to deaths of Japanese nationals in Japan. Data for 1947 and 1948 are from Final Annual Reports, Ministry of Welfare, for 1949 from Monthly Schedule Reports, Ministry of Welfare.

2/ Includes typhoid pneumonia.

3/ Includes croupous bronchitis.

This Table corresponds to Table 34, Statistical Annex, Public Health and Welfare in Japan, 1948.

TABLE 41. - 1/INFANT DEATH RATES FOR THE TEN LEADING CAUSES OF INFANT DEATHS IN 1949:
JAPAN, 1947-1949
(Rates per 1,000 live births each year)

List No.	Cause	* 1949	* 1948	1947
158	Congenital debility	13.7	14.1	15.6
107-109	<u>2</u> / Pneumonia (All forms)	10.2	8.4	13.2
119	Diarrhea, enteritis and ulceration of intestines	9.8	11.5	13.3
159	Premature birth	5.0	5.1	5.4
106	<u>3</u> / Bronchitis	3.5	3.2	4.3
161	Other diseases peculiar to first year of life	3.3	3.3	3.7
157	Congenital malformations	1.9	1.7	1.5
9	Whooping cough	1.8	0.9	3.2
35	Measles	1.7	0.7	2.3
68	Beriberi	1.3	1.4	1.8

TABLE 41. - 1/ INFANT DEATH RATES FOR THE TEN LEADING CAUSES OF INFANT DEATHS IN 1949
JAPAN, 1947-1949, Cont'd

Footnotes:

* All data for 1949 and birth data for 1948 are provisional. Rates for 1948 and 1949 are provisional.

1/ Rates are the number of infant deaths per 1,000 live births in the corresponding period, and were computed by the Public Health and Welfare Section, CHQ, SCAP. Data refer to events occurring to Japanese nationals in Japan. All data for 1947 and death data for 1948 are from vital Annual Reports, Ministry of Welfare. Birth data for 1948 and all data for 1949 are from Monthly Schedule Reports, Ministry of Welfare.

2/ Includes typhoid pneumonia.
3/ Includes croupous bronchitis.

This Table corresponds to Table 35, Statistical Annex, Public Health and Welfare in Japan, 1948.

TABLE 42. - STILLBIRTHS BY MONTH: JAPAN, 1947-1949

Year	Total stillbirths	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Un- known
*1949	192,970	14,885	14,195	16,044	15,603	16,238	14,828	16,653	18,042	17,899	16,934	15,276	16,373	-
*1948	144,095	12,286	12,171	13,427	11,304	11,402	10,400	11,139	11,852	12,479	12,915	12,273	13,447	-
1947	123,837	10,071	9,772	10,299	9,440	9,883	9,218	10,619	10,730	10,642	10,942	10,585	11,664	2

Footnotes: * Data for 1948 and 1949 are provisional. Data refer to stillbirths after the third month occurring to Japanese nationals in Japan. Data for 1947 are from Final Annual Report, Ministry of Welfare, and for 1948 and 1949 from Monthly Schedule Reports, Ministry of Welfare.
This Table corresponds to Table 36, Statistical Annex, Public Health and Welfare in Japan, 1948.

TABLE 43. - STILLBIRTH RATES BY MONTH: JAPAN, 1947-1949
(Rates per 1,000 live births each month)

Year	Annual	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
*1949	70.8	45.6	58.1	64.3	70.6	79.7	78.1	78.6	82.2	80.7	77.0	72.3	79.9
*1948	53.2	39.0	46.7	48.3	50.4	56.6	55.4	54.2	55.1	58.0	59.2	55.9	70.6
1947	46.2	34.1	43.2	43.7	45.1	50.5	47.4	46.9	45.3	45.9	47.8	50.1	62.4

Footnotes: * Data for 1948 and 1949 are provisional.

Rates are the number of stillbirths after the third month per 1,000 live births during the corresponding period, and were computed by Public Health and Welfare Section, CHQ, SCAP. Data for 1947 are from Final Annual Report, Ministry of Welfare, and for 1948 and 1949 from Monthly Schedule Reports, Ministry of Welfare.
This Table corresponds to Table 37, Statistical Annex, Public Health and Welfare in Japan, 1948.

TABLE 44. - STILLBIRTHS BY PREFECTURE: JAPAN, 1947-1949

All											
Year	Japan	Aichi	Akita	Aomori	Chiba	Ehime	Fukui	Fukuoka	Fukushima	Gifu	Gumma
* 1949	192,970	8,301	3,339	2,893	4,149	3,357	1,642	9,288	5,310	3,462	4,218
* 1948	144,095	5,539	2,550	2,103	3,490	2,608	1,306	6,534	3,687	2,432	3,168
1947	123,837	4,533	2,239	1,801	3,165	2,271	1,078	5,011	3,376	2,016	2,799
Year	Hiroshima	Hokkaido	Hyogo	Ibaraki	Ishikawa	Iwate	Kagawa	Kagoshima	Kanagawa	Kochi	Kumamoto
* 1949	4,054	9,938	8,466	4,827	2,008	3,458	2,323	4,082	4,698	1,632	3,993
* 1948	3,299	6,931	6,011	3,942	1,486	2,627	1,954	3,073	3,834	1,289	3,206
1947	3,006	6,079	4,961	3,344	1,428	2,361	1,618	2,606	3,515	1,077	2,788
											2,429
Year	Mie	Miyagi	Miyazaki	Nagano	Nagasaki	Nara	Niigata	Oita	Okayama	Osaka	Saitama
* 1949	3,204	4,282	3,189	5,322	3,884	1,526	6,578	3,098	4,421	9,547	2,308
* 1948	2,324	3,288	2,087	3,878	2,902	1,274	4,569	2,332	3,424	6,565	1,518
1947	1,999	2,793	1,862	3,481	2,307	1,190	3,908	1,957	2,798	5,665	1,173
											3,713
Year	Shiga	Shimane	Shizuoka	Tochigi	Tokushima	Tokyo	Tottori	Toyama	Wakayama	Yamagata	Yamanashi
1949	1,897	2,424	5,550	3,244	2,187	11,061	1,782	2,096	2,020	3,123	3,838
1948	1,433	1,990	4,515	2,787	1,940	8,636	1,238	1,450	1,572	2,123	2,650
1947	1,107	1,656	3,841	2,273	1,691	7,656	1,131	1,290	1,244	1,914	2,228
											1,359

Footnotes: * Data for 1948 and 1949 are provisional.

Data refer to stillbirths after the third month occurring to Japanese nationals in Japan. Data for 1947 are from Final Annual Report, Ministry of Welfare, and for 1948 and 1949 from Monthly Schedule Reports, Ministry of Welfare.

This Table corresponds to Table 38, Statistical Annex, Public Health and Welfare in Japan, 1948.

TABLE 45. - STILLBIRTH RATES BY PREFECTURE: JAPAN, 1947-1949
(Rates per 1,000 live births each year)

Year	Annual	Aichi	Akita	Aomori	Chiba	Ehime	Fukui	Fukuoka	Fukushima	Gifu	Gumma
*1949	70.8	77.5	69.9	56.0	60.9	63.6	65.9	74.0	71.9	69.3	80.5
*1948	53.2	49.9	57.9	44.7	51.8	48.7	50.0	54.7	51.2	45.6	60.4
1947	46.2	43.4	46.8	36.8	42.8	43.0	42.4	46.3	47.4	39.2	51.3

Year	Hiroshima	Hokkaido	Hyogo	Ibaraki	Ishikawa	Iwate	Kagawa	Kagoshima	Kanagawa	Kochi	Kumamoto	Kyoto
*1949	64.3	59.8	84.4	71.7	62.3	69.5	74.7	62.9	62.2	61.4	62.3	81.5
*1948	52.9	44.8	58.1	58.3	43.1	55.5	57.5	48.2	52.0	46.9	52.3	53.4
1947	44.4	42.9	50.9	46.3	38.3	51.2	47.2	44.3	47.1	37.6	45.5	45.1

Year	Mie	Miyagi	Miyazaki	Nagano	Nagasaki	Nara	Nilgata	Oita	Okayama	Osaka	Saga	Saitama
*1949	73.5	74.6	78.7	89.7	62.9	69.5	77.9	72.9	88.0	84.3	67.1	64.6
*1948	49.0	58.0	50.6	64.0	50.1	54.1	55.4	57.6	64.7	58.0	46.0	55.3
1947	43.2	50.3	48.2	56.2	43.5	49.6	45.3	46.2	52.3	54.6	36.7	48.1

Year	Shiga	Shimane	Shizuoka	Tochigi	Tokushima	Tokyo	Tottori	Toyama	Wakayama	Yamagata	Yamaguchi	Yamanashi
*1949	73.5	81.7	68.2	60.0	73.3	64.6	94.8	60.9	69.3	70.4	75.6	78.6
*1948	52.2	65.0	54.0	51.2	60.2	52.4	61.9	40.3	52.2	50.2	55.0	59.3
1947	42.8	50.9	47.1	43.2	53.4	48.7	58.5	31.1	40.4	43.6	45.2	51.7

Footnotes: *Data for 1948 and 1949 are provisional.

Rates are the number of stillbirths after the third month per 1,000 live births during the corresponding period, and were computed by Public Health and Welfare Section, GHQ, SCAP. Data for 1947 are from Final Annual Report, Ministry of Welfare, and for 1948 and 1949 from Monthly Schedule Report, Ministry of Welfare. This Table corresponds to Table 39, Statistical Annex, Public Health and Welfare in Japan, 1948.

TABLE 46. - MARRIAGES BY MONTH: JAPAN, 1947-1949

Total													
Year	Marriages	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
* 1949	846,014	82,805	92,232	90,231	85,271	83,220	61,169	54,808	54,026	53,987	56,226	56,225	75,144
* 1948	960,275	70,909	94,987	106,775	102,887	95,167	71,542	66,821	67,704	64,634	64,933	69,730	86,126
1947	934,170	70,965	72,762	84,012	74,741	111,959	81,021	73,173	62,937	71,405	73,505	64,587	93,103

Footnotes: * Data for 1948 and 1949 are provisional.

Data refer to marriages of Japanese nationals in Japan. Data for 1947 are from Final Annual Report, Ministry of Welfare, and for 1948 and 1949 from Monthly Schedule Reports, Ministry of Welfare.

This Table corresponds to Table 40, Statistical Annex, Public Health and Welfare in Japan, 1948.

TABLE 47. - MARRIAGE RATES BY MONTH: JAPAN, 1947-1949

Year	Annual	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
* 1949	10.3	11.9	14.6	12.9	12.6	11.9	9.1	7.9	7.7	8.0	8.1	8.4	10.8
* 1948	12.0	10.4	14.9	15.7	15.7	14.0	10.9	10.1	9.4	9.8	9.6	10.6	12.7
1947	12.0	10.7	12.2	12.7	11.7	16.9	12.6	11.0	9.5	11.1	11.1	10.1	14.1

Footnotes: * Data for 1948 and 1949 are provisional. Rates are the number of marriages per 1,000 population and were computed by Public Health and Welfare Section, HE2, SCAP. Data for 1947 are from Final Annual Report, Ministry of Welfare, and for 1948 and 1949 from Monthly Schedule Report, Ministry of Welfare.

This Table corresponds to Table 41, Statistical Annex, Public Health and Welfare in Japan, 1948.

TABLE 48. - MARRIAGES BY PREFECTURE: JAPAN, 1947-1949

Year	All Japan	Aichi	Akita	Aomori	Chiba	Ehime	Fukui	Fukuoka	Fukushima	Gifu	Gumma
1949	846,014	31,438	14,572	14,376	20,246	16,370	8,408	37,671	23,657	16,028	16,382
1948	960,275	37,244	14,923	14,565	22,615	19,048	9,468	42,808	25,331	19,036	18,004
1947	934,170	35,018	16,439	14,487	24,266	17,719	10,667	39,288	26,598	19,101	18,847

Year	Hiroshima	Hokkaido	Hyogo	Ibaraki	Ishikawa	Iwate	Kanawa	Karoshima	Kanagawa	Kochi	Kumamoto	Kyoto
1949	22,804	44,755	32,209	21,021	9,857	14,866	10,661	19,392	21,423	9,385	19,834	16,353
1948	26,266	45,793	36,731	23,413	11,670	15,188	13,549	22,786	24,190	11,220	22,860	19,766
1947	25,818	41,114	32,971	22,249	12,797	14,708	12,632	20,708	23,493	10,986	21,688	18,294

Year	Mie	Miyagi	Miyazaki	Nagano	Nagasaki	Nara	Niigata	Oita	Okayama	Osaka	Saga	Saitama	Shiga
1949	14,820	17,553	11,967	21,297	17,973	8,444	26,344	13,418	18,028	35,815	11,190	20,955	8,643
1948	17,478	19,691	14,251	23,351	21,058	10,210	29,340	16,395	21,463	41,248	12,662	23,220	10,571
1947	16,258	20,088	13,153	23,842	21,054	9,916	34,187	16,306	19,546	34,328	12,133	24,739	9,594

Year	Shimane	Shizuoka	Tochigi	Tokushima	Tokyo	Tottori	Toyama	Wakayama	Yamagata	Yamaguchi	Yamanashi	uncelebrated
1949	9,822	23,781	16,435	9,650	49,237	6,727	10,342	10,595	16,997	16,306	7,962	-
1948	11,407	27,592	15,672	11,983	56,810	7,711	11,654	12,571	16,665	18,962	8,776	-
1947	12,606	29,168	18,610	11,528	49,800	7,579	13,638	11,576	17,599	18,563	8,527	544

Footnotes: * Data for 1948 and 1949 are provisional.

Footnotes: * Data for 1948 and 1949 are provisional.
 Data refer to marriages of Japanese nationals in Japan. Data for 1947 are from Final Annual Report, Ministry of Welfare, and for 1948 and 1949 from Monthly Schedule Reports, Ministry of Welfare.

This Table corresponds to Table 42, Statistical Annex, Public Health and Welfare in Japan, 1948.

TABLE 49. - MARRIAGE RATES BY PREFECTURE: JAPAN, 1947-1949
(Rates per 1,000 population)

Year	Annual	Aichi	Akita	Aomori	Chiba	Ehime	Fukui	Fukuoka	Fukushima	Gifu	Gumma
*1949	10.3	9.5	11.1	11.5	9.2	10.8	11.2	11.1	11.4	10.3	9.9
*1948	12.0	11.5	11.7	12.0	10.6	12.9	12.9	12.9	12.5	12.5	11.2
1947	12.0	11.2	13.1	12.3	11.5	12.2	13.9	12.4	13.4	12.8	12.0

Year	Hiroshima	Hokkaido	Iiyogo	Ibaraki	Ishikawa	Iwate	Kacawa	Kagoshima	Kanagawa	Kochi	Kumamoto	Kyoto
*1949	10.9	10.9	10.0	10.0	10.2	11.2	11.1	10.7	9.0	10.6	10.8	8.9
*1948	12.8	11.4	11.6	11.5	12.4	11.7	14.5	12.9	10.4	13.0	12.8	11.1
1947	12.9	10.7	10.8	11.1	13.8	11.7	13.8	11.9	10.6	13.0	12.3	10.5

Year	Mie	Miyagi	Miyazaki	Narano	Narasaki	Nara	Niigata	Oita	Okayama	Osaka	Saga	Saitama
*1949	10.0	10.7	11.1	10.0	11.2	10.6	10.6	10.5	10.7	9.9	11.7	9.6
*1948	12.0	12.3	13.5	11.2	13.5	13.1	12.0	13.2	13.0	11.7	13.6	10.9
1947	11.5	12.8	12.8	11.6	13.8	12.7	14.2	13.2	12.1	10.3	13.2	11.8

Year	Shiga	Shimane	Shizuoka	Tochigi	Tokushima	Tokyo	Tottori	Toyama	Wakayama	Yamagata	Yamaguchi	Yamanashi
*1949	9.7	10.6	9.6	10.3	10.8	8.9	11.1	10.1	10.6	12.3	10.6	9.5
*1948	12.1	12.6	11.5	12.0	13.8	10.5	13.0	11.7	12.8	12.4	12.6	10.8
1947	11.2	14.1	12.4	12.1	13.5	10.0	12.9	13.9	12.1	13.2	12.6	10.6

Footnotes: * Data for 1948 and 1949 are provisional.

Rates are the number of marriages per 1,000 population estimated as of 1 July of the year concerned, and were computed by Public Health and Welfare Section, WHO, SCAP. Data for 1947 are from Final Annual Report, Ministry of Welfare, and for 1948 and 1949 from Monthly Schedule Reports, Ministry of Welfare.

This Table corresponds to Table 47, Statistical Annex, Public Health and Welfare in Japan, 1948.

TABLE 50. - DIVORCES BY MONTH: JAPAN, 1947-1949

Year	Total Divorces	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
*1949	82,872	6,302	7,113	7,720	7,164	7,286	5,933	6,078	7,171	7,551	6,994	6,171	7,389
*1948	79,415	3,801	6,268	7,612	7,311	7,449	6,067	6,415	6,780	7,410	6,724	6,252	7,329
1947	79,551	6,415	5,997	7,049	6,541	7,818	5,979	6,158	6,634	7,230	8,732	5,612	7,386

Footnotes:

*Data for 1948 and 1949 are provisional. Data refer to divorces of Japanese nationals in Japan. Data for 1947 are from Final Annual Report, Ministry of Welfare, and for 1948 and 1949 from Monthly Schedule Reports, Ministry of Welfare.

This Table corresponds to Table 44, Statistical Annex, Public Health and Welfare in Japan, 1948.

TABLE 51. - DIVORCE RATES BY MONTH: JAPAN, 1947-1949
(Rates per 1,000 population per annum)

Year	Annual	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
*1949	1.0	0.9	1.1	1.1	1.1	1.0	0.9	0.9	1.0	1.1	1.0	0.9	1.1
*1948	1.0	0.6	1.0	1.1	1.1	1.1	0.9	0.9	1.0	1.1	1.0	1.0	1.1
1947	1.0	1.0	1.0	1.1	1.0	1.2	0.9	0.9	1.0	1.1	1.0	0.9	1.1

Footnotes:

*Data for 1948 and 1949 are provisional. Rates are the number of divorces per 1,000 population, and were computed by Public Health and Welfare Section, GHQ, SCAP. Data for 1947 are from Final Annual Report, Ministry of Welfare and for 1948 and 1949 from Monthly Schedule Report, Ministry of Welfare.

This Table corresponds to Table 45, Statistical Annex, Public Health and Welfare in Japan, 1948.

TABLE 52. - DIVORCES BY PREFECTURE: JAPAN, 1947-1949

Year	All Japan	Aichi	Akita	Aomori	Chiba	Ehime	Fukui	Fukuoka	Fukushima	Gifu	Gumma
* 1949	82,872	2,904	1,785	1,529	1,614	1,808	876	3,853	2,269	1,454	1,432
* 1948	76,415	2,436	2,028	1,403	1,598	1,780	891	3,653	2,214	1,317	1,375
1947	79,551	2,451	2,131	1,197	1,536	1,748	935	3,918	2,181	1,298	1,223

Year	Hiroshima	Hokkaido	Hyogo	Ibaraki	Ishikawa	Iwate	Karawa	Kagoshima	Kanagawa	Kochi	Kumamoto	Kyoto
* 1949	2,604	4,070	3,207	1,441	1,184	1,461	1,246	1,893	1,921	1,171	1,921	1,700
* 1948	2,503	3,603	2,977	1,329	1,224	1,590	1,090	1,771	1,874	1,175	1,906	1,629
1947	2,678	3,061	3,254	1,270	1,234	1,518	1,127	2,045	1,928	1,089	1,803	1,750

Year	Mie	Miyagi	Miyazaki	Nagano	Nagasaki	Nara	Niigata	Oita	Okayama	Osaka	Saga	Saitama
* 1949	1,340	1,452	1,099	1,598	2,042	819	2,799	1,323	1,860	3,850	1,030	1,552
* 1948	1,248	1,417	1,101	1,596	1,989	774	2,829	1,267	1,684	3,456	1,020	1,491
1947	1,247	1,273	1,111	1,558	1,916	822	2,660	1,333	1,639	3,833	1,031	1,523

Year	Shiga	Shimane	Shizuoka	Tochigi	Tokushima	Tokyo	Tottori	Toyama	Wakayama	Yamagata	Yamanashi	
* 1949	734	991	2,368	1,297	933	5,368	702	1,260	1,106	1,589	1,788	629
* 1948	729	1,019	2,339	1,261	867	4,733	705	1,392	1,091	1,642	1,694	605
1947	732	1,177	2,354	1,154	857	4,661	716	1,512	1,089	1,534	1,823	621

Footnotes: * Data for 1948 and 1949 are provisional. Data refer to divorces of Japanese nationals in Japan. Data for 1947 are from Final Annual Report, Ministry of Welfare, and for 1948 and 1949 from Monthly Schedule Reports, Ministry of Welfare.

This Table corresponds to Table 46, Statistical Annex, Public Health and Welfare in Japan, 1948.

TABLE 53. - DIVORCE RATES BY PREFECTURE: JAPAN, 1947-1949
(Rates per 1,000 population)

Year	Annual	Aichi	Akita	Aomori	Chiba	Ehime	Fukui	Fukuoka	Fukushima	Gifu	Gumma
*1949	1.0	0.9	1.4	1.2	0.7	1.2	1.2	1.1	1.1	0.9	0.9
*1948	1.0	0.8	1.6	1.2	0.7	1.2	1.2	1.1	1.1	0.9	0.9
1947	1.0	0.8	1.7	1.0	0.7	1.2	1.3	1.2	1.1	0.9	0.8

Year	Hiroshima	Hokkaido	Hyogo	Ibaraki	Ishikawa	Iwate	Karawa	Kagoshima	Kanagawa	Kochi	Kumamoto	Kyoto
*1949	1.2	1.0	1.0	0.7	1.2	1.1	1.3	1.0	0.8	1.3	1.0	0.9
*1948	1.2	0.9	0.9	0.7	1.3	1.2	1.2	1.0	0.8	1.4	1.1	0.9
1947	1.3	0.8	1.1	0.6	1.3	1.2	1.2	1.2	0.9	1.3	1.0	1.0

Year	Mie	Miyagi	Miyazaki	Nagano	Nagasaki	Nara	Milgata	Oita	Okayama	Osaka	Saga	Saitama
*1949	0.9	0.9	1.0	0.7	1.3	1.0	1.1	1.0	1.1	1.1	1.1	0.7
*1948	0.9	0.9	1.0	0.8	1.3	1.0	1.2	1.0	1.0	1.0	1.1	0.7
1947	0.9	0.8	1.1	0.8	1.3	1.1	1.2	1.1	1.0	1.2	1.1	0.7

Year	Shira	Shimane	Shizuoka	Tochigi	Tokushima	Tokyo	Tottori	Toyama	Wakayama	Yamagata	Yamaguchi	Yamanashi
*1949	0.8	1.1	1.0	0.8	1.0	1.0	1.2	1.2	1.1	1.2	1.2	0.8
*1948	0.8	1.1	1.0	0.8	1.0	0.9	1.2	1.4	1.1	1.2	1.1	0.7
1947	0.9	1.3	1.0	0.8	1.0	0.9	1.2	1.5	1.1	1.2	1.2	0.8

Footnotes: *Data for 1948 and 1949 are provisional. Rates are the number of divorces per 1,000 population estimated as of 1 July of the year concerned, and were computed by Public Health and Welfare Section, H2, SCAP. Data for 1947 are from Final Annual Report, Ministry of Welfare, and for 1948 and 1949 from Monthly Schedule Reports, Ministry of Welfare.

This Table corresponds to Table 47, Statistical Annex, Public Health and Welfare in Japan, 1948.

TABLE 54 - VENEREAL DISEASE CLINIC ACTIVITIES,
JAPAN, 1947 - 1949

Activity	1949	1948	1/ 1947
Average monthly case load			
All venereal diseases	79,681	59,510	29,537
Syphilis	52,128	37,794	18,107
Primary	10,459	7,816	3,150
Congenital	3,137	1,959	891
All other	38,532	28,019	14,066
Gonorrhea	23,216	18,333	9,923
Acute	5,328	12,207	6,320
Chronic	7,888	6,126	3,603
Chancroid	3,831	3,077	1,391
Other venereal diseases	506	306	116
2/ Total new cases not previously reported:			
All venereal diseases	176,642	212,889	25,669
Syphilis	85,716	96,819	10,131
Gonorrhea	80,740	101,563	13,211
Chancroid	9,287	13,492	2,216
Other venereal diseases	899	1,015	111
Total cases discontinuing treatment without permission:			
All venereal diseases	92,923	96,544	10,673
Syphilis	69,431	65,101	6,317
Gonorrhea	20,195	27,438	3,779
Chancroid	2,726	3,731	548
Other venereal diseases	571	274	29
Total contacts reported:			
All venereal diseases	97,815	99,848	13,205
Syphilis	67,473	62,584	8,733
Gonorrhea	28,105	26,611	3,885
Chancroid	1,967	8,337	555
Other venereal diseases	270	2,316	32
Total visits for contact tracing:			
All venereal diseases	74,289	75,013	10,635
Syphilis	51,852	44,271	7,187
Gonorrhea	20,772	20,252	2,943
Chancroid	1,447	8,221	482
Other venereal diseases	218	2,269	23

FOOTNOTES:

1/ Data available for November and December only.

2/ Venereal disease cases not previously treated by any clinic or physician.

SOURCE:

Monthly reports, Ministry of Welfare.

TABLE 55.- CONSUMPTION OF TOTAL AND RATIONED FOOD PER CAPITA PER DAY IN CALORIES,
Tokyo, Other Large Cities, and Rural Areas, 1947-1949

Kind of food and year	Tokyo			Other Large Cities 1/			Rural Areas 2/		
	Calories		%	Calories		%	Calories		%
	Total	Rationed food		Total	Rationed food		Total	Rationed food	
Total Food 3/									
1949	1982	1392	70.2	1983	1289	65.0	2140	394	18.4
1948	1924	1285	66.7	1959	1010	51.6	2059	457	22.2
1947	1852	1176	63.5	1858	1138	61.2	2130	210	9.9
Staple Foods 4/									
1949	1641	1206	73.3	1644	1225	74.5	1856	360	19.4
1948	1624	1148	70.7	1625	915	56.3	1798	448	24.9
1947	1601	1070	66.8	1573	1095	69.6	1888	195	10.3
Rice									
1949	956	832	87.0	905	735	81.2	1208	253	21.0
1948	860	704	81.8	967	594	61.4	1098	327	29.8
1947	654	472	72.3	817	602	73.7	1113	132	11.9
Wheat									
1949	362	272	75.0	373	288	77.4	155	46	29.7
1948	395	281	70.9	260	158	60.8	148	40	27.0
1947	417	338	80.9	304	241	79.3	150	22	14.4
Barley									
1949	134	84	62.8	209	120	57.4	248	42	17.1
1948	117	40	34.2	168	66	39.1	273	52	19.0
1947	84	50	59.4	99	55	55.8	265	19	7.0
Other Grains 5/									
1949	19	14	74.1	23	17	75.7	46	5	9.7
1948	19	10	54.6	21	14	66.7	34	4	11.5
1947	107	87	81.4	105	90	85.1	52	8	16.0
Sweet Potatoes									
1949	111	47	42.2	99	55	55.4	144	12	8.3
1948	157	76	49.9	152	74	48.7	166	20	12.1
1947	275	114	41.3	164	84	51.4	221	12	5.6

TABLE 55.-- CONSUMPTION OF TOTAL AND RATIONED FOOD PER CAPITA PER DAY IN CALORIES:
Tokyo, Other Large Cities, and Rural Areas, 1947-1949

Kind of food and year	Tokyo			Other Large Cities 1/			Rural Areas 2/		
	Total	Calories	%	Total	Calories	%	Total	Calories	%
		Rationed food	Rationed		Rationed food	Rationed		Rationed food	Rationed
Other Potatoes 5/									
1949	59	37	61.9	35	10	27.3	55	2	3.3
1948	76	35	46.2	57	9	15.5	79	5	6.4
1947	64	9	14.3	84	23	27.3	87	2	1.8
Legumes									
1949	83	42	50.5	67	24	35.5	89	11	12.3
1948	61	31	50.2	72	20	27.9	95	12	12.3
1947	97	77	79.4	60	19	32.0	93	7	7.3
Fish									
1949	108	31	28.8	113	11	9.5	75	8	10.8
1948	99	52	52.8	97	29	29.6	50	9	18.5
1947	79	21	26.8	94	11	11.8	42	4	10.1
Meat, Poultry, Eggs, Milk and Milk Products									
1949	32	4	13.4	34	3	8.2	11	7/	2.7
1948	22	7	31.0	24	3	14.0	8	7/	5.0
1947	10	1	11.7	15	2	15.6	6	1	11.7
Leafy Green and Yellow Vegetables									
1949	19	1	5.7	20	7/	1.5	23	7/	0.0
1948	30	11	36.0	26	5	17.6	30	7/	0.3
1947	20	1	3.4	35	2	6.6	33	7/	0.0
Other Fruits and Vegetables									
1949	28	1	2.2	40	1	1.2	36	7/	0.0
1948	36	8	22.2	46	6	13.0	36	7/	0.6
1947	27	2	5.5	42	2	4.1	35	7/	0.0

See footnotes at end of table.

TABLE 55.- CONSUMPTION OF TOTAL AND RATIONED FOOD PER CAPITA PER DAY IN CALORIES:
Tokyo, Other Large Cities, and Rural Areas, 1947-1949

FOOTNOTES:

- 1/ Includes Nagoya, Osaka, Kure, Fukuoka, Sendai, Sapporo, Kurezawa, and Matsuyama 1947-1949; includes Kobe, Yokohama and Kyoto 1949.
- 2/ Includes 27 prefectures in 1947; includes all 46 prefectures 1948-1949.
- 3/ Total food includes all edible foods consumed. Only certain specific foods are identified in this table.
- 4/ Staple foods includes rice, wheat, barley, other grains, sweet and white potatoes, yams and taro.
- 5/ Other grains includes all cereal grains except rice, wheat, and barley.
- 6/ Other potatoes includes white, yams, and taro.
- 7/ Less than one calorie per capita per day.
0.0 indicates less than 0.1 percent.

SOURCES:

Quarterly Nutritional Surveys, Ministry of Welfare.

TABLE 56. - FOOD CONSUMPTION IN GRAMS PER CAPITA PER DAY:
TOKYO, OTHER LARGE CITIES, AND RURAL AREAS, 1947-1949

Kind of Food	Tokyo		1/Other Large Cities		2/Rural Areas	
	1949	1948	1947	1948	1949	1947
<u>Grains</u>						
Rice	425.0	412.2	356.8	416.0	485.4	447.5
Wheat	276.7	248.4	187.4	277.3	354.1	323.3
Barley	114.1	128.5	118.4	84.0	46.5	44.1
Other grains 3/	38.8	32.8	23.9	60.3	71.4	73.8
	5.4	5.5	27.1	6.1	13.4	14.4
	0.5	0.8	0.4	0.6	0.8	0.4
<u>Nuts, etc.</u>						
Potatoes	158.8	221.5	227.3	113.4	180.5	223.6
Sweet	83.0	128.9	215.7	73.0	115.9	131.7
White	60.6	85.1	57.6	29.4	41.4	72.5
Other 4/	10.2	7.5	14.0	11.0	23.2	19.4
	8.6	6.8	1.0	7.1	4.2	3.5
<u>Sugars</u>						
	3.4	2.0	1.7	2.8	1.5	0.7
<u>Oils</u>						
Legumes	49.0	35.4	41.8	42.9	49.9	49.4
Soya	1.3	0.7	0.9	2.1	2.7	3.0
Soya products	43.0	28.8	17.3	37.0	41.3	39.1
Other beans	4.7	5.9	23.4	3.8	5.9	7.6
	91.7	81.0	62.3	98.9	56.9	38.1
<u>Animal Foods</u>						
Fish	67.9	67.0	55.3	73.2	31.4	26.0
Meat Poultry	13.3	9.3	4.7	14.8	3.1	1.9
Eggs	5.3	3.2	1.6	4.9	2.4	1.1
Milk	5.2	1.5	0.7	6.0	3.5	2.4
<u>Leafy, Green & Yellow Vegetables</u>	67.2	111.2	81.8	72.6	76.5	94.1
						95.8

TABLE 56.-FOOD CONSUMPTION IN GRAYS PER CAPITA PER DAY:
TOKYO, OTHER LARGE CITIES, AND RURAL AREAS, 1947-1949 - Cont'd

Kind of Food	Tokyo			1/Other Large Cities			2/Rural Areas		
	1949	1948	1947	1949	1948	1947	1949	1948	1947
Other Fruits & Vegetables	111.2	104.9	126.7	150.7	139.6	200.4	142.2	154.6	132.0
Citrus & Tomatoes	20.3	23.2	20.2	20.5	20.0	12.9	8.5	10.0	11.9
Other Fruits	17.9	18.4	5.9	21.6	21.7	12.1	13.0	12.9	9.0
Other Vegetables	71.7	66.3	100.6	106.6	157.9	175.4	126.7	131.7	141.1
Seaweeds	4.2	3.6	3.7	5.3	4.1	6.2	2.8	1.9	1.8
Processed Vegetables	31.6	28.8	24.8	39.5	50.6	43.0	55.2	64.4	70.1
Dried	1.5	1.4	0.8	1.9	3.1	3.6	3.1	3.0	3.2
Pickled	30.1	27.4	24.0	36.6	47.5	39.4	52.1	61.4	66.9
Flavors	28.7	23.3	17.2	32.2	26.6	26.4	28.1	19.1	21.2

Footnotes:

1/ Includes Nagoya, Osaka, Kure, Fukuoka, Sendai, Sapporo, Kurazawa, Matsuyama, Kobe, Yokohama, and Kyoto.

2/ Includes all rural areas in 46 prefectures.

3/ Other grains includes all cereal grains except rice, wheat and barley.

4/ Other potatoes includes yams and taro.

Sources: Quarterly Nutrition Surveys, Ministry of Welfare.

TABLE 57.- NUTRITION VALUE OF FOOD CONSUMED DAILY.
Tokyo, Other Large Cities, and Rural Areas, 1947 - 49

	Tokyo		Other Large Cities 1/				Rural Areas 2/	
	1949	1948	1947	1949	1948	1947	1949	1948
Calories	1982	1924	1852	1983	1959	1858	2140	2059
Total protein (grams)	66	63	61	69	66	62	65	61
Animal protein (grams)	18	16	13	20	18	16	12	8
Fat (grams)	19	17	12	18	15	13	15	13
Carbohydrates (grams)	381	381	364	388	393	364	437	426
Calcium (grams)	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
Phosphorus (grams)	1.7	1.7	1.7	1.7	1.8	1.7	1.7	1.7
Iron (grams)	45	46	52	47	49	41	46	48
Vitamins:								
A (Int. Units)	2267	3498	2485	2452	2933	3295	2385	2866
B1 (mgm)	1.5	1.6	1.7	1.6	1.6	1.6	1.6	1.6
B2 (mgm)	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7
Niacin (mgm)	9.5	10.0	9.7	9.7	10.2	9.9	9.6	9.1
C (mgm)	99	142	149	102	141	152	117	138
								155

Footnotes:

1/ Includes Nagoya, Osaka, Kure, Fukuoka, Sendai, Sapporo, Kurezawa, and Matsuyama 1947-1949; includes Kobe, Yokohama and Kyoto 1949.

2/ Includes 27 prefectures in 1947; includes all 46 prefectures 1948-1949

Sources: Quarterly Nutritional Surveys, Ministry of Welfare.

TABLE 58. - AVERAGE BODY HEIGHT AND WEIGHT*
Japan, 1947-1949

Age in Years	MALE				FEMALE			
	Height in Cms.		Weight in Kms.		Height in Cms.		Weight in Kms.	
	Normal	1947	1948	1949	Normal	1947	1948	1949
0	-	64.4	65.4	64.9	-	63.3	63.8	64.2
1	75.9	75.4	74.5	74.4	74.7	73.9	73.4	73.6
2	84.5	83.5	82.4	83.2	83.3	82.3	81.9	82.0
3	91.3	90.2	89.8	90.3	90.2	88.7	89.3	89.2
4	97.5	96.2	96.0	96.4	96.3	95.0	95.2	95.5
5	103.1	101.7	101.8	102.7	102.2	101.4	100.5	101.4
6	108.1	107.1	107.3	108.2	107.2	106.0	105.6	107.1
7	111.0	111.8	112.3	113.2	110.8	110.8	110.7	112.5
8	116.0	116.2	116.9	117.5	115.0	115.7	115.8	117.2
9	121.0	121.5	121.6	122.0	119.0	120.5	121.0	121.3
10	125.0	125.5	125.9	126.0	124.0	125.5	126.0	126.8
11	129.0	129.7	130.3	130.2	129.0	129.9	130.9	130.8
12	134.0	133.6	135.0	134.5	136.0	134.8	136.0	136.1
13	140.0	138.6	139.7	139.8	141.0	139.6	140.7	141.1
14	148.0	143.8	144.9	145.3	145.0	144.0	144.5	145.0
15	153.0	150.3	150.2	151.3	148.0	147.9	147.3	147.7
16	157.0	155.6	154.6	155.7	149.0	149.3	149.2	149.1
17	159.0	158.9	158.2	158.4	149.0	150.2	150.1	150.2
18	161.0	160.1	160.3	160.0	149.0	150.3	150.7	150.7
19	161.0	160.9	160.9	160.8	149.0	150.6	151.1	151.0
20	161.0	161.5	161.2	161.2	150.0	150.4	150.9	151.1
21	161.0	161.0	161.5	161.6	150.0	150.2	150.6	151.0
22	161.0	161.3	161.6	161.5	150.0	150.3	150.5	150.9
23	162.0	161.3	161.6	161.6	149.0	149.4	150.4	150.5
24	161.0	161.0	161.6	161.8	149.0	150.1	150.6	150.3
25	161.0	161.3	161.6	161.8	150.0	150.0	150.5	150.1
26-30	160.3	160.7	161.2	161.2	149.4	149.8	149.5	149.9
31-40	160.1	160.0	160.3	160.3	148.7	148.8	148.6	149.1
41-50	158.7	158.8	159.1	159.1	147.0	147.4	147.3	147.7
51-60	157.9	157.4	157.5	157.5	145.5	145.6	145.5	145.2
61-70	155.6	155.8	155.7	155.8	143.3	143.2	143.3	143.2
71 & over	154.2	154.2	153.6	153.6	140.5	140.4	140.6	140.9
				48.5	48.6	48.8	49.1	41.7
				6.6	7.1	6.8	8.8	8.7
				9.0	9.0	8.7	11.0	10.6
				11.2	10.8	12.6	12.8	12.4
				12.6	14.1	15.7	15.8	15.8
				14.1	16.9	17.6	18.6	19.4
				16.9	19.0	20.8	22.4	23.3
				20.8	22.9	25.0	27.5	28.4
				25.0	27.5	30.9	31.5	32.1
				31.5	35.1	35.5	35.9	36.3
				35.1	39.4	43.6	40.4	40.5
				43.6	47.1	46.5	45.9	47.1
				47.1	49.0	49.1	48.4	50.3
				49.1	50.5	51.1	48.6	50.8
				50.5	51.2	51.0	48.7	50.8
				50.8	50.7	50.8	48.5	50.3
				50.3	50.3	50.3	47.9	50.1
				50.1	50.0	50.2	47.8	50.2
				50.2	49.9	49.9	49.9	49.9
				49.9	49.2	49.1	49.1	48.5
				48.5	47.7	47.4	47.7	48.5
				46.4	43.2	46.4	43.2	44.5
				44.5	40.9	41.1	40.9	41.7

SOURCE: Nutrition Survey for month of May of year cited, Ministry of Welfare.
*Height is stated in centimeters and weight in kilograms.

TABLE 59. - SCHOOL LUNCH PROGRAM: JAPAN, OCT. 1947 - DEC. 1949

Month and Year	Sponsored by National Government			Sponsored by Local Government		
	Urban		No. of schools	Rural		No. of schools
	No. of schools	No. of children		No. of schools	No. of children	
Total	No. of schools	No. of children	No. of schools	No. of schools	No. of children	No. of children
1947						
October	4391	2308396	3549	2832364	842	76032
November	4840	4092879	3558	3023167	1282	1069712
December	5467	4137975	3571	3054586	1896	1083399
1948						
January	6693	4773277	3579	3074586	1965	1420757
February	6612	4804302	3576	3077265	1977	1422669
March	6961	4834589	3579	3079148	1981	1402804
April	6858	5209554	3584	3104329	1984	1496967
May	6958	5261192	3593	3188204	1987	1493001
June	7490	5385700	3598	3205674	2011	1490950
July	7826	5517983	3621	3177912	2018	1408686
August	(120011)		*	*	*	*
September	8606	6141040	3617	3286526	2826	1546609
October	8647	6108032	3622	3288438	2884	1550909
November	8672	6127258	3620	3289714	2899	1552647
December	9036	6292311	3627	3287322	2874	1540292
1949						
January	9099	6277754	3629	3300575	2911	1449425
February	9350	6465842	3697	3415147	2843	1334863
March	9758	6710501	3680	3734812	2760	1415488
April	9859	6473737	3606	3449783	2934	1300217
May	10541	6562229	3809	3606591	2731	1143409
June	10654	6982227	3704	3559444	2836	1190556
July	10411	7026670	3733	3586944	2807	1163056
August	(125401)		*	*	*	*
September	10583	7053694	3751	3596962	2769	1153038
October	10578	7086407	3712	3555570	4498	2544430
November	10594	7173224	3769	3611490	4441	2498502
December	10882	7178557	3739	3552192	4471	2547808
Footnotes:	* Data for summer school program during month of August not available except for total number of children.					

Source: Special Report, Ministry of Education.

TABLE 60. - PUBLIC ASSISTANCE PROGRAM BY MONTH: JAPAN, 1946-1949

	Total	Persons Assisted ^{1/}		Total	Cost ^{2/}		Kind
		Institutional	Non-Institutional		Cash		
1946							
August	2953298	66991	2886307	101785455	91030771		10754684
September	2856911	65821	2797390	127786099	118879647		8915362
October	2707439	67047	2636392	137063874	120478159		16486715
November	2658769	80885	2577884	130922273	115235943		15686330
December	2837207	92778	2744429	206121565	144075959		62045606
1947							
January	2891336	100545	2790791	251604569	160259872		91344697
February	2701670	98837	2602833	177036655	156671451		20365804
March	2882023	108420	2773603	252964185	201012453		51961732
April	2767041	112769	2650272	234480696	197818652		36642044
May	2763335	126052	2637281	235030843	208811628		26219215
June	2747512	131628	2615884	234333722	213863215		20470507
July	2656478	130575	2525963	259804242	274359213		25445029
August	2688891	146885	2542006	315187198	293439251		22127937
September	3210865	223742	2987123	409654297	339282546		70371751
October	2947295	147734	2799561	380556040	358731288		21824752
November	2721316	138596	2582720	516877433	366094622		50782811
December	2841273	128623	2702650	439342881	399638024		39704857
1948							
January	2199594	137796	2061798	407902232	393573119		14329113
February	2156094	136684	2020410	428190769	376219148		51971621
March	2215226	138195	2077031	464416236	408285028		56131208
April	2149871	135944	2013927	425886272	366200622		59685650
May	2048564	136898	1911666	412721121	377359593		35361528
June	1982020	140591	1841429	427371772	397062940		30308832

TABLE 60. - PUBLIC ASSISTANCE PROGRAM BY MONTH, JAPAN 1946-1949 Cont'd

	Total	Persons Assisted ^{1/}		Cost ^{2/}		Kind
		Institutional	Non-Institutional	Total	Cash	
1948 (Cont'd)						
July	1955569	142198	1813371	509782304	399900645	109881659
August	1903539	144613	1758926	479065048	420431374	58633714
September	1917990	144882	1773108	542504234	519237684	27215546
October	1863912	143806	1720106	590155490	565029472	25126018
November	1834516	143940	1690576	598595126	579688317	18906809
December	1844225	140284	1703841	647591778	627365281	20226497
1949						
January	1765110	139160	1625950	645977507	631149911	14833592
February	1715732	138575	1577157	675709384	650124954	25673430
March	1721991	139312	1582679	853997166	801720789	52276377
April	1651233	133412	1517821	713351879	660104765	53247114
May	1650109	130847	1519262	700215745	661773531	38442214
June	1661223	132310	1528913	762449085	767113119	15735966
July	1689174	133865	1534509	861415878	848419481	12996497
August	1681166	132721	1548845	392149052	878768129	13350923
September	1990547	132583	1857864	953171843	892053116	64118727
October	1699184	135551	1563533	910772293	897392836	13379457
November	1723369	131385	1591984	929627627	916409402	13218225
December	1725728	130609	1595119	1034769935	991028790	43740245

Footnotes: ^{1/} The number of persons assisted refers to the number of times assistance was given to welfare cases during the month indicated. This number exceeds the actual number of individuals on welfare rolls by an unknown amount.

^{2/} Increase of per capita cost of public assistance is attributable to rising cost of items on welfare budget because of devaluation of the yen and increase in number of items provided.

SOURCE: Monthly Reports, Ministry of Welfare.

TABLE 61. - LIVE BIRTHS, DEATHS, INFANT DEATHS, STILLBIRTHS,
MARRIAGES AND DIVORCES BY PREFECTURE: JAPAN, 1949

Area	Live Births	Deaths	*Infant Deaths	Stillbirths	Marriages	Divorces
All Japan	2,723,873	950,719	168,989	192,970	946,014	82,872
All "Shi"	928,949	311,683	51,164	88,862	278,456	31,580
All "Gun"	1,794,924	639,036	117,825	104,108	567,558	51,292
Aichi	107,117	36,463	6,986	8,301	31,438	2,904
Akita	47,736	17,233	4,059	3,339	14,572	1,785
Aomori	51,703	16,842	4,643	2,893	14,376	1,589
Chiba	68,116	26,037	4,278	4,149	20,246	1,614
Ehime	52,797	17,645	3,111	3,357	16,370	1,808
Fukui	24,925	9,954	1,848	1,642	8,408	876
Fukuoka	125,514	40,041	7,362	9,288	37,671	3,853
Fukushima	73,867	23,841	4,665	5,310	23,657	2,269
Gifu	49,989	18,536	3,523	3,462	16,028	1,454
Gumma	52,417	18,452	2,935	4,218	16,382	1,432
Hiroshima	63,039	23,512	3,422	4,054	22,804	2,604
Hokkaido	166,215	48,361	10,467	9,938	44,765	4,070
Hyogo	100,354	34,695	5,710	8,466	32,209	3,207
Ibaraki	67,326	24,854	4,436	4,827	21,021	1,441
Ishikawa	32,250	13,015	2,654	2,008	9,857	1,184
Iwate	49,740	18,427	4,443	3,458	14,866	1,461
Kagawa	31,107	11,247	1,970	2,323	10,661	1,246
Kagoshima	64,875	21,455	3,374	4,082	19,392	1,893
Kanagawa	75,555	22,985	3,297	4,698	21,423	1,921
Kochi	26,587	10,577	1,548	1,632	9,385	1,171
Kumamoto	63,551	21,401	3,327	3,393	19,834	1,921
Kyoto	52,930	19,707	2,954	4,312	16,353	1,700
Mie	43,610	17,526	3,134	3,204	14,820	1,340
Niyagi	57,430	18,184	3,587	4,282	17,553	1,452
Miyazaki	40,508	12,913	2,272	3,189	11,967	1,099
Nagano	59,316	22,818	3,099	5,322	21,297	1,598
Nagasaki	61,711	19,709	3,631	3,884	17,978	2,042
Nara	21,946	9,584	1,674	1,526	8,444	819
Niigata	84,471	31,550	5,471	6,578	26,344	2,799
Oita	42,511	17,001	2,856	3,098	13,418	1,323
Okayama	50,235	19,557	3,046	4,421	18,028	1,860
Osaka	113,241	38,883	7,136	9,547	35,815	3,850
Saga	34,391	12,306	2,392	2,308	11,190	1,030
Saitama	71,855	25,698	4,514	4,639	20,955	1,552
Shiga	25,794	11,217	1,903	1,897	8,643	734
Shimane	29,678	12,033	1,970	2,424	9,822	991
Shizuoka	81,390	25,380	4,549	5,550	23,781	2,368
Tochigi	54,054	17,855	2,931	3,244	16,435	1,297
Tokushima	29,838	11,081	1,878	2,187	9,650	933
Tokyo	171,309	54,240	7,959	11,061	49,237	5,368
Tottori	18,801	7,356	1,159	1,782	6,727	702
Toyama	34,392	14,287	3,203	2,096	10,342	1,260
Wakayama	29,139	11,183	1,734	2,020	10,595	1,106
Yamagata	44,340	17,656	3,615	3,123	16,997	1,589
Yamaguchi	50,760	18,508	2,897	3,838	16,306	1,788
Yamanashi	25,443	8,914	1,367	2,000	7,962	629

See footnotes on next page

TABLE 61. - LIVE BIRTHS, DEATHS, INFANT DEATHS, STILLBIRTHS,
MARRIAGES AND DIVORCES BY PREFECTURE: JAPAN, 1949 Cont'd

FOOTNOTES:-

*Deaths under one year of age.

Data refer to vital events of Japanese nationals in Japan.

SOURCE:-

Monthly Vital Statistics Schedule Reports, Ministry of Welfare.

TABLE 62. - 1/ LIVE BIRTH, DEATH, INFANT DEATH, STILLBIRTH,
MARRIAGE AND DIVORCE RATES BY PREFECTURE: JAPAN, 1949

Area	Live Birth Rate	Death Rate	Infant Death Rate	Stillbirth Rate	Marriage Rate	Divorce Rate
All Japan	33.1	11.6	62.0	70.8	10.3	1.0
All "Shi"	32.7	11.0	55.1	95.7	9.8	1.1
All "Gun"	33.4	11.9	65.6	58.0	10.5	1.0
Aichi	32.4	11.0	65.2	77.5	9.5	0.9
Akita	36.3	13.1	85.0	69.9	11.1	1.4
Aomori	41.4	13.5	89.8	56.0	11.5	1.2
Chiba	31.1	11.9	62.8	60.9	9.2	0.7
Ehime	34.8	11.6	58.9	63.6	10.8	1.2
Fukui	33.2	13.2	74.1	65.9	11.2	1.2
Fukuoka	37.0	11.8	58.7	74.0	11.1	1.1
Fukushima	35.6	11.5	63.2	71.9	11.4	1.1
Gifu	32.0	11.9	70.5	69.3	10.3	0.9
Gunma	31.8	11.2	56.0	80.5	9.9	0.9
Hiroshima	30.1	11.2	54.3	64.3	10.9	1.2
Hokkaido	40.3	11.7	63.0	59.8	10.9	1.0
Hyogo	31.0	10.7	56.9	84.4	10.0	1.0
Ibaraki	32.1	11.9	65.9	71.7	10.0	0.7
Ishikawa	33.4	13.5	82.3	62.3	10.2	1.2
Iwate	37.5	13.9	89.3	69.5	11.2	1.1
Kagawa	32.5	11.7	63.3	74.7	11.1	1.3
Kagoshima	35.8	11.9	52.0	62.9	10.7	1.0
Kanagawa	31.8	9.7	43.6	62.2	9.6	0.8
Kochi	29.9	11.9	58.2	61.4	10.6	1.3
Kumamoto	34.7	11.7	52.4	62.8	10.8	1.0
Kyoto	28.9	10.8	55.8	81.5	8.9	0.8
Mie	29.3	11.8	71.9	73.5	10.0	0.9
Miyagi	35.1	11.1	62.5	74.6	10.7	0.9
Miyazaki	37.6	12.3	58.8	62.9	11.2	1.3
Nagano	27.8	10.7	52.2	89.7	10.0	0.7
Nagasaki	38.5	12.3	58.8	62.9	11.2	1.3
Nara	27.5	12.0	76.3	69.5	10.6	1.0
Niigata	33.8	12.6	64.8	77.9	10.6	1.1
Oita	33.3	13.3	67.2	72.9	10.5	1.0
Okayama	29.7	11.6	60.6	88.0	10.7	1.1
Osaka	31.4	10.8	63.0	84.3	9.9	1.1
Saga	36.0	12.9	69.6	67.1	11.7	1.1
Saitama	32.9	11.8	62.8	64.6	9.6	0.7
Shiga	28.8	12.5	73.8	73.5	9.7	0.8
Shimane	32.1	13.0	66.4	81.7	10.6	1.1
Shizuoka	33.0	10.3	55.9	68.2	9.6	1.0
Tochigi	33.9	11.2	54.2	60.0	10.3	0.8
Tokushima	33.5	12.4	62.9	73.3	10.8	1.0
Tokyo	30.9	9.8	46.5	64.6	8.9	1.0
Tottori	30.9	12.1	61.6	94.8	11.1	1.2
Toyama	33.6	14.0	93.1	60.9	10.1	1.2
Wakayama	29.0	11.1	59.5	69.3	10.6	1.1
Yamagata	32.1	12.8	81.5	70.4	12.3	1.2
Yamaguchi	32.9	12.0	57.1	75.6	10.6	1.2
Yamanashi	30.4	10.7	53.7	78.6	9.5	0.8

See footnotes on next page

TABLE 62. - 1/ LIVE BIRTH, DEATH, INFANT DEATH, STILLBIRTH
MARRIAGE AND DIVORCE RATES BY PREFECTURE: JAPAN, 1949 Cont'd

FOOTNOTES:-

- 1/ Birth, death, marriage and divorce rates are per 1,000 population, estimated as of 1 July 1949 and the infant death and stillbirth rates are per 1,000 live births during 1949.

Data refer to vital events of Japanese nationals in Japan.

SOURCES:-

Rates were computed by Public Health and Welfare Section, GHQ, SCAP.
Sources of original data were Monthly Vital Statistics Schedule Reports, Ministry of Welfare.

TABLE 63 - LIVE BIRTHS BY MONTH BY PREFECTURE: JAPAN, 1949

Area	Total Live Births	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
All Japan	2,723,873	326,166	244,596	249,484	220,891	203,738	189,937	211,921	219,500	221,808	219,877	211,220	204,825
All "Shi"	928,949	109,858	81,195	82,192	72,725	68,357	67,602	76,387	77,122	76,243	73,520	72,297	71,951
All "Gun"	1,794,924	216,308	163,311	167,292	148,166	135,381	122,335	135,534	142,378	145,565	146,357	138,923	132,874
Aichi	107,117	13,572	9,544	9,162	9,091	7,944	7,295	8,498	8,807	8,425	8,265	8,383	8,131
Akita	47,736	5,365	4,598	4,567	4,126	3,339	2,861	3,401	3,833	4,277	4,273	3,835	3,261
Amori	51,703	5,318	4,273	4,628	4,535	4,198	3,688	3,907	4,252	4,577	4,866	4,032	3,429
Osaka	66,116	8,191	5,894	6,337	5,543	5,092	4,514	4,974	5,494	5,714	5,600	5,292	5,471
Osaka	52,797	6,474	4,783	4,642	4,102	3,805	3,570	4,230	4,265	4,394	4,342	4,170	4,020
Fukuoka	24,925	3,217	2,426	2,345	2,010	1,913	1,855	1,950	1,964	2,027	1,984	1,693	1,541
Fukuoka	12,514	1,908	1,233	1,314	9,533	8,784	8,744	10,164	10,352	10,567	9,892	9,504	9,519
Fukuoka	73,867	7,973	6,292	6,735	6,121	5,778	4,987	5,248	6,010	6,089	6,704	6,288	5,642
Gifu	49,989	6,480	4,619	4,615	4,244	3,780	3,476	3,844	3,926	3,717	3,860	3,789	3,639
Gumma	6,355	4,776	4,776	4,767	4,236	4,185	3,775	4,039	4,129	4,165	4,002	4,019	3,969
Hiroshima	63,039	8,055	5,665	5,554	5,074	4,706	4,595	5,452	5,224	5,025	4,816	4,474	4,399
Hokkaido	166,215	19,202	15,005	16,385	13,725	13,124	12,296	12,459	13,054	13,833	13,577	12,188	11,367
Hyogo	100,354	12,150	9,359	9,633	7,845	7,664	7,600	8,497	8,402	7,770	7,148	7,103	7,183
Ibaraki	67,326	7,546	5,874	6,278	5,787	5,363	4,661	5,022	5,295	5,231	5,371	5,184	5,714
Ishikawa	32,250	4,151	3,262	3,367	2,801	2,439	2,102	2,317	2,446	2,539	2,435	2,286	2,105
Iwate	49,740	5,536	4,672	4,912	4,817	4,014	3,353	3,568	3,687	3,677	4,143	3,899	3,522
Kagawa	31,107	4,294	2,903	2,910	2,344	2,157	2,152	2,570	2,616	2,464	2,348	2,174	2,175
Kagoshima	64,875	7,775	5,379	5,359	5,363	4,544	4,299	4,684	5,156	5,394	5,704	5,576	5,682
Kanagawa	75,555	8,904	6,376	6,688	5,652	5,819	5,773	6,028	5,950	6,234	6,028	5,876	6,227
Kochi	28,587	3,162	2,328	2,292	1,965	1,918	1,926	2,106	2,212	2,167	2,204	2,154	2,133

TABLE 63 - LIVE BIRTHS BY MONTH BY PREFECTURE: JAPAN, 1949 Cont'd

Area	Total Live Births	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Kumamoto	63,551	7,698	5,619	5,530	4,711	4,547	4,218	4,840	5,140	5,293	5,266	5,568	5,091
Kyoto	52,930	6,458	4,961	5,157	4,431	3,975	3,822	4,251	4,201	3,975	3,820	3,909	3,988
Mie	43,610	5,410	3,931	3,835	3,392	3,170	2,996	3,258	3,720	3,421	3,693	3,572	3,282
Miyagi	57,430	6,594	5,869	5,609	5,035	4,152	3,594	4,119	4,260	4,570	5,057	4,832	4,339
Miyazaki	40,508	3,479	3,467	3,481	3,173	3,159	2,768	3,058	3,357	3,290	3,248	3,328	3,471
Nagano	59,316	7,324	5,581	5,542	4,969	4,869	4,116	4,247	4,337	4,844	4,862	4,490	4,135
Nagasaki	61,711	6,893	5,522	5,349	4,797	4,371	4,057	5,067	5,021	5,235	5,094	5,257	5,048
Nara	21,946	2,766	1,952	1,991	1,727	1,619	1,554	1,762	1,774	1,701	1,646	1,764	1,690
Niigata	84,471	9,107	7,680	8,628	7,038	6,397	5,361	6,121	6,909	7,679	7,501	6,577	5,473
Oita	42,511	5,455	4,097	3,851	3,591	3,085	2,849	3,300	3,433	3,372	3,285	3,167	3,022
Okayama	50,235	6,616	4,840	4,546	4,066	3,600	3,406	4,166	4,135	3,862	3,757	3,625	3,616
Osaka	113,241	13,009	10,245	10,218	8,441	8,500	8,867	10,165	9,621	8,794	8,533	8,478	8,370
Saga	34,391	4,442	3,154	3,091	2,573	2,456	2,111	2,556	2,707	2,852	2,816	2,826	2,807
Saitama	71,855	8,452	6,315	6,564	5,606	5,148	4,992	5,639	5,868	6,053	5,780	5,713	5,725
Shiga	25,794	3,051	2,472	2,602	2,148	2,091	1,831	1,941	2,088	1,946	1,913	1,822	1,889
Shimane	29,678	3,558	2,758	2,927	2,535	2,188	2,019	2,274	2,284	2,345	2,503	2,228	2,059
Shizuoka	81,390	9,985	7,167	6,671	6,832	5,873	5,648	6,173	6,355	6,534	6,771	6,589	6,792
Tochigi	54,054	6,238	4,784	5,112	4,455	4,206	3,762	4,038	4,425	4,293	4,359	4,132	4,250
Tokushima	27,838	3,838	2,793	2,785	2,498	2,157	2,013	2,214	2,325	2,315	2,369	2,306	2,404
Tokyo	171,509	20,695	14,374	14,859	13,210	12,453	12,713	13,906	14,012	14,277	13,689	13,914	13,807
Tottori	18,801	2,526	1,704	1,582	1,664	1,401	1,245	1,425	1,384	1,437	1,641	1,388	1,404
Toysama	34,392	4,110	3,190	3,484	3,047	2,699	2,264	2,535	2,806	2,914	2,706	2,450	2,187
Wakayama	29,139	3,590	2,640	2,686	2,256	2,068	1,932	2,108	2,378	2,383	2,247	2,290	2,261
Yamagata	54,125	3,797	4,158	3,634	3,331	3,005	3,331	3,588	3,933	3,875	3,996	3,511	3,052
Yamaguchi	50,760	6,211	4,575	4,035	4,035	3,575	3,473	4,306	4,433	4,248	3,996	3,735	3,677
Yamaguchi	25,443	3,288	2,318	2,250	2,293	2,082	1,839	1,863	1,865	1,974	1,924	1,890	1,857

Data refer to births of Japanese nationals in Japan.

Sources: Monthly Vital Statistics Schedule Reports, Ministry of Welfare

TABLE 64. - LIVE BIRTH RATES BY MONTH BY PREFECTURE: JAPAN, 1949
(Rates per 1,000 population per annum)

Area	Annual	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
All Japan	33.1	46.7	38.8	35.7	32.7	29.2	28.1	30.4	31.4	32.8	31.5	31.3	29.3
All "Shin"	32.7	45.5	37.3	34.1	30.9	28.3	29.0	31.7	32.0	32.7	30.5	31.0	29.4
All "Gum"	33.4	47.3	39.6	36.6	33.6	29.6	27.7	29.7	31.2	32.9	32.0	31.4	29.1
Aichi	32.4	48.3	37.6	32.6	33.5	28.3	26.8	30.3	31.4	31.0	29.4	30.9	29.0
Akita	36.3	48.0	45.6	40.9	38.2	29.9	26.5	30.4	34.5	39.6	38.2	35.5	29.2
Aomori	41.4	50.2	44.6	43.6	44.2	39.6	35.9	36.8	40.1	44.6	45.9	39.3	32.3
Chiba	31.1	44.0	35.0	34.0	30.7	27.3	25.0	26.7	29.5	31.7	30.1	29.4	29.4
Ehime	34.8	50.2	41.1	36.0	32.9	29.5	28.6	32.8	33.1	35.2	33.7	33.4	31.2
Fukuoka	33.2	50.4	42.1	36.7	32.5	30.0	30.0	30.6	30.8	32.8	31.1	27.4	24.1
Fukushima	37.0	55.2	43.1	39.2	34.2	30.5	31.3	35.3	35.9	37.9	34.3	34.1	33.0
Gifu	35.6	45.2	36.5	38.2	35.9	32.8	29.2	29.8	34.1	35.7	38.0	36.8	32.0
Gumma	32.0	48.8	38.5	34.8	33.0	28.5	27.1	29.0	38.6	28.9	29.1	29.5	27.4
	31.8	45.4	37.8	34.0	31.3	29.9	27.9	28.8	29.5	30.7	28.6	29.7	28.3
Hiroshima	30.1	45.2	35.2	31.2	29.4	26.4	26.7	30.6	29.3	29.2	27.0	26.0	24.7
Hokkaido	40.3	54.9	47.5	46.8	40.5	37.5	36.3	35.6	37.3	40.8	38.8	36.0	32.5
Ibama	31.0	44.2	37.7	35.1	29.5	27.9	28.6	30.9	30.6	29.2	26.0	26.7	26.1
Ishikawa	32.1	42.4	36.5	35.3	33.6	30.1	27.1	28.2	29.8	30.4	30.2	30.1	32.1
Iwate	33.4	50.6	44.1	41.1	35.3	29.8	26.5	28.3	29.8	32.0	29.7	28.8	25.7
Kagawa	37.5	49.1	45.9	43.6	44.2	35.6	30.8	31.7	32.7	33.7	36.8	35.2	31.3
Kagoshima	35.5	52.8	39.5	35.8	29.8	26.5	27.4	31.6	32.2	31.3	28.9	27.6	26.8
Kanagawa	31.8	50.6	34.9	34.9	36.0	29.6	28.6	30.5	33.5	36.3	37.1	37.5	37.0
Kochi	29.9	44.1	35.0	33.2	29.0	28.9	29.6	29.9	29.5	31.9	29.9	30.1	30.9
		41.9	34.2	30.4	27.2	25.4	26.4	27.9	29.3	29.7	29.2	29.5	28.3

TABLE 64. - LIVE BIRTH RATES BY MONTH BY PREFECTURE: JAPAN, 1949 Cont'd
(Rates per 1,000 population per annum)

Area	Annual	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Kumamoto	34.7	49.5	40.2	35.6	31.3	29.3	28.0	31.1	33.1	35.2	33.9	37.0	32.8
Kyoto	28.9	41.6	35.4	33.2	29.5	25.6	25.4	27.4	27.0	26.3	24.6	26.0	25.7
Mie	29.3	42.8	34.5	30.3	27.8	25.1	24.5	25.8	29.5	28.0	28.8	29.2	26.0
Miyagi	35.1	47.5	42.0	40.4	37.4	29.9	26.7	28.6	30.7	34.0	36.4	35.9	31.2
Miyazaki	37.6	51.3	42.0	38.0	35.8	34.5	31.2	33.4	36.6	37.1	35.5	37.5	37.9
Nagano	27.8	40.5	34.1	30.6	28.4	26.9	23.5	23.5	24.0	27.7	26.9	25.6	22.8
Nagasaki	38.5	50.6	44.9	39.3	36.4	32.1	30.8	37.2	36.8	39.7	37.4	39.9	37.0
Nara	27.5	40.8	31.9	29.4	26.3	23.9	23.7	28.0	26.2	25.9	24.3	26.9	24.9
Niigata	33.8	43.0	40.1	40.7	34.3	30.2	26.1	28.9	32.6	37.4	35.4	32.1	25.8
Oita	33.3	50.3	41.8	35.5	34.2	28.5	27.2	30.4	31.7	32.1	30.3	30.2	27.9
Osayama	29.7	46.1	37.3	31.7	29.3	25.1	24.5	29.0	28.8	27.8	26.2	26.1	25.2
Osaka	31.4	42.5	37.1	33.4	28.5	27.8	29.9	33.2	31.4	29.7	27.9	28.6	27.4
Saga	36.0	54.8	43.1	38.1	32.8	30.3	26.9	31.5	33.4	36.4	34.7	36.0	34.6
Saitama	32.9	45.5	37.7	35.4	31.2	27.7	27.8	30.4	31.6	33.7	31.1	31.8	30.9
Shiga	28.8	40.2	36.0	34.3	29.2	27.5	24.9	25.6	27.5	26.5	25.2	24.8	24.9
Shimane	32.1	45.2	38.8	37.2	33.3	27.8	26.5	28.9	29.0	30.8	31.8	29.3	26.2
Shizuoka	33.0	47.7	37.9	31.8	33.7	28.0	27.9	29.5	30.3	32.2	32.3	32.5	32.4
Tochigi	33.9	46.0	39.1	37.7	34.0	31.0	28.7	29.8	32.6	32.7	32.1	31.5	31.3
Tokushima	33.5	50.7	40.9	36.8	31.4	28.5	27.5	29.3	30.7	31.6	31.6	31.5	31.8
Tokyo	30.9	42.6	33.8	31.5	28.5	26.4	27.9	29.5	29.7	31.3	29.0	30.5	29.3
Tottori	30.9	49.0	36.6	30.7	33.3	27.2	24.9	27.6	26.8	28.8	31.8	27.8	27.2
Toyama	33.6	47.3	40.6	40.1	36.2	31.1	26.9	29.2	32.3	34.7	31.1	29.1	25.2
Wakayama	29.0	42.1	34.3	31.5	27.3	24.2	23.4	28.2	27.9	28.9	26.3	27.7	26.5
Yamagata	32.1	43.7	35.9	35.5	32.0	28.4	26.5	28.4	30.6	34.7	33.1	31.0	26.0
Yamaguchi	32.9	47.4	38.7	34.3	31.8	27.3	27.4	32.9	33.8	33.5	30.5	29.5	28.1
Yamanashi	30.4	46.3	36.2	31.7	33.4	29.3	26.8	26.3	26.3	28.7	27.1	27.5	26.2

Data refer to births of Japanese nationals in Japan.

Rates were computed by Public Health and Welfare Section, CHQ, SCAP.

Sources: Sources of original data were Monthly Vital Statistics Schedule Reports, Ministry of Welfare.

TABLE 65. - DEATHS BY MONTH BY PREFECTURE: JAPAN, 1949

Area	Total Deaths	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
All Japan	950719	86814	77763	90267	84155	78768	74297	79443	77981	73630	72120	73347	82134
All "Shi"	317683	28793	25359	28970	27913	26044	24638	25934	25161	24226	23615	23943	27087
All "Gun"	639036	58021	52404	61297	56242	52724	49659	53509	52820	49404	48505	49404	55047
Aichi	36463	3386	2961	3514	3323	2991	2945	3312	3018	2620	2636	2748	3009
Akita	17233	1522	1375	1708	1668	1475	1252	1282	1365	1340	1324	1385	1497
Aomori	16842	1440	1211	1654	1624	1498	1373	1369	1417	1333	1237	1272	1414
Chiba	26037	2342	2182	2597	2203	2045	2006	2149	2075	1997	2028	2067	2446
Ehime	17645	1675	1546	1642	1514	1419	1478	1446	1420	1350	1343	1389	1423
Fukui	9954	899	812	951	911	830	702	875	772	857	744	738	863
Fukuoka	40041	3671	3237	3723	3532	3199	3223	3389	3189	3167	3219	3186	3406
Fukushima	23841	1987	1885	2381	2151	2034	1889	1996	1949	1778	1712	1928	2151
Gifu	18536	1593	1542	1686	1705	1571	1441	1660	1662	1415	1403	1339	1519
Gumma	18452	1661	1598	1764	1552	1433	1521	1780	1491	1454	1282	1384	1572
Hiroshima	23512	2140	1830	2145	2075	1922	1798	1933	1904	1953	1857	1871	2084
Hokkaido	48361	4386	4017	4781	4566	4348	3831	3887	4001	3768	3521	3421	3834
Hyogo	34695	3035	2765	3213	2961	2824	2781	3017	2904	2805	2768	2873	2949
Ibaraki	24854	2248	2146	2520	2095	1904	1689	2269	2071	1757	1781	1857	2217
Ishikawa	13015	1144	1048	1278	1208	1205	1008	1030	1002	1001	1013	1023	1055
Iwate	18427	1581	1443	1896	1835	1707	1479	1371	1461	1462	1241	1385	1566
Kagawa	11247	1124	972	1017	907	904	854	888	934	890	882	865	1000
Kagoshima	21455	2169	1719	1848	1662	1562	1752	1748	1878	1733	1747	1706	1931
Kanagawa	22985	2133	1955	2143	1957	1863	1709	1794	1829	1888	1805	1795	2114
Kochi	10577	1097	951	1005	844	806	767	780	862	843	812	863	947

TABLE 65. - DEATHS BY MONTH BY PREFECTURE: JAPAN, 1949 Cont'd

Area	Total Deaths	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Kumamoto	21401	1956	1674	1822	1778	1661	1554	1797	1782	1770	1686	1928	1983
Kyoto	19707	1805	1583	1942	1802	1698	1566	1555	1510	1604	1500	1520	1622
Mie	17526	1658	1536	1777	1493	1413	1297	1481	1494	1309	1333	1297	1438
Miyagi	18184	1654	1544	1936	1761	1607	1474	1431	1383	1187	1238	1400	1579
Miyazaki	12913	1284	1062	1147	1004	973	952	1065	1127	1050	1047	993	1209
Nagano	22818	2001	1921	2194	2144	1921	1746	1875	1831	1657	1786	1783	1959
Nagasaki	19709	1891	1603	1634	1676	1582	1592	1631	1597	1429	1590	1663	1821
Nara	9584	890	732	908	816	867	744	862	802	749	687	723	804
Niigata	31550	2548	2337	3100	3022	2887	2570	2764	2601	2445	2333	2441	2502
Oita	17001	1591	1482	1612	1423	1398	1301	1375	1380	1372	1306	1346	1415
Okayama	19557	1830	1677	1722	1670	1621	1397	1584	1648	1697	1617	1475	1615
Osaka	38883	3469	3041	3727	3480	3340	3162	3423	3168	2898	2947	2803	3325
Saga	12306	1196	1029	1113	1042	996	938	992	1032	928	952	980	1108
Saitama	25698	2336	2094	2345	2130	2015	1995	2344	2211	1923	1892	1998	2415
Shiga	11217	969	871	1058	1060	969	915	1022	954	862	828	759	910
Shimane	12033	1058	1013	1208	1087	1059	870	869	978	1008	950	932	1001
Shizuoka	25380	2362	2107	2423	2203	2056	1962	2188	2209	1844	1807	1956	2263
Tochigi	17855	1620	1599	1773	1519	1342	1312	1531	1503	1378	1325	1421	1532
Tokushima	11081	1088	928	1023	895	846	797	886	933	925	863	872	1025
Tokyo	54240	5407	4549	5149	4784	4427	4349	4392	4148	3975	3984	4347	4929
Tottori	7356	656	613	652	673	602	534	565	611	620	550	595	685
Toyama	14287	1123	985	1223	1339	1340	1167	1318	1196	1235	1078	1100	1183
Wakayama	11183	1062	903	1040	973	865	866	907	961	924	911	813	958
Yamagata	17656	1604	1432	1701	1685	1490	1376	1413	1431	1354	1360	1375	1435
Yamaguchi	18508	1738	1555	1711	1633	1529	1340	1537	1518	1505	1467	1328	1627
Yamanashi	8914	765	748	861	770	724	683	751	729	671	718	704	790

Data refer to deaths of Japanese Nationals in Japan

Sources: Monthly Vital Statistics Schedule Reports, Ministry of Welfare.

TABLE 66. - DEATH RATES BY MONTH BY PREFECTURE: JAPAN, 1949
(Rate per 1,000 population per annum)

Area	Annual	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
All Japan	11.6	12.4	12.3	12.9	12.5	11.3	11.0	11.4	11.2	10.9	10.3	10.9	11.8
All "Shi"	11.0	11.9	11.6	12.0	12.0	10.8	10.6	10.8	10.4	10.4	9.8	10.3	11.2
All "Gun"	11.9	12.7	12.7	13.4	12.7	11.5	11.2	11.7	11.6	11.2	10.6	11.2	12.0
Aichi	11.0	12.1	11.7	12.5	12.2	10.7	10.8	11.8	10.7	9.6	9.4	10.1	10.7
Akita	13.1	13.6	13.6	15.3	15.4	13.2	12.0	11.5	12.2	12.4	11.9	12.8	13.4
Aomori	13.5	13.6	12.6	15.6	15.8	14.1	13.4	12.9	13.4	13.0	11.7	12.4	13.3
Chiba	11.9	12.6	13.0	13.9	12.2	11.0	11.1	11.5	11.1	10.5	10.9	11.5	13.1
Ehime	11.6	13.0	13.3	12.7	12.1	11.0	11.8	11.2	11.0	10.8	10.4	11.1	11.0
Fuku	13.2	14.1	14.1	14.9	14.7	13.0	11.4	13.7	12.1	13.9	11.7	11.9	13.5
Fukuoka	11.8	12.7	12.4	12.9	12.7	11.1	11.6	11.4	11.1	11.4	11.2	11.4	11.8
Fukushima	11.5	11.3	11.8	13.5	12.6	11.5	11.1	11.3	11.1	10.4	9.7	11.3	12.2
Gifu	11.9	12.0	12.9	12.7	13.3	11.8	11.2	12.5	12.5	11.0	10.6	10.4	11.4
Gunma	11.2	11.9	12.3	12.6	11.5	10.2	11.2	12.7	10.6	10.7	9.2	10.2	11.2
Hiroshima	11.2	12.0	13.4	12.0	12.0	10.8	10.4	10.9	10.7	11.3	10.4	10.9	11.7
Hokkaido	11.7	12.5	12.7	13.7	13.5	12.4	11.3	11.1	11.4	11.1	10.1	10.1	11.0
Ibigo	10.7	11.0	11.1	11.7	11.1	10.3	10.5	11.0	10.6	10.5	10.1	10.1	10.7
Ibaraki	11.9	12.6	13.4	14.2	12.2	10.7	11.6	12.8	11.6	10.2	10.0	10.8	12.5
Ishikawa	13.5	14.0	14.2	15.6	15.2	14.7	12.7	12.6	12.2	12.6	12.4	12.9	12.9
Iwate	13.9	14.0	14.2	16.8	16.8	15.2	13.6	12.2	13.0	13.4	11.0	12.7	13.9
Kagawa	11.7	13.8	13.2	12.5	11.5	11.1	10.9	11.0	11.5	11.3	10.8	11.0	12.3
Kagoshima	11.9	14.1	12.4	12.0	11.2	10.2	11.8	11.4	12.2	11.6	11.4	11.5	12.6
Kanagawa	9.7	10.6	10.7	10.6	10.0	9.2	8.8	8.9	9.1	9.7	8.9	9.2	10.5
Kochi	11.9	14.5	14.0	13.3	11.6	10.7	10.5	10.3	11.4	11.6	10.8	11.8	12.6

TABLE 66. - DEATH RATES BY MONTH BY PREFECTURE: JAPAN, 1949 Cont'd
(Rates per 1,000 population per annum)

Area	Annual	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Kumamoto	11.7	12.6	11.9	11.7	11.8	10.7	10.3	11.6	11.5	11.8	10.9	12.8	12.8
Kyoto	10.8	11.6	11.3	12.5	12.0	10.9	10.4	10.0	9.7	10.7	9.7	10.1	10.4
Mie	11.8	13.1	13.5	14.1	12.2	11.2	11.0	11.7	11.8	10.7	10.6	10.6	11.4
Miyagi	11.1	11.9	12.2	13.9	13.1	11.6	11.0	10.3	10.0	8.8	8.9	10.4	11.4
Miyazaki	12.0	14.0	12.8	12.5	11.3	10.6	10.7	11.6	12.3	11.8	11.4	11.2	13.2
Nagano	10.7	11.1	11.8	12.1	12.2	10.6	10.0	10.4	10.1	9.5	9.9	10.2	10.8
Nagasaki	12.3	13.9	13.0	12.0	12.7	11.6	12.1	12.0	11.7	10.8	11.7	12.6	13.4
Nara	12.0	13.1	12.0	13.4	12.4	12.8	11.3	12.7	11.8	11.4	10.1	11.0	11.9
Niigata	12.6	12.0	12.2	14.6	14.7	13.6	12.5	13.0	12.3	11.9	11.0	11.9	11.8
Oita	13.3	14.7	15.1	14.9	13.6	12.9	12.4	12.7	12.7	13.1	12.0	12.8	13.1
Okayama	11.6	12.7	12.9	12.0	12.0	11.3	10.1	11.0	11.5	12.2	11.3	10.6	11.3
Oseka	10.8	11.3	11.0	12.2	11.8	10.9	10.7	11.2	10.4	9.8	9.6	9.8	10.9
Sega	12.9	14.8	14.1	13.7	13.3	12.3	12.0	12.2	12.7	11.8	11.7	12.5	13.7
Saitama	11.8	12.6	12.5	12.6	11.9	10.9	11.1	12.6	11.9	10.7	10.2	11.1	13.0
Shiga	12.5	12.8	12.7	13.9	14.4	12.8	12.4	13.5	13.1	11.7	10.9	10.3	12.0
Shimane	13.0	13.5	14.3	15.4	14.3	13.5	11.4	11.1	12.4	13.2	12.1	12.2	12.7
Shizuoka	10.3	11.3	11.1	11.6	10.9	9.8	9.7	10.4	10.5	9.1	8.6	9.6	10.8
Tochigi	11.2	11.9	13.1	13.1	11.6	9.9	10.0	11.3	11.1	10.5	9.8	10.8	11.3
Tokushima	12.4	14.4	13.6	13.5	12.2	11.2	10.9	11.7	12.3	12.6	11.4	11.9	13.5
Tokyo	9.8	11.5	10.7	10.9	10.5	9.4	9.5	9.3	8.8	8.7	8.4	9.1	10.5
Tottori	12.1	12.7	13.2	12.6	13.5	11.7	10.7	11.0	11.8	12.4	10.7	11.9	13.3
Toyama	14.0	12.9	12.6	14.1	15.9	15.4	13.9	15.2	13.8	14.7	12.4	13.1	13.6
Wakayama	11.1	12.5	11.7	12.2	11.8	10.1	10.5	10.6	11.3	11.2	10.7	9.9	11.2
Yamagata	12.8	13.7	13.5	14.5	14.9	12.7	12.1	12.1	11.9	11.9	11.6	12.1	12.2
Yamanashi	12.0	13.4	13.1	13.1	12.9	11.7	10.6	11.7	11.6	11.9	11.2	10.5	12.4
Yamaguchi	10.7	10.8	11.7	12.1	11.2	10.2	9.9	10.6	10.3	9.8	10.1	10.3	11.1

Data refer to deaths of Japanese nationals in Japan.

Sources: Rates were computed by Public Health and Welfare Section, GHQ, SCAP.

Sources of original data were Monthly Vital Statistics Schedule Reports, Ministry of Welfare.

TABLE 67. - DEATHS FROM SELECTED CAUSES BY MONTH: JAPAN, 1949

List No.	Cause of Death	Total Deaths	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
	All Causes	950719	86814	77763	90267	84155	78768	74297	79443	77981	73630	72120	73347	82134
1	Typhoid fever	970	57	57	72	54	68	82	96	147	123	90	59	65
2	Paratyphoid fever	116	7	3	13	4	4	11	16	15	20	10	9	4
3	Plague	-	-	-	-	-	-	-	-	-	-	-	-	-
4	Cholera	-	-	-	-	-	-	-	-	-	-	-	-	-
6	Cerebrospinal (meningococcus) meningitis	497	37	35	59	44	33	48	30	46	80	38	23	24
7	Anthrax	-	-	-	-	-	-	-	-	-	-	-	-	-
8	Scarlet fever	51	6	4	1	7	5	5	3	5	2	3	7	3
9	Whooping cough	9147	566	556	530	580	728	928	1070	1186	922	591	588	902
10	Diphtheria	1659	245	196	184	165	115	75	82	49	60	100	178	210
12	Tetanus	1964	136	142	136	158	156	170	188	212	217	169	145	135
13-22	Tuberculosis (all forms)	138764	11135	10106	12279	12493	13005	12287	12095	11833	11526	11240	10525	10240
13	Tuberculosis of the respiratory system	116295	9820	8625	10444	10553	10936	10196	10062	9717	9446	9397	8843	8556
23	Leprosy	162	17	15	12	18	14	12	12	6	9	15	19	13
26a	Glanders	1	-	-	-	-	1	-	-	-	-	-	-	-
27	Dysentery	7824	47	54	60	111	225	534	1850	2430	1638	569	191	115
28	Malaria	122	13	11	19	7	9	9	9	8	17	6	9	5
30	Syphilis	5492	518	467	518	467	397	390	404	393	399	477	515	547
33	Influenza	502	57	78	62	53	51	25	13	16	20	19	32	76
34	Smallpox	21	-	2	2	5	6	3	1	1	1	1	-	-
35	Measles	12440	575	681	1403	1787	2758	2209	1442	679	233	169	229	275
36	Acute poliomyelitis and acute poliomyelitis	1024	76	60	75	67	81	89	109	81	110	113	79	84
37b	Japanese "B" encephalitis	1182	3	6	8	2	5	14	24	133	174	169	21	3
38a	Yellow fever	-	-	-	-	-	-	-	-	-	-	-	-	-
38b	Rabies	76	2	4	3	6	6	5	6	12	10	10	6	6
39	Typhus fever and typhus-like diseases	67	4	8	7	5	6	10	3	9	8	3	2	2

TABLE 67. - DEATHS FROM SELECTED CAUSES BY MONTH: JAPAN, 1949 - Cont'd

List No.	Cause of Death	Total Deaths	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
45-55	Cancer and other	58890	4741	4239	4737	4647	4910	4937	5137	5718	5155	5122	4916	5127
61	malignant tumors	18835	201	179	202	175	151	130	139	127	129	152	141	159
82	Diabetes mellitus	5591	597	539	603	553	474	411	373	412	361	381	394	493
81a	Simple meningitis	9108	767	705	870	919	889	759	760	771	753	645	615	654
83	Intracranial lesions													
	of vascular origin	100650	9581	8446	9811	9006	7947	7358	7132	6724	7713	8517	8753	9631
90-95	Diseases of the heart	52825	5111	4572	5229	4078	4196	3953	3864	3677	3655	4221	4408	5071
106	Bronchitis	25605	3109	3060	3370	2844	2153	1688	1389	1136	1142	1471	1749	2584
107-109	Pneumonia (all forms)	56636	7529	6816	7277	5378	4718	3482	2928	2025	2251	2692	4043	5427
110	Pleurisy (not specified													
	as tuberculosis)	6030	515	503	612	548	537	517	540	506	451	469	422	400
117	Ulcer of the stomach													
	or duodenum	19621	1835	1592	1976	1692	1826	1475	1469	1357	1343	1612	1758	1926
119-120	Diarrhea, enteritis and													
	ulceration of intestines													
119	(all ages)	72186	4824	4046	4755	4857	5375	6737	9328	9603	7424	5327	5044	4836
	Diarrhea, enteritis and													
	ulceration of intestines													
120	(under 2 years of age)	41822	2025	2295	2396	2024	3263	4424	5676	5371	3734	2631	3069	3014
	ulceration of intestines													
	(2 years of age and over)													
121	Appendicitis	30364	1999	1751	2059	2033	2112	2343	3642	4232	3690	2695	1975	1822
130-132	Nephritis	33859	266	231	284	270	290	255	334	348	305	238	227	215
			3265	2910	3330	2887	2587	2421	2514	2473	2591	2886	2926	3069

TABLE 67. - DEATHS FROM SELECTED CAUSES BY MONTH: JAPAN, 1949 - Cont'd

List No.	Cause of Death	Total Deaths	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
140-150	Diseases of pregnancy, child birth and the puerperium	4495	404	346	406	413	373	342	345	415	361	369	354	367
140-142a and 147	Puerperal fever	543	44	43	55	68	51	37	45	51	43	40	31	35
143	Hemorrhage of pregnancy (death before delivery)	237	30	24	17	17	24	19	22	26	29	10	10	9
144	Toxemias of pregnancy	1354	142	106	142	108	107	104	106	115	139	95	110	110
146	Hemorrhage of childbirth and the puerpium													
	(gestation of 8 months or over or unspecified)	891	68	76	85	75	71	62	59	85	76	81	73	80
147	Infection during child-birth and the puerperium (gestation of 3 months or over or unspecified)													
	Toxemias during child-birth and the puerperium (excluding death before delivery)	181	13	7	18	12	6	21	18	17	19	26	14	10
157	Congenital malformations	6586	641	573	678	560	493	463	459	411	495	536	605	642
158	Congenital debility	37285	4885	4024	4664	3397	2524	2224	2385	2174	2000	2298	3009	3701
159	Premature birth	13667	1484	1272	1362	1112	1015	972	1003	1002	989	963	1117	1376
160	Injury at birth	1082	89	91	92	96	75	85	95	90	91	101	85	92
162	Senility	66699	6635	5893	6815	5927	4749	4268	4558	4784	4861	5545	5853	6811
163-164	Suicide	14211	864	860	1118	1342	1593	1250	1396	1288	1169	1140	1091	1100
165-168	Homicide	1669	199	119	145	132	150	115	144	171	164	135	143	142
169-195	Accidental deaths	34498	2253	1984	2520	2537	2871	3125	4320	4463	3166	2528	2240	2491

TABLE 67. - DEATHS FROM SELECTED CAUSES BY MONTH: JAPAN, 1949 - Cont'd

List No.	Cause of Death	Total Deaths	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
199-200	Sudden death, ill-defined and unknown causes	16984	1659	1506	1721	1605	1340	1283	1307	1243	1119	1259	1364	1573
199	Sudden death	48	7	2	4	5	2	5	5	5	3	4	3	3
200a	Ill-defined causes	16548	1617	1473	1690	1577	1300	1247	1270	1200	1078	1222	1344	1540
200b	Found dead (cause unknown)	36	-	1	1	5	5	2	4	2	1	7	3	5
200c	Unspecified causes	352	35	30	36	18	33	34	28	36	37	26	14	25
	All other causes	125349	11949	10771	12383	11347	9869	9236	10170	10182	9764	9717	9453	10508

Footnotes: Data refer to deaths of Japanese nationals in Japan.

A dash (-) indicates that no deaths were reported.

Sources: Monthly Vital Statistics Schedule Reports, Ministry of Welfare.

TABLE 68. - DEATH RATES FOR SELECTED CAUSES BY MONTH: JAPAN, 1949
(Rates per 100,000 population per annum)

List No.	Cause of Death	Annual	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
	All Causes	1156.6	1247.5	1233.2	1233.0	1245.6	1123.3	1099.7	1137.9	1117.0	1089.8	1037.0	1085.6	1176.5
1	Typhoid fever	1.2	0.8	0.9	1.0	0.8	1.0	1.2	1.4	2.1	1.8	1.3	0.9	0.9
2	Paratyphoid fever	0.1	0.1	0.0	0.2	0.1	0.1	0.2	0.2	0.2	0.3	0.1	0.1	0.1
3	Plague	-	-	-	-	-	-	-	-	-	-	-	-	-
4	Cholera	-	-	-	-	-	-	-	-	-	-	-	-	-
6	Cerebrospinal (meningo-coccus) meningitis	0.6	0.5	0.6	0.8	0.7	0.5	0.7	0.4	0.7	1.2	0.5	0.3	0.3
7	Anthrax	-	-	-	-	-	-	-	-	-	-	-	-	-
8	Scarlet fever	0.1	0.1	0.1	0.0	0.1	0.1	0.1	0.0	0.1	0.0	0.0	0.1	0.1
9	Whooping cough	11.1	8.1	8.8	7.6	8.6	10.4	13.7	15.3	17.0	13.6	8.5	8.7	12.9
10	Diphtheria	2.0	3.5	3.1	2.6	2.4	1.6	1.1	1.2	0.7	0.9	1.4	2.6	3.0
12	Tetanus	2.4	1.9	2.3	1.9	2.3	2.2	2.5	2.7	3.0	3.2	2.4	2.1	1.9
13-22	Tuberculosis (all forms)	168.8	159.5	160.3	175.9	184.9	186.3	181.9	173.2	169.5	170.6	161.0	155.8	146.7
13	Tuberculosis of the respiratory system	141.5	136.4	135.8	149.6	156.2	156.6	150.9	144.1	139.2	139.8	134.6	130.9	122.6
23	Leprosy	0.2	0.2	0.2	0.2	0.3	0.2	0.2	0.2	0.1	0.1	0.2	0.3	0.2
26a	Glanders	0.0	-	-	-	-	-	-	-	-	-	-	-	-
27	Dysentery	9.5	0.7	0.9	0.9	1.6	3.2	7.9	26.5	34.8	24.2	8.2	2.8	1.6
28	Malaria	0.1	0.2	0.2	0.3	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
30	Syphilis	6.7	7.4	7.4	7.4	6.9	5.7	5.8	5.8	5.6	5.9	6.8	7.6	7.8
33	Influenza	0.6	0.8	1.2	0.9	0.8	0.7	0.4	0.2	0.2	0.3	0.4	0.4	1.1
34	Smallpox	0.0	-	0.0	0.0	0.1	0.1	0.0	0.0	0.0	-	0.0	-	-
35	Measles	15.1	82.	10.8	20.1	26.4	39.5	32.7	20.7	9.7	3.4	2.4	3.4	3.9
36	Acute poliomyelitis and													
37b	Acute poliomyelitis	1.2	1.1	1.0	1.1	1.0	1.2	1.3	1.6	1.2	1.6	1.6	1.2	1.2
38a	Japanese "B" encephalitis	1.4	0.0	0.1	0.1	0.0	0.1	0.2	0.3	2.2	11.5	2.4	0.3	0.0
38b	Yellow fever	-	-	-	-	-	-	-	-	-	-	-	-	-
39	Rabies	0.1	0.0	0.1	0.0	0.1	0.1	0.1	0.1	0.2	0.1	0.1	0.1	0.1
39	Typhus fever and typhus-like diseases	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.0	0.1	0.1	0.0	0.0	0.0

TABLE 68. - DEATH RATES FOR SELECTED CAUSES BY MONTH: JAPAN, 1949 - Cont'd
(Rates per 100,000 population per annum)

List No.	Cause of Death	Annual	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
45-55	Cancer and other malignant tumors	71.6	67.9	67.2	67.9	68.8	70.3	71.6	73.6	76.2	76.3	73.4	72.8	73.4
61	Diabetes mellitus	2.3	2.9	2.8	2.9	2.6	2.2	1.9	2.0	1.8	1.9	2.2	2.1	2.3
68	Beriberi	6.8	8.6	8.5	8.6	8.2	6.8	6.1	5.3	5.9	5.3	5.5	5.8	7.1
81a	Simple meningitis	11.1	11.0	11.2	12.5	13.6	12.7	11.2	10.9	11.0	11.1	9.2	9.1	9.4
85	Intracranial lesions of vascular origin	122.4	137.2	133.9	140.5	133.3	113.8	108.9	102.2	96.3	114.2	122.0	129.6	138.4
90-95	Diseases of the heart	64.3	73.2	72.5	74.9	72.2	60.0	58.5	55.3	52.7	54.1	60.5	65.2	72.6
106	Bronchitis	31.3	44.5	48.5	48.3	42.1	30.8	25.0	19.9	16.3	16.9	21.1	25.9	37.0
107-109	Pneumonia (all forms)	68.9	107.8	108.1	104.2	94.4	70.4	51.1	40.5	29.0	33.3	33.6	58.8	92.1
110	Pleurisy (not specified as tuberculosis)	7.3	7.4	8.0	8.8	8.1	7.7	7.7	7.7	7.2	6.8	6.7	6.2	5.7
117	Ulcer of the stomach or duodenum	23.7	26.3	25.2	26.3	25.0	23.3	21.8	21.0	19.4	19.9	23.1	26.0	27.6
119-120	Diarrhea, enteritis and ulceration of intestines (all ages)	87.8	69.1	64.2	68.1	71.9	77.0	100.2	133.6	137.6	109.9	76.3	74.7	69.3
119	Diarrhea, enteritis and ulceration of intestines (under two years of age)	50.9	40.4	36.4	36.6	41.8	45.7	65.5	81.3	76.9	55.3	37.7	45.4	43.2
120	Diarrhea, enteritis and ulceration of intestines (2 years of age and over)	36.9	28.6	27.8	29.5	30.1	30.3	34.7	52.3	60.6	54.6	38.6	29.2	26.1
121	Appendicitis	4.0	3.8	3.7	4.1	4.0	4.2	3.8	4.8	5.0	4.5	4.3	3.4	3.1
130-132	Nephritis (all forms)	41.2	46.8	46.1	47.7	42.7	37.1	35.8	36.0	35.4	38.4	41.3	43.3	44.0

TABLE 68. - DEATH RATES FOR SELECTED CAUSES BY MONTH: JAPAN, 1949 - Cont'd
(Rates per 100,000 population per annum)

List No.	Cause of Death	Annual	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
140-150	Diseases of pregnancy, childbirth and the puerperium	5.5	5.8	5.5	5.8	6.1	5.3	5.1	4.9	5.9	5.3	5.3	5.2	5.3
140-142a and 147	Puerperal fever	0.7	0.6	0.7	0.8	1.0	0.7	0.4	0.6	0.7	0.6	0.6	0.5	0.5
143	Hemorrhage of pregnancy (death before delivery)	0.3	0.4	0.4	0.2	0.3	0.3	0.3	0.3	0.4	0.4	0.1	0.1	0.1
144	Toxemias of pregnancy	1.6	2.0	1.7	2.0	1.6	1.5	1.5	1.5	1.6	1.6	1.4	1.6	1.6
146	Hemorrhage of childbirth and the puerperium													
	(restoration of 8 months or over or unspecified)	1.1	1.0	1.2	1.2	1.1	1.0	0.9	0.8	1.2	1.1	1.2	1.1	1.1
147	Infection during childbirth and the puerperium													
	(restoration of 3 months or over or unspecified)	0.5	0.4	0.5	0.5	0.7	0.5	0.3	0.5	0.5	0.5	0.3	0.3	0.3
148	Toxemias during childbirth and the puerperium (excluding death before delivery)	0.2	0.2	0.1	0.3	0.2	0.1	0.3	0.3	0.2	0.3	0.4	0.2	0.1
157	Congenital malformation	8.0	9.2	9.1	9.7	8.3	7.1	6.9	6.6	5.9	7.3	7.7	9.0	9.2
158	Constitutional debility	45.4	70.0	63.8	66.8	50.3	36.2	32.9	34.2	31.1	29.6	32.9	44.5	53.0
159	Premature birth	16.6	21.3	20.2	19.5	16.5	14.5	14.4	14.4	14.4	14.6	13.8	16.5	19.7
160	Injury at birth	1.3	1.3	1.4	1.3	1.4	1.1	1.3	1.4	1.3	1.3	1.4	1.3	1.3
162	Senility	81.1	95.0	93.5	97.6	87.7	68.0	63.2	65.3	68.5	71.9	79.4	86.6	97.6
163-164	Suicide	17.3	19.4	18.6	16.0	19.9	22.8	18.5	20.0	18.4	17.3	16.3	16.1	15.8
165-168	Homicide	2.0	1.6	1.9	2.1	2.0	2.1	1.7	2.1	2.4	2.4	1.9	2.1	2.0
169-195	Accidental deaths	42.0	32.3	31.5	36.1	37.6	41.1	46.3	61.9	63.9	46.9	36.2	33.2	35.7

TABLE 68. - DEATH RATES FOR SELECTED CAUSES BY MONTH: JAPAN, 1949 - Cont'd
(Rates per 100,000 population per annum)

List No.	Cause of Death	Annual	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
199-200 a,b,c	Sudden death, ill-defined and unknown causes	20.7	23.8	23.9	24.7	23.8	19.2	19.1	18.7	17.8	16.6	18.0	20.2	22.5
199	Sudden death	0.1	0.1	0.0	0.1	0.1	0.0	0.1	0.1	0.1	0.0	0.1	0.0	0.0
200a	Ill-defined causes	20.1	23.2	23.4	24.1	23.3	18.6	18.5	18.2	17.2	16.0	17.5	19.9	22.1
200b	Found dead (cause unknown)	0.0	-	0.0	0.0	0.1	0.1	0.0	0.1	0.0	0.0	0.1	0.0	0.1
200c	Unspecified causes	0.4	0.5	0.5	0.5	0.3	0.5	0.5	0.4	0.5	0.5	0.4	0.2	0.4
	All other causes	152.5	171.2	170.8	177.4	168.0	141.4	136.7	145.7	145.8	144.5	139.2	139.9	150.5

Footnotes:

Base data refer to deaths of Japanese nationals in Japan.

A dash (-) indicates that no deaths were reported.

A rate of 0.0 indicates that there were some deaths but the rate was less than 0.1.

Sources:

Rates were computed by Public Health and Welfare Section, GHQ, SCAP.

Sources of original data were Monthly Vital Statistics Schedule Report, Ministry of Welfare.

TABLE 69.- DEATHS FROM SELECTED CAUSES BY PREFECTURE: JAPAN, 1949

Area	All Causes	Typhus	Paratyphoid	Plague	Cholera	Cerebrospinal (meningococcus)			Anthrax	Scarlet	Whooping	Diphtheria	Tetanus	Tuberculosis
		fever (1)	fever (2)	(3)	(4)	Meningitis (6)	(7)	fever (8)	cough (9)	(10)	(12)	(13-22)		
All Japan	950,719	970	116	-	-	497	-	51	9,147	1,659	1,964	138,764	1,964	138,764
All "Shi"	311,683	536	57	-	-	294	-	26	3,412	703	676	60,630	676	60,630
All "Gun"	639,036	434	59	-	-	203	-	25	5,735	956	1,288	78,134	1,288	78,134
Aichi	36,463	47	-	-	-	6	-	-	402	28	72	5,383	72	5,383
Akita	17,233	8	2	-	-	8	-	1	164	31	20	2,120	20	2,120
Aomori	16,842	22	4	-	-	11	-	1	236	49	26	2,710	26	2,710
Chiba	26,037	27	2	-	-	11	-	3	227	31	118	2,961	118	2,961
Ehime	17,645	7	1	-	-	4	-	-	233	24	62	2,268	62	2,268
Fukui	9,954	11	1	-	-	1	-	-	66	16	11	1,166	11	1,166
Fukuoka	40,041	23	2	-	-	21	-	1	484	86	74	6,918	74	6,918
Fukushima	23,841	19	6	-	-	18	-	1	302	40	37	2,997	37	2,997
Gifu	18,536	28	3	-	-	4	-	3	176	27	30	2,575	30	2,575
Gumma	18,452	10	2	-	-	8	-	-	297	32	66	2,007	66	2,007
Hiroshima	23,512	44	3	-	-	10	-	-	175	36	47	3,324	47	3,324
Hokkaido	48,361	33	3	-	-	51	-	5	634	142	50	10,284	50	10,284
Ibogo	34,695	58	1	-	-	12	-	3	340	63	40	5,478	40	5,478
Ibaraki	24,854	15	2	-	-	10	-	-	124	25	103	2,315	103	2,315
Ishikawa	13,015	8	2	-	-	2	-	-	131	20	25	1,631	25	1,631
Iwate	18,427	13	5	-	-	8	-	1	183	43	15	2,737	15	2,737
Kagawa	11,247	8	1	-	-	2	-	-	91	17	40	1,398	40	1,398
Kagoshima	21,455	4	-	-	-	5	-	-	103	63	99	2,771	99	2,771
Kanagawa	22,985	36	4	-	-	23	-	2	271	45	51	4,240	51	4,240
Kochi	10,577	21	2	-	-	6	-	-	32	11	32	1,358	32	1,358

TABLE 69.- DEATHS FROM SELECTED CAUSES BY PREFECTURE: JAPAN, 1949

Area	All Causes	Typhus		Paratyphoid		Flague	Cholera	Cerebrospinal (meningococcus)		Anthrax	Scarlet Whooping cough		Diphtheria	Tetanus	Tuberculosis (all forms)
		fever (1)	fever (2)	fever (3)	(4)			(5)	(6)		(7)	(8)			
Kumamoto	21,401	3	1	-	-	-	-	5	-	-	1	99	39	42	2,736
Kyoto	19,707	20	-	-	-	-	-	22	-	-	1	162	13	19	3,915
Mie	17,526	34	7	-	-	-	-	7	-	-	-	113	17	33	2,239
Miyagi	18,184	25	8	-	-	-	-	14	-	-	1	290	29	28	2,690
Miyazaki	12,913	3	-	-	-	-	-	5	-	-	1	46	55	60	1,810
Nagano	22,818	17	1	-	-	-	-	8	-	-	1	241	22	50	2,620
Nagasaki	19,709	10	-	-	-	-	-	7	-	-	1	109	61	48	2,717
Nara	9,584	21	-	-	-	-	-	1	-	-	1	76	17	16	1,232
Niigata	31,550	25	4	-	-	-	-	5	-	-	1	259	56	41	4,237
Oita	17,001	4	1	-	-	-	-	6	-	-	-	97	42	34	2,248
Okayama	19,557	45	2	-	-	-	-	4	-	-	1	96	27	36	2,443
Osaka	26,883	55	6	-	-	-	-	50	-	-	4	433	45	42	7,259
Saga	12,306	6	-	-	-	-	-	4	-	-	-	92	39	41	1,643
Saitama	25,698	31	5	-	-	-	-	11	-	-	3	344	16	59	2,918
Shiga	11,217	7	3	-	-	-	-	3	-	-	-	121	9	7	1,350
Shimane	12,033	8	1	-	-	-	-	5	-	-	1	53	18	14	1,772
Shizuoka	25,360	25	6	-	-	-	-	11	-	-	1	302	37	67	3,204
Tochigi	17,855	19	2	-	-	-	-	2	-	-	2	177	27	48	1,852
Tokushima	11,081	10	-	-	-	-	-	-	-	-	-	40	15	32	1,533
Tokyo	54,240	99	16	-	-	-	-	72	-	-	7	574	127	99	11,632
Tottori	7,356	4	-	-	-	-	-	10	-	-	1	57	13	7	1,013
Toyama	14,287	17	2	-	-	-	-	2	-	-	-	151	28	11	1,906
Wakayama	11,183	16	3	-	-	-	-	-	-	-	1	31	5	18	1,422
Yamagata	17,656	8	1	-	-	-	-	14	-	-	1	296	21	20	2,048
Yamaguchi	18,508	11	-	-	-	-	-	4	-	-	-	166	39	46	2,915
Yamanashi	8,914	3	1	-	-	-	-	4	-	-	-	51	11	29	769

See footnotes at end of table.

TABLE 69.- DEATHS FROM SELECTED CAUSES BY PREFECTURE, JAPAN, 1949 - Cont'd

Area	Leptosy (23)	Giardiasis (24)	Dysentery (25)	Malaria (26)	Syphilis (27)	Influenza (28)	Smallpox (29)	Measles (30)	Acute poliomyelitis (31)	Japanese B* Yellow encephalitis fever (32)	Hepatitis (33)	Typhus fever and typhus- like diseases (34)
All Japan	162	1	7,824	122	5,492	502	21	12,440	1,024	1,182	76	67
All "Shi"	13	-	2,644	29	2,570	135	16	4,629	321	639	33	22
All "Gun"	149	1	5,180	93	2,922	367	5	7,811	703	543	43	45
Aichi	7	-	464	4	199	16	-	557	56	72	-	1
Akita	3	-	54	4	77	8	-	238	18	7	-	1
Aomori	1	-	49	-	83	24	-	239	23	9	-	3
Chiba	1	-	310	3	179	13	1	274	27	21	19	3
Ehime	3	-	101	1	67	23	1	137	21	18	-	6
Fukui	2	-	34	2	32	7	-	78	6	26	-	-
Fukuoka	1	-	144	5	330	27	-	820	46	31	-	2
Fukushima	2	-	213	3	117	7	-	378	25	1	-	-
Gifu	-	-	226	3	85	9	-	160	18	23	-	-
Gumma	12	-	401	2	109	8	-	283	26	6	7	-
Hiroshima	1	-	209	3	93	20	-	164	31	51	-	3
Hokkaido	2	-	96	3	331	17	-	1,156	59	2	-	-
Hyogo	3	-	149	7	177	15	2	393	38	71	1	4
Ibaraki	1	-	552	1	128	1	-	262	23	24	3	-
Ishikawa	1	-	50	1	55	6	-	180	9	10	-	-
Iwate	2	-	118	-	76	8	-	216	20	2	-	-
Kagawa	1	-	102	-	54	11	-	27	14	15	-	-
Kagoshima	9	-	98	5	139	11	-	228	34	15	-	-
Kanagawa	-	-	240	2	197	13	-	279	28	45	8	-
Kochi	3	-	55	2	57	4	-	21	10	1	-	-

TABLE 69.- DEATHS FROM SELECTED CAUSES BY PREFECTURE, JAPAN, 1949 - Cont'd

Area	Leprosy (23)	Glanders (26a)	Dysentery (27)	Malaria (28)	Syphilis (30)	Influenza (33)	Smallpox (34)	Measles (35)	Acute polio- myelitis and acute polio- encephalitis (36)	Japanese #3* Yellow encephalitis (37b)	Yellow fever (38a)	Rabies (38b)	Typhus fever and typhus- like diseases (39)
Kumamoto	56	-	110	2	107	16	-	225	24	80	-	-	3
Kyoto	-	-	88	-	165	9	-	358	10	18	-	-	3
Mie	4	-	89	5	89	12	-	257	13	26	-	-	1
Miyagi	9	1	53	2	117	2	2	240	25	6	-	-	4
Miyazaki	3	-	136	-	81	7	-	97	17	35	-	-	-
Nagano	-	-	92	4	112	6	-	173	25	22	-	-	-
Nagasaki	6	-	87	4	145	9	-	311	21	7	-	-	-
Nara	-	-	25	2	62	4	-	198	12	15	-	-	-
Niigata	2	-	372	3	104	19	-	575	24	11	-	-	3
Oita	4	-	105	1	82	12	-	244	39	8	-	-	4
Okayama	-	-	123	1	86	7	-	54	21	35	-	-	11
Osaka	1	-	153	4	300	22	7	178	45	52	-	-	2
Saga	-	-	52	1	84	14	-	154	13	8	-	-	-
Seitama	3	-	575	2	123	3	-	271	17	40	-	14	1
Shiga	2	-	18	6	43	3	-	188	8	30	-	-	-
Shimane	-	-	124	4	61	10	-	145	11	17	-	-	1
Shizuoka	2	-	355	1	147	19	1	264	38	64	-	-	2
Tochigi	1	-	307	4	124	7	1	98	19	7	-	1	-
Tokushima	2	-	71	6	58	6	-	47	8	28	-	-	-
Tokyo	4	-	835	4	511	27	2	605	43	86	-	22	4
Tottori	3	-	34	2	34	1	-	30	2	4	-	-	-
Toyama	-	-	68	1	45	12	-	211	11	44	-	-	-
Wakayama	1	-	42	3	49	7	1	169	11	29	-	-	4
Yamagata	2	-	69	1	63	4	-	195	11	2	-	-	-
Yamaguchi	-	-	127	6	85	12	1	231	18	46	-	-	-
Yamanashi	2	-	60	2	30	4	-	132	6	12	-	-	1

See footnotes at end of table.

TABLE 69.- DEATHS FROM SELECTED CAUSES BY PREFECTURE: JAPAN, 1949 - Cont'd

Area	Cancer and other malignant tumors (45-55)	Diabetes mellitus (61)	Intracranial		Pneumon- not specifi-		Fleurisy (110)	Ulcer of stomach or of intestines duodenum (all ages)		Diarrhea en-teritis and ulceration of intestines (under 2 yrs.)		Diarrhea en-teritis and ulceration of intestines (2 yrs. and over)	
			Simple menin- gitis (81a)	of vascular origin (83)	of the Bron- chitis forms) (90-95)	ia (all fied as tu- berculous (107-109)		(117)	(119-120)	(119)	(120)		
All Japan	58,890	1,885	5,591	9,108	100,650	52,825	25,695	56,696	6,030	19,521	72,186	41,822	30,364
All 'Shi'	22,356	517	1,828	2,923	28,366	16,719	6,069	19,416	1,944	6,436	19,130	11,996	7,134
All 'Gun'	36,534	1,368	3,763	6,185	72,284	36,106	19,626	37,220	4,086	13,085	53,056	29,826	23,230
Aichi	2,368	88	222	445	3,906	2,198	1,058	2,258	221	661	2,376	1,342	1,034
Akita	735	18	135	213	2,476	911	488	832	98	240	1,655	1,171	484
Aomori	644	15	207	130	1,712	662	524	1,149	98	218	1,798	1,426	372
Chiba	1,802	51	113	290	3,340	1,642	657	1,568	182	637	1,421	750	671
Edo	1,110	37	46	199	1,614	969	545	1,082	103	299	1,203	594	609
Fuku	554	27	55	103	902	520	285	490	60	184	1,087	560	527
Fukuoka	2,642	80	216	322	3,594	1,847	869	2,508	263	860	2,897	1,977	920
Fukushima	1,334	30	119	260	3,019	1,337	660	1,792	175	476	1,757	1,112	645
Gifu	1,047	37	86	267	1,918	1,051	616	981	123	337	1,585	682	603
Gumma	1,180	29	70	155	2,201	1,236	442	1,342	85	397	1,246	543	703
Hiroshima	1,557	81	128	234	2,415	1,322	634	1,154	138	588	1,665	819	846
Hokkaido	2,445	85	347	593	3,670	2,250	3,276	3,276	305	600	4,383	3,231	1,152
Ibigo	2,270	68	214	363	3,112	1,860	885	1,798	233	710	2,942	1,592	1,350
Ibaraki	1,570	43	135	180	3,348	1,829	798	1,308	171	594	1,680	922	758
Ishikawa	805	45	113	159	1,235	711	425	747	61	294	1,167	793	374
Iwate	635	21	187	177	2,407	865	599	1,367	90	302	1,582	1,200	382
Kagawa	769	38	18	126	1,116	624	275	590	61	253	704	333	371
Kagoshima	997	37	104	199	2,324	1,122	578	1,285	186	507	1,815	1,072	713
Kanagawa	1,586	38	113	210	2,514	1,263	392	1,647	139	547	1,094	631	463
Kochi	610	53	67	104	1,284	456	262	482	77	194	639	254	385

TABLE 69.-DEATHS FROM SELECTED CAUSES BY PREFECTURE: JAPAN, 1949 - Cont'd

Area	Cancer and other malign- ant tumors (45-55)	Diabetes mellitus (61)	Beriberi (68)	Simple menin- geal lesions (81a)	Intracran- ial lesions (83)	Diseases of the heart (90-95)	Bron- chitis (106)	Pneumon- ia (all forms) (107-109)	Pleurisy not speci- fied as tu- berculosis (110)	Ulcer of stomach or duodenum (117)	Diarrhea en- teritis and ulceration of intestines (all ages) (119-120)	Diarrhea en- teritis and ulceration of intestines (under 2 yrs.) (119)	Diarrhea en- teritis and ulceration of intestines (2 yrs. and over) (120)
Kumamoto	1,223	40	96	169	2,258	1,239	599	1,259	197	535	1,684	849	835
Kyoto	1,420	43	146	186	1,732	1,054	362	990	110	429	1,282	695	597
Maie	1,098	34	110	207	1,788	1,038	650	888	101	409	1,450	775	675
Miyagi	1,205	24	145	144	2,284	1,086	383	1,250	97	304	1,247	884	363
Miyazaki	658	22	58	143	1,279	769	364	678	102	290	1,010	562	448
Nagano	1,704	44	77	152	3,223	1,604	625	1,272	121	533	1,403	610	793
Nagasaki	1,050	45	91	208	1,660	995	644	1,265	158	388	1,418	958	460
Nara	793	17	58	92	960	520	237	489	46	221	812	450	362
Niigata	2,075	72	216	310	3,985	1,651	1,161	1,880	201	555	2,803	1,440	1,363
Oita	885	35	82	186	1,697	910	575	936	130	431	1,324	701	623
Okeayama	1,308	58	111	202	2,077	1,077	551	876	147	447	1,474	557	917
Osaka	2,771	52	387	328	3,036	2,146	702	2,731	205	806	2,934	1,765	1,169
Saga	754	25	110	110	1,217	616	340	562	71	263	934	645	289
Saitama	1,652	40	103	216	3,004	1,607	917	1,570	158	502	1,978	928	1,050
Shiga	679	24	73	80	1,083	657	308	522	63	244	1,204	604	600
Shimane	708	40	51	94	1,376	743	353	577	83	301	939	467	472
Shizuoka	1,418	48	134	283	2,863	1,477	644	1,517	131	539	1,830	1,003	827
Tochigi	1,189	22	80	131	2,271	1,085	561	1,118	104	408	1,626	717	909
Tokushima	646	24	30	125	962	682	344	620	72	216	853	364	469
Tokyo	3,980	75	302	328	5,241	3,131	849	3,657	352	1,129	2,244	1,539	705
Tottori	505	26	52	52	813	381	193	307	54	180	608	319	289
Toyama	837	34	113	153	1,307	700	504	814	76	266	1,757	1,028	729
Wakayama	859	35	35	86	1,173	597	269	562	66	234	988	550	498
Yamagata	1,075	28	120	162	2,382	995	587	1,068	118	321	1,728	1,083	645
Yamanashi	1,081	40	102	178	1,825	973	417	1,056	149	468	1,224	716	508
Yamanashi	617	17	14	54	1,027	617	304	536	69	204	736	398	338

See footnotes at end of table.

TABLE 69.- DEATHS FROM SELECTED CAUSES BY PREFECTURE: JAPAN, 1949 - Cont'd

Area	Diseases of										Sudden death, ill-defined and unknown causes (199-200a,b,c)	All other causes
	Appendicitis (121)	Nephritis (130-132)	pregnancy and the puerperium (140-150)	Congenital debility (158)	Premature birth (159)	Injury at birth (160)	Senility (162)	Suicide (163-164)	Homicide (165-168)	Accidental deaths (169-195)		
All Japan	3,267	33,859	4,495	6,556	13,667	1,082	66,699	14,211	1,669	34,498	16,984	125,349
All "Shi"	1,554	10,598	1,605	2,262	4,252	414	16,366	5,320	715	11,555	5,677	38,376
All "Gun"	1,713	23,321	2,890	4,294	9,415	668	50,333	8,891	954	22,943	11,307	86,973
Aichi	118	1,010	115	254	603	40	2,685	558	62	1,168	540	4,511
Akita	54	674	121	99	327	22	1,036	201	21	567	289	2,077
Aomori	53	383	93	125	251	11	965	133	19	490	381	2,157
Chiba	97	1,076	123	174	525	42	2,188	377	87	719	462	3,332
Ehime	47	562	77	130	252	28	1,776	235	25	769	297	2,589
Fukui	31	276	45	57	156	9	895	157	10	428	148	1,419
Fukuoka	133	1,478	175	249	484	36	2,348	534	132	1,945	674	5,203
Fukushima	84	737	130	209	379	23	1,599	254	35	933	443	2,945
Gifu	51	521	75	121	321	19	1,399	340	28	610	271	2,440
Gunma	51	685	82	171	227	21	1,140	327	29	630	273	2,550
Hiroshima	83	811	102	166	288	30	2,024	371	35	938	365	3,363
Hokkaido	184	1,115	291	367	519	46	2,269	557	92	2,232	1,213	5,790
Iyogo	128	1,158	180	211	506	35	2,533	693	71	1,361	670	4,678
Ibaraki	80	1,075	119	195	518	22	1,871	260	54	719	413	3,363
Ishikawa	45	369	50	90	207	12	1,076	162	14	398	178	2,033
Iwate	63	505	106	117	373	18	1,104	198	24	732	320	2,251
Kagawa	37	419	44	73	166	16	1,056	140	21	441	302	1,611
Kanagawa	45	1,046	148	123	707	22	1,664	224	18	861	324	3,328
Kanagawa	86	772	110	159	546	34	1,398	445	57	1,084	463	2,610
Kochi	20	426	54	56	191	20	1,179	132	20	446	203	1,564

TABLE 69.- DEATHS FROM SELECTED CAUSES BY PREFECTURE, JAPAN, 1949 - Cont'd

Area	Appendi- citis (121)	Nephritis (all forms) (130-132)	Diseases of pregnancy childbirth and the (malforma- tion) (140-150)	Congenital debility (158)	Premature birth (159)	Injury at birth (160)	Sani- tary (162)	Suicide (163-164)	Homicide (165-168)	Accidental deaths (169-195)	Sudden death, ill- defined and unknown causes (199-200a,b,c)		all other causes
Kumamoto	61	871	124	129	807	239	23	1,693	238	69	761	306	3,172
Kyoto	89	563	82	135	660	298	37	1,418	479	33	573	288	2,495
Mie	44	617	76	113	698	289	18	1,262	251	31	523	342	2,556
Miyagi	74	619	83	171	798	271	35	1,070	172	28	693	312	2,153
Miyazaki	40	622	87	91	595	170	20	928	152	14	466	174	1,893
Nagano	82	947	102	143	642	291	16	1,731	516	40	790	385	2,981
Nagasaki	42	697	98	135	889	239	17	1,709	182	30	822	296	3,087
Nara	31	394	30	50	408	133	10	713	170	10	253	243	1,212
Niigata	144	1,155	128	196	1,045	418	29	1,729	503	28	1,036	513	3,976
Oite	61	661	74	92	707	212	22	1,493	221	29	548	252	2,507
Okayama	69	659	79	128	890	288	24	1,971	279	21	695	308	2,826
Osaka	179	1,582	186	267	1,385	559	37	2,033	709	60	1,277	686	4,467
Saga	45	579	48	90	669	172	17	853	125	24	532	210	1,789
Saitama	70	1,016	133	198	1,137	400	24	1,770	391	42	669	431	3,293
Shiga	43	425	28	54	489	167	14	870	221	15	263	193	1,700
Shimane	41	434	50	59	460	181	10	968	163	15	414	151	1,577
Shizuoka	68	878	109	200	1,026	332	21	1,859	476	50	885	440	3,586
Tochigi	58	676	86	156	605	228	15	1,216	261	29	566	324	2,342
Tokushima	22	397	48	72	542	172	8	1,036	165	12	409	214	1,564
Tokyo	289	2,068	266	385	1,376	702	80	2,599	1,137	131	1,900	1,199	6,082
Tottori	25	374	37	62	293	93	10	730	78	12	236	83	937
Toyama	46	510	54	102	596	268	26	664	201	9	570	250	1,921
Wakayama	38	267	37	75	347	162	15	972	263	12	338	268	1,673
Yamagata	61	667	69	104	905	313	29	1,017	249	18	582	268	2,034
Yamaguchi	54	679	79	113	676	149	13	1,591	297	39	888	332	2,378
Yamanashi	32	404	42	88	265	99	6	599	102	14	358	189	1,404

See footnotes at end of table.

TABLE 69.- DEATHS FROM SELECTED CAUSES BY PREFECTURE; JAPAN, 1949 - Cont'd

FOOTNOTES:

Data refer to deaths of Japanese nationals in Japan.
A dash (-) indicates that no deaths were reported.

SOURCES:

Monthly Vital Statistics Schedule Reports, Ministry of Welfare.

TABLE 70.- DEATH RATES FOR SELECTED CAUSES BY PREFECTURE, JAPAN, 1947

Area	All Causes	Typhoid			Cerebrospinal (Meningococcus)			Scarlet fever (8)	Whooping cough (9)	Diphtheria (10)	Tetanus (12)	Tuberculosis all forms (13-22)
		fever (1)	Paratyphoid fever (2)	Plague (3)	Cholera (4)	meningitis (6)	Anthrax (7)					
All Japan	1156.6	1.2	0.1	-	-	0.6	-	0.1	11.1	2.0	2.4	168.8
All 'Shi'	1097.4	1.9	0.2	-	-	1.0	-	0.1	12.0	2.5	2.4	213.5
All 'Gun'	1187.8	0.8	0.1	-	-	0.4	-	0.0	10.7	1.8	2.4	145.2
Aichi	1103.0	1.4	-	-	-	0.2	-	-	12.2	0.8	2.2	162.8
Akita	1310.1	0.6	0.2	-	-	0.6	-	0.1	12.5	2.4	1.5	161.2
Aomori	1349.1	1.8	0.3	-	-	0.9	-	0.1	18.9	3.9	2.1	217.1
Chiba	1187.1	1.2	0.1	-	-	0.5	-	0.1	10.3	1.4	5.4	135.0
Ehime	1162.6	0.5	0.1	-	-	0.3	-	-	15.4	1.6	4.1	149.4
Fukui	1324.6	1.5	0.1	-	-	0.1	-	-	8.8	2.1	1.5	155.2
Fukuoka	1179.6	0.7	0.1	-	-	0.6	-	0.0	14.3	2.5	2.2	203.8
Fukushima	1148.1	0.9	0.3	-	-	0.9	-	0.0	14.5	1.9	1.8	144.3
Gifu	1186.3	1.8	0.2	-	-	0.3	-	0.2	11.3	1.7	1.9	164.8
Gumma	1119.2	0.6	0.1	-	-	0.5	-	-	18.0	1.9	4.0	121.7
Hiroshima	1121.5	2.1	0.1	-	-	0.5	-	-	8.3	1.7	2.2	158.5
Hokkaido	1173.7	0.8	0.1	-	-	1.2	-	0.1	15.4	3.4	1.2	249.6
Ibigo	1072.5	1.8	0.0	-	-	0.4	-	0.1	10.5	1.9	1.2	169.3
Ibaraki	1186.3	0.7	0.1	-	-	0.5	-	-	5.9	1.2	4.9	110.5
Ishikawa	1348.6	0.8	0.2	-	-	0.2	-	-	13.6	2.1	2.6	169.0
Iwate	1369.5	1.0	0.4	-	-	0.6	-	0.1	13.8	3.2	1.1	206.4
Kagawa	1175.0	0.8	0.1	-	-	0.2	-	-	9.5	1.8	4.2	146.1
Kagoshima	1185.2	0.2	-	-	-	0.3	-	-	5.7	3.5	5.5	153.1
Kanagawa	967.9	1.6	0.2	-	-	1.0	-	0.1	11.4	1.9	2.1	178.5
Kochi	1191.4	2.4	0.2	-	-	0.7	-	-	3.6	1.2	3.6	153.0

TABLE 70.- DEATH RATES FOR SELECTED CAUSES BY PREFECTURE: JAPAN, 1947

Area	All Causes	Typhoid fever (1)	Paratyphoid fever (2)	Plague (3)	Cholera (4)	Cerebrospinal (Meningococcus) meningitis		Scarlet fever (8)	Whooping cough (9)	Diphtheria (10)	Tetanus (12)	Tuberculosis all forms (13-22)
						Anthrax (7)	(6)					
Kumamoto	1169.3	0.2	0.1	-	-	-	0.3	0.1	5.4	2.1	2.3	149.5
Kyoto	1077.5	1.1	-	-	-	-	1.2	0.1	8.9	0.7	1.0	214.1
Mie	1178.6	2.3	0.5	-	-	-	0.5	-	7.6	1.1	2.2	150.6
Miyagi	1111.6	1.5	0.5	-	-	-	0.9	0.1	17.7	1.8	1.7	164.4
Miyazaki	1197.3	0.3	-	-	-	-	0.5	0.1	4.3	5.1	5.6	167.8
Nagano	1070.7	0.8	0.0	-	-	-	0.4	0.0	11.3	1.0	2.3	122.9
Nagasaki	1228.5	0.6	-	-	-	-	0.4	0.1	6.8	3.8	3.0	169.4
Nara	1201.2	2.6	-	-	-	-	0.1	0.1	9.5	2.1	2.0	154.4
Niigata	1264.2	1.0	0.2	-	-	-	0.2	0.0	10.4	2.2	1.6	169.8
Oita	1331.8	0.3	0.1	-	-	-	0.5	-	7.6	3.3	2.7	176.1
Okayama	1156.5	2.7	0.1	-	-	-	0.2	0.1	5.7	1.6	2.1	144.5
Osaka	1079.5	1.5	0.2	-	-	-	1.4	0.1	12.0	1.2	1.2	201.5
Saga	1289.4	0.6	-	-	-	-	0.4	-	9.6	4.1	4.5	172.2
Saitama	1176.2	1.4	0.2	-	-	-	0.5	0.1	15.7	0.8	2.7	133.6
Shiga	1254.1	0.8	0.3	-	-	-	0.3	-	13.5	1.0	0.8	150.9
Shimane	1299.6	0.9	0.1	-	-	-	0.5	0.1	5.7	1.9	1.5	191.4
Shizuoka	1028.9	1.0	0.2	-	-	-	0.4	0.0	12.2	1.5	2.7	129.9
Tochigi	1118.5	1.2	0.1	-	-	-	0.1	0.1	11.1	1.7	3.0	116.0
Tokushima	1243.9	1.1	-	-	-	-	-	-	4.5	1.7	3.6	172.1
Tokyo	977.0	1.6	0.3	-	-	-	1.3	0.1	10.3	2.3	1.8	209.5
Tottori	1210.9	0.7	-	-	-	-	1.6	0.2	9.4	2.1	1.2	166.7
Toysama	1396.6	1.7	0.2	-	-	-	0.2	-	14.8	2.7	1.1	186.3
Wakayama	1113.6	1.6	0.3	-	-	-	-	-	3.1	0.5	1.8	141.6
Yamagata	1279.6	0.6	0.1	-	-	-	1.0	0.1	21.5	1.5	1.4	148.4
Yamaguchi	1199.7	0.7	-	-	-	-	0.3	-	10.8	2.5	3.0	189.0
Yamaguchi	1066.8	0.4	0.1	-	-	-	0.5	-	6.1	1.3	3.5	92.0

See footnotes at end of table.

TABLE 70.- DEATH RATES FOR SELECTED CAUSES BY PREFECTURE: JAPAN, 1949 - Cont'd
(Rates per 100,000 population)

Area	Leprosy (23)	Glanders (26a)	Dysentery (27)	Malaria (28)	Syphilis (30)	Influenza (33)	Smallpox (34)	Acute Poliomyelitis and acute polio encephalitis			Yellow fever (38a)	Rabies (38b)	Typhus fever and typhus like diseases (39)
								(35)	(36)	(37b)			
All Japan	0.2	0.0	9.5	0.1	6.7	0.6	0.0	15.1	1.2	1.4	-	0.1	0.1
All "Shis"	0.0	-	9.3	0.1	9.0	0.5	0.0	16.3	1.1	2.2	-	0.1	0.1
All "Guns"	0.3	0.0	9.6	0.2	5.4	0.7	0.0	14.5	1.3	1.0	-	0.1	0.1
Aichi	0.2	-	14.0	0.1	6.0	0.5	-	16.8	1.7	2.2	-	-	0.0
Akita	0.2	-	4.1	0.3	5.9	0.6	-	18.1	1.4	0.5	-	-	0.1
Aomori	0.1	-	3.9	-	8.6	1.9	-	19.1	1.8	0.7	-	-	0.2
Chiba	0.0	-	14.1	0.1	8.2	0.6	0.0	12.5	1.2	1.0	-	0.9	0.1
Chino	0.2	-	6.7	0.1	4.4	1.5	0.1	9.0	1.4	1.2	-	-	0.4
Fukui	0.3	-	4.5	0.3	4.3	0.9	-	10.4	0.8	3.5	-	-	-
Fukuoka	0.0	-	4.2	0.1	9.7	0.8	-	24.2	1.4	0.9	-	-	0.1
Fukushima	0.1	-	10.3	0.1	5.6	0.3	-	18.2	1.2	0.0	-	-	-
Gifu	-	-	14.5	0.2	5.4	0.6	-	10.2	1.2	1.5	-	-	-
Gunma	0.7	-	24.3	0.1	6.6	0.5	-	17.2	1.6	0.4	-	0.4	-
Hiroshima	0.0	-	10.0	0.1	4.4	1.0	-	7.8	1.5	2.4	-	-	0.1
Hokkaido	0.0	-	2.3	0.1	8.0	0.4	0.0	28.1	1.4	0.0	-	-	-
Iyogo	0.1	-	4.6	0.2	5.5	0.5	0.1	12.1	1.2	2.2	-	0.0	0.1
Ibaraki	0.0	-	26.3	0.0	6.1	0.0	-	12.5	1.1	1.1	-	0.1	-
Ishikawa	0.1	-	5.2	0.1	5.7	0.6	-	18.7	0.9	1.0	-	-	-
Iwate	0.2	-	8.9	-	5.7	0.6	-	16.3	1.5	0.2	-	-	-
Izumi	0.1	-	10.7	-	5.6	1.1	-	2.8	1.5	1.6	-	-	-
Kagawa	0.5	-	5.4	0.3	7.7	0.6	-	12.6	1.9	0.8	-	-	-
Kanagawa	-	-	10.1	0.1	9.3	0.5	-	11.7	1.2	1.9	-	0.3	-
Kochi	0.3	-	6.2	0.2	6.4	0.5	-	2.4	1.1	0.1	-	-	-

TABLE 70.- DEATH RATES FOR SELECTED CAUSES BY PREFECTURE: JAPAN, 1949 - Cont'd
(Rates per 100,000 population)

Area	Leprosy (23)	Glanders (24a)	Dysentery (27)	Malaria (28)	Syphilis (30)	Influenza (33)	Smallpox (34)	Measles (35)	Acute P. oligomyelitis and acute polio encephalitis				Typhus fever and typhus like diseases		
									(36)	(37b)	(38a)	(38b)	(39)	(39b)	(39c)
Kumamoto	3.1	-	6.0	0.1	5.8	0.9	-	12.3	1.3	4.4	-	-	-	0.2	-
Kyoto	-	-	4.8	-	9.0	0.5	-	19.6	0.5	1.0	-	-	-	0.2	-
Mie	0.3	-	5.1	0.3	6.0	0.8	-	17.3	0.9	1.7	-	-	-	0.1	-
Miyagi	0.6	0.1	3.2	0.1	7.2	0.1	0.1	14.7	1.5	0.4	-	-	-	0.2	-
Miyazaki	0.3	-	12.8	-	7.5	0.6	-	9.0	1.6	3.2	-	-	-	-	-
Nagano	-	-	4.3	0.2	5.3	0.3	-	8.1	1.2	1.0	-	-	-	-	-
Nagasaki	0.4	-	5.4	0.2	9.0	0.6	-	19.4	1.3	0.4	-	0.1	-	-	-
Nara	-	-	3.1	0.3	7.8	0.5	-	24.6	1.5	1.9	-	-	-	-	-
Niigata	0.1	-	14.9	0.1	4.2	0.8	-	23.0	1.0	0.4	-	-	-	0.1	-
Oita	0.3	-	8.2	0.1	6.4	0.9	-	19.1	3.1	0.6	-	-	-	0.3	-
Okayama	-	-	7.3	0.1	5.1	0.4	-	3.2	1.2	2.1	-	-	-	0.7	-
Oosaka	0.0	-	4.2	0.1	8.3	0.6	0.2	24.4	1.2	1.4	-	-	-	0.1	-
Saga	-	-	5.4	0.1	8.8	1.5	-	16.1	1.4	0.8	-	-	-	-	-
Saitama	0.1	-	26.3	0.1	5.6	0.1	-	12.4	0.8	1.8	-	0.6	-	0.0	-
Shiga	0.2	-	2.0	0.7	4.8	0.3	-	21.0	0.9	3.4	-	-	-	-	-
Shimane	-	-	13.4	0.4	6.6	1.1	-	15.7	1.2	1.8	-	-	-	0.1	-
Shizuoka	0.1	-	14.4	0.0	6.0	0.8	0.0	10.7	1.5	2.6	-	-	-	0.1	-
Tochigi	0.1	-	19.2	0.3	7.8	0.4	0.1	6.1	1.2	0.4	-	0.1	-	-	-
Tokushima	0.2	-	8.0	0.7	6.5	0.7	-	5.3	0.9	3.1	-	-	-	-	-
Tokyo	0.1	-	15.0	0.1	9.2	0.5	0.0	10.9	0.8	1.5	-	0.4	-	0.1	-
Tottori	0.5	-	5.6	0.3	5.6	1.2	-	4.9	0.3	0.7	-	-	-	-	-
Toyama	-	-	6.6	0.1	4.4	0.2	-	20.6	1.1	4.3	-	-	-	-	-
Wakayama	0.1	-	4.2	0.3	4.9	0.7	0.1	16.8	1.1	2.9	-	-	-	0.4	-
Yamagata	0.1	-	5.0	0.1	4.5	0.3	-	14.1	0.8	0.1	-	-	-	-	-
Yamaguchi	-	-	8.2	0.4	5.5	0.8	0.1	15.0	1.2	3.0	-	-	-	-	-
Yamaguchi	0.2	-	7.2	0.2	3.6	0.5	-	15.8	0.7	1.4	-	-	-	-	-

See footnotes at end of table.

TABLE 70.-- DEATH RATES FOR SELECTED CAUSES BY PREFECTURE: JAPAN, 1949 - Cont'd
(Rates per 100,000 population)

Area	Cancer and other malig- nant tumors (45-55)	Diabetes Mellitus (61)	Beriberi (68)	Simple Meningitis (81a)	Intercranial le- sions of vascular origin (83)		Diseases of the heart (90-95)		Pneumonia Bronchitis (all forms) (107-109)		Pleu- sury (not specified as tuberculosis) (110)		Ulcer of the stomach or duodenum (117)	
All Japan	71.6	2.3	6.8	11.1	122.4	64.3	31.3	68.9	7.3	23.7				
All "Shi"	76.7	1.8	6.4	10.3	99.9	58.9	21.4	68.4	6.8	22.7				
All "Gun"	67.9	2.5	7.0	11.5	134.4	67.1	36.5	69.2	7.6	24.3				
Aichi	71.6	2.7	6.7	13.5	118.2	66.5	32.0	68.3	6.7	20.0				
Akita	57.4	1.4	10.3	16.2	188.2	69.3	37.1	63.3	7.5	18.2				
Aomori	53.2	1.2	16.6	10.4	137.1	53.0	42.0	92.0	7.9	17.5				
Chiba	82.2	2.3	5.2	13.2	152.3	74.9	30.0	71.5	8.3	29.0				
Enme	73.1	2.4	3.0	13.1	106.3	63.8	35.9	71.3	6.8	19.7				
Fuku	73.7	3.6	7.3	13.7	120.0	69.2	38.5	65.2	8.0	24.5				
Fukuoka	77.8	2.4	6.4	9.5	105.9	54.4	25.6	73.9	7.7	25.3				
Fukushima	64.2	1.4	5.7	12.5	145.4	64.4	31.8	86.3	8.4	22.9				
Gifu	67.0	2.4	5.5	17.1	122.8	67.3	39.4	62.8	7.9	21.6				
Gunma	71.6	1.8	4.2	9.4	133.5	75.0	26.8	81.4	5.2	24.1				
Hiroshima	74.3	3.9	6.1	11.2	115.2	63.1	30.2	55.0	6.6	28.0				
Hokkaido	59.3	2.1	8.4	14.4	89.1	54.6	30.5	79.5	7.4	14.6				
Iwate	70.2	2.1	6.6	11.2	96.2	57.5	27.4	55.6	7.2	21.9				
Ibaraki	74.9	2.1	6.4	8.6	159.8	77.8	38.1	62.4	8.2	28.4				
Ishikawa	83.4	4.7	11.7	16.5	128.0	73.7	44.0	77.4	6.3	30.5				
Iwate	47.9	1.6	14.1	13.3	181.5	65.2	15.2	103.1	6.8	22.8				
Kagawa	80.3	4.0	1.9	13.2	116.6	65.2	28.7	61.6	6.4	26.4				
Kagoshima	55.1	2.0	5.7	11.0	128.4	62.0	31.9	69.9	9.2	28.0				
Kanagawa	66.8	1.6	4.8	8.8	105.9	53.2	16.5	69.4	5.9	23.0				
Kochi	68.7	6.0	7.5	11.7	144.6	51.4	29.5	54.3	8.7	21.9				

TABLE 70.- DEATH RATES FOR SELECTED CAUSES BY PREFECTURE: JAPAN, 1949 - Cont'd
(Rates per 100,000 population)

Area	Cancer and other malignant tumors (45-55)	Diabetes Mellitus (61)	Beriberi (68)	Simple Meningitis (81a)	Intracranial lesions of vascular origin (83)	Diseases of the heart (90-95)	Bronchitis (106)	Pneumonia (all forms) (107-109)	Fleury (not specified as tuberculosis) (110)	Ulcer of the stomach or duodenum (117)
Kimamoto	66.8	2.2	5.2	9.2	123.4	67.7	32.7	68.8	10.8	29.2
Kyoto	77.6	2.4	8.0	10.2	94.7	57.6	19.8	54.1	6.0	23.5
Kie	73.8	2.3	7.4	13.9	120.2	69.8	43.7	59.7	6.8	27.5
Miyagi	73.7	1.5	8.9	8.8	139.6	66.4	23.4	76.4	5.9	18.6
Miyazaki	61.0	2.0	5.4	13.3	118.6	71.3	33.8	62.9	9.5	26.9
Nagano	80.0	2.1	3.6	7.1	151.2	75.3	29.3	59.7	5.7	25.0
Nagasaki	65.4	2.8	5.7	13.0	103.5	62.0	40.1	78.9	9.8	24.2
Nara	99.4	2.1	7.3	11.5	120.3	65.2	29.7	61.3	5.8	27.7
Niigata	83.1	2.9	8.7	12.4	159.7	66.2	46.5	75.3	8.1	22.2
Oita	69.3	2.7	6.4	14.6	132.9	71.3	45.0	73.3	10.2	33.8
Okayama	77.3	3.4	6.6	11.9	122.8	63.7	32.6	51.8	8.7	26.4
Osaka	76.9	1.4	10.7	9.1	84.3	59.6	19.5	75.8	5.7	22.4
Saga	79.0	2.6	11.5	11.5	127.5	64.5	35.6	58.9	7.4	27.6
Saitama	73.6	1.8	4.7	9.9	137.5	73.6	42.0	71.9	7.2	23.0
Shiga	75.9	2.7	8.2	8.9	121.1	73.5	34.4	58.4	7.0	27.3
Shimane	76.5	4.3	5.5	10.2	148.6	80.2	28.1	62.3	9.0	32.5
Shizuoka	57.5	1.9	5.4	11.5	116.9	59.9	26.1	61.5	5.3	21.9
Tochigi	74.5	1.4	5.0	8.2	142.3	68.0	35.1	70.0	6.5	25.6
Tokushima	72.5	2.7	3.4	14.0	108.0	76.6	38.6	69.6	8.1	24.2
Tokyo	71.7	1.4	5.4	5.9	94.4	56.4	15.3	65.9	6.3	20.3
Tottori	83.1	4.3	8.6	8.6	113.8	62.7	31.8	50.5	8.9	29.6
Toyama	81.8	3.3	11.0	15.0	127.8	68.4	49.3	79.6	7.4	26.0
Wakayama	85.5	3.5	3.5	8.6	116.8	59.5	26.8	56.0	6.6	23.3
Yamagata	77.9	2.0	8.7	11.7	172.6	72.1	42.5	77.4	8.6	23.3
Yamaguchi	70.1	2.6	6.6	11.5	118.3	63.1	27.0	68.5	9.7	30.3
Yamomashi	73.8	2.0	1.7	6.5	122.9	73.8	36.4	64.1	8.3	24.4

See footnotes at end of table.

TABLE 70.- DEATH RATES FOR SELECTED CAUSES BY PREFECTURE: JAPAN, 1949 - Cont'd
(Rates per 100,000 population)

Area	Diarrhea enteritis and ulceration of the in- testines (all ages)		Diarrhea enteritis and ulceration of the in- testines (under 2 years)		Diarrhea enteritis and ulceration of the in- testines (2 years and over)		Appendicitis (121)	Nephritis (all forms) (130-132)	Diseases of pregnancy childbirth and the puerperium (140-150)
	(119-120)	(119)	(119)	(120)	(120)	(121)			
All Japan	87.8	50.9		36.9		4.0		41.2	5.5
All "Shi"	67.4	42.2		25.1		5.5		37.1	5.7
All "Gun"	98.6	55.4		43.2		3.2		43.3	5.4
Aichi	71.9	40.6		31.3		3.6		30.6	4.4
Akita	125.8	89.0		36.8		4.1		51.2	9.2
Aomori	144.0	114.2		29.8		4.2		30.7	7.4
Chiba	64.8	34.2		30.6		4.4		49.1	5.6
Ehime	79.3	39.1		40.1		3.1		37.0	5.1
Fukui	144.6	74.5		70.1		4.1		36.7	6.0
Fukuoka	85.3	58.2		27.1		3.9		43.5	5.2
Fukushima	84.6	53.5		31.1		4.0		35.5	6.3
Gifu	101.4	57.8		43.6		3.3		33.3	4.8
Gunma	75.6	32.9		42.6		3.1		41.5	5.0
Hiroshima	79.4	39.1		40.4		4.0		38.7	4.9
Hokkaido	106.4	76.4		28.0		4.5		27.1	7.1
Hyogo	90.9	49.2		41.7		4.0		35.8	5.6
Ibaraki	80.2	44.0		36.2		3.8		51.3	5.7
Ishikawa	120.9	82.2		38.8		4.7		38.2	5.2
Iwate	119.3	90.5		28.8		4.8		38.1	8.0
Kagawa	73.5	34.8		38.8		3.9		43.8	4.6
Kagoshima	100.3	59.2		41.0		2.5		57.8	8.2
Kanagawa	46.1	26.6		19.5		3.6		32.5	4.6
Kochi	72.0	28.6		43.4		2.3		48.0	6.1

TABLE 70.- DEATH RATES FOR SELECTED CAUSES BY PREFECTURE: JAPAN, 1949 - Cont'd
(Rates per 100,000 population)

Area	Diarrhea enteritis and ulceration of the in- testines (all ages) (119-120)	Diarrhea enteritis and ulceration of the in- testines (under 2 years) (119)	Diarrhea enteritis and ulceration of the in- testines (2 years and over) (120)	Appendicitis (121)	Nephritis (all forms) (130-132)	Diseases of pregnancy childbirth and the puerperium (140-150)
Kumamoto	92.0	46.4	45.6	3.3	47.6	6.8
Kyoto	70.1	37.5	32.6	4.9	30.8	4.5
Mie	97.5	52.1	45.4	3.0	41.5	5.1
Miyagi	76.2	54.0	22.2	4.5	37.8	5.1
Miyazaki	93.6	52.1	41.5	3.7	57.7	8.1
Nagano	65.8	28.6	37.2	3.8	44.4	4.8
Nagasaki	88.4	59.7	28.7	2.6	43.4	6.1
Nara	101.8	56.4	45.4	3.9	49.4	3.8
Niigata	112.3	57.7	54.6	5.8	46.3	5.1
Oita	103.7	54.9	48.8	4.8	51.8	5.8
Okayama	87.2	32.9	54.2	4.1	39.0	4.7
Osaka	81.5	49.0	32.5	5.0	43.9	5.2
Saga	97.9	67.6	30.3	4.7	60.7	5.0
Saitama	90.5	42.5	48.1	3.2	46.5	6.1
Shiga	134.6	67.5	67.1	4.8	47.5	3.1
Shimane	101.4	50.4	51.0	4.4	46.9	5.4
Shizuoka	74.2	40.7	33.5	2.4	35.6	4.4
Tochigi	101.9	44.9	56.9	3.6	42.3	5.4
Tokushima	95.8	40.9	54.9	2.5	44.6	4.3
Tokyo	40.4	27.7	12.7	4.6	37.2	4.8
Tottori	100.1	52.5	47.6	4.1	61.6	6.1
Toyama	171.7	100.5	71.3	4.5	49.9	5.3
Wakayama	98.4	54.8	43.6	3.8	26.6	3.7
Yamagata	125.2	78.5	46.7	4.4	48.3	5.0
Yamaguchi	79.3	46.4	32.9	3.5	44.0	5.1
Yamanashi	88.1	47.6	40.4	3.8	48.3	5.0

See footnotes at end of table.

TABLE 70.- DEATH RATES FOR SELECTED CAUSES BY PREFECTURE: JAPAN, 1949 - Cont'd
(Rates per 100,000 population)

Area	Congenital Malformations (157)	Congenital debility (158)	Premature birth (159)	Injury at birth (160)	Senility (162)	Suicide (163-164)	Homicide (165-168)	Accidental deaths (169-195)	Sudden death ill-defined and unknown causes (199-200)	All other causes
All Japan	8.0	45.4	16.6	1.3	81.1	17.3	2.0	42.0	20.7	152.5
All "Shi"	8.0	35.1	15.0	1.5	57.6	18.7	2.5	40.7	20.0	135.1
All "Gun"	8.0	50.8	17.5	1.2	93.6	16.5	1.8	42.6	21.0	161.7
Aichi	7.7	50.0	18.2	1.2	81.2	16.9	1.9	35.3	16.3	136.5
Akita	7.5	86.7	24.9	1.7	78.8	15.3	1.6	44.6	22.0	157.9
Aomori	10.0	89.2	20.1	0.9	77.3	10.7	1.5	39.3	30.5	172.8
Chiba	7.9	41.1	23.9	1.9	99.8	17.2	4.0	32.8	21.1	151.9
Ehime	8.6	44.4	16.6	1.8	117.0	15.5	1.6	50.7	19.6	170.6
Fukui	7.6	65.9	20.8	1.2	119.1	20.9	1.3	57.0	32.7	188.8
Fukuoke	7.3	45.3	14.3	1.1	69.2	15.7	3.9	57.3	19.9	153.3
Fukushima	10.1	45.4	18.3	1.1	77.0	12.3	1.7	44.9	21.3	141.8
Gifu	7.7	59.0	20.5	1.2	89.5	21.8	1.8	39.0	17.3	156.2
Gumma	10.4	36.8	13.8	1.3	69.1	19.8	1.8	38.2	16.6	154.7
Hiroshima	7.9	38.4	13.7	1.4	96.5	17.7	1.7	44.7	17.4	160.4
Hokkaido	8.9	39.2	12.6	1.1	55.1	13.5	2.2	54.2	29.4	140.3
Iyogo	6.5	35.9	15.6	1.1	78.3	21.4	2.2	42.1	20.7	144.6
Ibarak	9.3	53.5	24.7	1.1	89.3	12.4	2.6	34.3	19.7	160.5
Ishikawa	9.3	50.6	21.4	1.2	111.5	16.8	1.5	41.2	18.4	210.7
Iwate	8.8	70.7	28.1	1.4	83.2	14.9	1.8	55.2	24.1	169.7
Kagawa	7.6	59.1	17.3	1.7	110.3	14.6	2.2	46.1	31.6	168.3
Kagoshima	6.8	39.1	13.6	1.2	91.9	12.4	1.0	47.6	17.9	183.8
Kenagawa	6.7	23.0	9.3	1.4	58.9	17.5	2.4	45.6	19.5	109.9
Kochi	6.5	50.8	21.5	2.3	132.8	14.9	2.3	46.9	22.9	176.2


TABLE 70.- DEATH RATES FOR SELECTED CAUSES BY PREFECTURE: JAPAN, 1949 - Cont'd
(Rates per 100,000 population)

Area	Congenital Malformations (157)	Congenital debility (158)	Premature birth (159)	Injury at birth (160)	Senility (162)	Suicide (163-164)	Homicide (165-168)	Accidental deaths (169-195)	Sudden death ill-defined and unknown causes (199-200)	All other causes
Kumamoto	7.0	44.1	16.3	1.3	92.5	13.0	3.8	41.6	16.7	173.3
Kyoto	7.4	36.1	16.3	2.0	77.5	26.2	1.8	31.3	15.7	136.4
Mie	7.6	46.9	19.4	1.2	84.9	16.9	2.1	35.2	17.9	171.9
Miyagi	16.5	48.8	16.6	2.1	65.4	10.5	1.7	41.8	19.1	131.6
Miyazaki	8.4	54.2	15.8	1.9	86.0	14.1	1.3	43.2	16.1	170.0
Nagano	6.7	30.1	13.7	0.8	81.2	24.2	1.9	37.1	18.1	139.9
Nagasaki	8.4	55.4	14.9	1.1	106.5	11.3	1.9	51.2	18.5	192.4
Nara	6.3	51.1	16.7	1.3	89.4	21.3	1.3	31.7	30.5	151.9
Niigata	7.9	41.9	16.7	1.2	69.3	20.2	1.1	41.5	20.6	159.3
Oita	7.2	55.4	16.6	1.7	117.0	17.3	2.3	42.9	19.7	196.4
Okayama	7.6	52.6	17.0	1.4	116.6	16.5	1.2	41.1	18.2	167.1
Osaka	7.4	38.4	15.5	1.0	56.4	19.7	1.7	35.5	19.0	124.0
Saga	9.4	70.1	18.0	1.8	89.4	13.1	2.5	55.7	22.0	187.4
Saitama	2.1	52.0	18.3	1.1	81.0	15.1	1.9	30.6	19.7	150.7
Shiga	6.0	54.7	18.7	1.6	97.3	24.7	1.7	29.4	21.6	190.1
Shimane	6.4	49.7	19.5	1.1	104.5	17.6	1.6	44.7	16.3	170.3
Shizuoka	8.1	41.6	13.5	0.9	75.4	19.3	2.0	35.9	17.8	145.4
Tochigi	9.8	37.9	14.3	0.9	76.2	16.3	1.8	35.5	20.3	146.7
Tokushima	8.1	60.8	19.3	0.9	116.3	18.5	1.3	45.9	24.0	175.8
Tokyo	6.9	24.8	12.6	1.4	46.8	20.5	2.4	34.2	21.6	109.6
Tottori	10.2	48.2	15.3	1.6	120.2	12.8	2.0	38.8	13.7	154.2
Toysama	10.0	59.3	26.2	2.5	64.9	19.6	0.9	55.7	24.4	187.8
Wakayama	7.5	34.6	16.1	1.5	96.8	26.2	1.2	33.7	26.7	166.6
Yamagata	7.5	65.6	22.7	2.1	73.7	18.0	1.3	42.2	19.4	147.4
Yamaguchi	7.3	43.8	9.7	0.8	103.1	19.3	2.5	57.6	21.5	154.1
Yamaguchi	10.5	31.7	11.8	0.7	71.7	12.2	1.7	42.8	22.6	168.0


Footnotes: Data refer to deaths of Japanese nationals in Japan. A dash (-) indicates that no deaths were reported and the death rate was zero. A rate of 0.0 indicates that there were some deaths but the rate was less than 0.1.

Sources: Rates were computed by Public Health and Welfare Section, GHQ, SCAP. Sources of death data were Monthly Vital Statistics Schedule Reports, Ministry of Welfare.

BIRTH AND DEATH RATES: JAPAN, 1875-1949




BIRTH RATES BY MONTH: JAPAN 1947-1949 AND MEDIAN RATE 1935-1941




PREFECTURAL VITAL STATISTICS


1949 CALENDAR YEAR



BIRTH AND DEATH RATES BY MONTH: JAPAN, 1939-1949



DEATH RATES BY MONTH: JAPAN 1947-1949 AND MEDIAN RATE 1935-1941









PREFECTURAL MORBIDITY STATISTICS


1949 CALENDAR YEAR

LEGEND

PREFECTURAL RATE FOR EACH DISEASE IS
EXPRESSED AS A PERCENTAGE OF THE
NATIONAL RATE FOR THE SAME DISEASE

	0.0 - 39.9 %
	40.0 - 79.9 %
	80.0 - 119.9 %
	120.0 - 159.9 %
	160.0 - 199.9 %
	200.0 % AND OVER

TUBERCULOSIS




DYSENTERY



MALARIA



DEATH RATES FOR TEN LEADING CAUSES OF DEATH IN 1949: JAPAN, 1923-1943 AND 1947-1949









PREFECTURAL MORBIDITY STATISTICS


1949 CALENDAR YEAR

LEGEND

PREFECTURAL RATE FOR EACH DISEASE IS
EXPRESSED AS A PERCENTAGE OF THE
NATIONAL RATE FOR THE SAME DISEASE

	0.0 — 39.9 %
	40.0 — 79.9 %
	80.0 — 119.9 %
	120.0 — 159.9 %
	160.0 — 199.9 %
	200.0 % AND OVER


INFLUENZA



SMALLPOX



MEASLES









PREFECTURAL MORBIDITY STATISTICS


1949 CALENDAR YEAR

LEGEND

PREFECTURAL RATE FOR EACH DISEASE IS
EXPRESSED AS A PERCENTAGE OF THE
NATIONAL RATE FOR THE SAME DISEASE

	0.0 — 39.9 %
	40.0 — 79.9 %
	80.0 — 119.9 %
	120.0 — 159.9 %
	160.0 — 199.9 %
	200.0 % AND OVER

SCARLET FEVER



WHOOPING COUGH



DIPHTHERIA









PREFECTURAL MORBIDITY STATISTICS


1949 CALENDAR YEAR

LEGEND

PREFECTURAL RATE FOR EACH DISEASE IS EXPRESSED AS A PERCENTAGE OF THE NATIONAL RATE FOR THE SAME DISEASE

	0.0 - 39.9 %
	40.0 - 79.9 %
	80.0 - 119.9 %
	120.0 - 159.9 %
	160.0 - 199.9 %
	200.0 % AND OVER


JAPANESE "B" ENCEPHALITIS



TYPHUS FEVER



PNEUMONIA









PREFECTURAL MORBIDITY STATISTICS


1949 CALENDAR YEAR

LEGEND


PREFECTURAL RATE FOR EACH DISEASE IS
EXPRESSED AS A PERCENTAGE OF THE
NATIONAL RATE FOR THE SAME DISEASE

	0.0 — 39.9 %
	40.0 — 79.9 %
	80.0 — 119.9 %
	120.0 — 159.9 %
	160.0 — 199.9 %
	200.0 % AND OVER


TYPHOID FEVER




PARATYPHOID FEVER




EPIDEMIC MENINGITIS



INFANT DEATH AND STILLBIRTH RATES: JAPAN, 1899-1949



INFANT DEATH RATES BY MONTH: JAPAN 1947-1949 AND MEDIAN RATE 1938-1942



PREFECTURAL VITAL STATISTICS








1949 CALENDAR YEAR

INFANT DEATH RATE

N


STILLBIRTH RATE

LEGEND
PREFECTURAL RATE EXPRESSED AS
A PERCENTAGE OF NATIONAL RATE

	0.0 — 49.9 %
	50.0 — 69.9 %
	70.0 — 89.9 %
	90.0 — 109.9 %
	110.0 — 129.9 %
	130.0 — 149.9 %
	150.0 % AND OVER


(A-14)

DEATH RATES FOR 8 LEADING CAUSES OF INFANT DEATHS IN 1949: JAPAN, 1923-1943 AND 1947-1949




INFANT DEATH AND STILLBIRTH RATES BY MONTH


JAPAN, 1939-1949




STILLBIRTH RATES BY MONTH: JAPAN 1947-1949 AND MEDIAN RATE 1935 - 1941



MARRIAGE AND DIVORCE RATES BY MONTH: JAPAN, 1939-1949



MARRIAGE RATES BY MONTH: JAPAN 1947-1949 AND MEDIAN RATE 1932-1938










PREFECTURAL VITAL STATISTICS

1949 CALENDAR YEAR

MARRIAGE RATE


DIVORCE RATE

LEGEND
PREFECTURAL RATE EXPRESSED AS
A PERCENTAGE OF NATIONAL RATE


	0.0 - 49.9 %
	50.0 - 69.9 %
	70.0 - 89.9 %
	90.0 - 109.9 %
	110.0 - 129.9 %
	130.0 - 149.9 %
	150.0 % AND OVER

(A-20)

MARRIAGE AND DIVORCE RATES: JAPAN, 1883-1949



DIVORCE RATES BY MONTH: JAPAN 1947-1949 AND MEDIAN RATE 1932-1938



PATIENT NO.		EPIDEMIOLOGICAL CASE CARD										HEALTH CENTER NO.	
1. NAME AND SEX <input type="checkbox"/> MALE <input type="checkbox"/> FEMALE		2. DATE OF BIRTH		3. ADDRESS									
				4. PLACE OF OCCURRENCE									
5. ONSET DATE OF FIRST EXAMINATION DATE CONFIRMED		6. INDUSTRY		7. OCCUPATION		8. OF FAMILY		9. PRINCIPAL OCCUPATION		10. JAPANESE AGE EUROPEAN AGE YEARS MONTHS			
11. IMMUNIZATION DATE OF LAST IMMUNIZATION NUMBER OF SMALLPOX VACCINATIONS DATE		12. DATE OF FIRST EXAMINATION DATE CONFIRMED		13. DISEASES IN PAST ASSOCIATED WITH THIS DISEASE		14. DATE OF ONSET DATE OF FIRST EXAMINATION DATE CONFIRMED		15. DATE OF LAST IMMUNIZATION DATE		16. DATE OF FIRST EXAMINATION DATE CONFIRMED			
17. BACTERIOLOGICAL TEST		18. BLOOD PHARYNX PLACE OF TEST		19. SPINAL FLUID OTHER		20. DATE DATE		21. DATE DATE		22. DATE DATE			
19. SOURCE		20. TO BE THOUGHT AS SOURCE OF INFECTION		21. PERSONS MATTERS		22. TIME OF CONTACT		23. PLACE		24. CONDITION OF CONTACT			
20. SIMILAR PATIENTS IN THE NEIGHBORHOOD <input type="checkbox"/> THERE IS <input type="checkbox"/> THERE IS NOT		21. TRIP DATE (TO WHERE)		22. CAME BACK DATE		23. OTHER SOURCES		24. DISEASES IN PAST ASSOCIATED WITH THIS CASE		25. OFFICE (OR SCHOOL)			
21. FAMILY		22. NAME SEX AGE		23. DISEASES IN PAST ASSOCIATED WITH THIS CASE		24. TEST (OR SCHOOL)		25. NAME SEX AGE		26. DISEASES IN PAST ASSOCIATED WITH THIS CASE			
22. PRECAUTIONS TAKEN		23. ISOLATION OF HEALTHY PEOPLE		24. DATE RELEASED DATE		25. DATE OF CLEANING PATIENT'S HOUSE		26. INTERCEPTION OF COMMUNICATION FROM OUTSIDE		27. DATE RELEASED DATE			
23. REMARKS		24. OTHER		25. DATE OF CLEANING PATIENT'S HOUSE		26. INTERCEPTION OF COMMUNICATION FROM OUTSIDE		27. DATE RELEASED DATE		28. DATE RELEASED DATE			
24. DIED		25. CURED		26. DIED		27. CURED		28. DIED		29. CURED			

EPIDEMIOLOGICAL CASE CARD

HEALTH CENTER NO.

PATIENT NO.

☐ MALE

☐ FEMALE

1. NAME AND SEX

2. DATE OF BIRTH

JAPANESE AGE

EUROPEAN AGE

YEARS

MONTHS

3. ADDRESS

PRESENT

AT TIME OF ONSET

4. PLACE OF ONSET

5. OCCUPATION

RELATIONSHIP WITH PATIENT

6. ADDRESS OF HEAD OF FAMILY

7. OF FAMILY

8. WITH PATIENT

9. FAMILY CONTACTS

DENSITY OF HOUSES

NO. OF MATS PER ONE PERSON

CONDITION OF HOUSE

GOOD

BAD

11. HOUSING & ENVIRONMENT

RELATIONSHIP WITH BUILDING FAMILY

SEPARATE ROOM

COM-PART-MENT

TOGETHER WITH OTHER PEOPLE

12. SICK ROOM

CLASS OF LIVING CONDITION

YEN A MONTH

13. CLASS OF LIVING CONDITION

SUBSIDY YES () PUBLIC () PRIVATE () NO () NEED () NEED NOT ()

14. PHYSICIAN REPORTING AND CASE FINDING

NAME OF INSTITUTION

HOSPITAL

CLINIC

HEALTH CENTER

CASE FOUND BY

MASS EXAMINATION

PATIENT VISITED

OTHER

LOCATION OF INSTITUTION

15. ONSET

DATE OF REPORT

NAME

17. REACTION TO TUBERCULIN TEST

WHETHER RECEIVED TUBERCULOSIS TEST

YES () TIMES () NO

DATE REACTION BECAME POSITIVE

DATE

BCG INOCULATION

WHETHER INOCULATED

YES () TIMES () NO

18. DISEASES IN PAST

YEAR

MONTH

19. TREATMENT

IN HOSPITAL

HEALTH CENTER

VISITING HOSPITAL

REST IN OWN HOUSE

TREATED OUTSIDE HOSPITAL

LEAVE AS IT IS

INJECTION

SURGICAL TREATMENT

PNEUMOTHORAX

TAKING DRUGS

20. MEASURES HEREAFTER

HOSPITALIZATION

VISITING HOSPITAL

HEALTH CENTER

LEAVE AS IN PRESENT CONDITION

16. PRESENT SYMPTOMS

GERM TEST AND RESULTS

DONE (POSITIVE, NEGATIVE)

UNDONE

METHOD OF TEST

SMEAR

CULTURE

COLONY PICKING

COUGH

HEMOPTYSIS

FEVER

SPUTUM

OTHER

NUTRITION

GOOD

MEDIUM

BAD

PROCESS AND OUTCOME

YEAR

MONTH

DAY

YEAR

MONTH

DAY

YEAR

MONTH

DAY

YEAR

MONTH

DAY

REMARKS

CURED



MORBIDITY SCHEDULE

CITY, TOWN AND VILLAGE OFFICE'S MARK	
HEALTH CENTER NO.	
PATIENT NO.	

NAME OF DISEASE	NAME	
DATE OF ONSET	SEX	<input type="checkbox"/> MALE <input type="checkbox"/> FEMALE
DATE OF FIRST EXAMINATION	DATE OF BIRTH	
DATE CONFIRMED	AGE (EUROPEAN)	YEARS MONTHS
DATE REPORTED	ADDRESS	
DATE OF LAST IMMUNIZATION	PLACE OF ONSET	
TOTAL AMOUNT OF VACCINE INNOCULATED	INDUSTRY PATIENT WORKS IN	
REACTION TO SMALLPOX	PATIENT'S OCCUPATION	
FREQUENCY OF INFECTION	PRINCIPAL OCCUPATION IN THE FAMILY	
CONTACT SOURCE	<input type="checkbox"/> WITHIN FAMILY <input type="checkbox"/> OFFICE <input type="checkbox"/> SCHOOL <input type="checkbox"/> TRIP <input type="checkbox"/> OTHER <input type="checkbox"/> UNKNOWN	SICK PERSONS IN FAMILY
	<input type="checkbox"/> WATER (<input type="checkbox"/> CHLORINATED <input type="checkbox"/> WELL <input type="checkbox"/> SPRING <input type="checkbox"/> RIVER <input type="checkbox"/> OTHER) <input type="checkbox"/> CANDIES <input type="checkbox"/> MILK <input type="checkbox"/> FRUITS & VEGETABLE <input type="checkbox"/> FISH, SHELL AND SEA PRODUCTS <input type="checkbox"/> OTHER	PERSONS OUT OF PERSONS
		NUMBER OF MATS PER ONE PERSON
		LOCATION OF INSTITUTION WHERE REPORTER WORKS

T**MORBIDITY SCHEDULE**

CITY, TOWN AND VILLAGE OFFICE'S MARK
HEALTH CENTER NO.
PATIENT NO.

DIAGNOSIS		NAME	
COMPLICATIONS		SEX	<input type="checkbox"/> MALE <input type="checkbox"/> FEMALE
METHOD OF CONFIRMATION	<input type="checkbox"/> GERM TEST <input type="checkbox"/> X-RAY <input type="checkbox"/> ANSCULTATION OR PERCUSSION <input type="checkbox"/> SURGICAL <input type="checkbox"/> OTHER	DATE OF BIRTH	
DATE OF ONSET		AGE (EUROPEAN)	
DATE OF FIRST EXAMINATION		ADDRESS	
DATE CONFIRMED		PLACE OF ONSET	
DATE REPORTED		OCCUPATION OF PATIENT	PRESENT <input type="checkbox"/> STILL WORKING <input type="checkbox"/> NOT AT WORK
CAUSE OF DISEASE	<input type="checkbox"/> MAL-NUTRITION <input type="checkbox"/> BAD ENVIRONMENT <input type="checkbox"/> OVER FATIGUE <input type="checkbox"/> CONCEPTION <input type="checkbox"/> ANXIETY <input type="checkbox"/> CONVALESCENCE <input type="checkbox"/> UNKNOWN <input type="checkbox"/> OTHERS	AT THE TIME OF ONSET	
GERM TEST AND RESULTS	<input type="checkbox"/> DONE <input type="checkbox"/> POSITIVE <input type="checkbox"/> NEGATIVE <input type="checkbox"/> UNDONE	PRINCIPAL OCCUPATION OF FAMILY	
DATE REACTION TO TUBERCULIN TEST BECAME POSITIVE		CONTACT	<input type="checkbox"/> FAMILY <input type="checkbox"/> OTHER <input type="checkbox"/> UNKNOWN
WHETHER INNOCULATED WITH BCG	<input type="checkbox"/> YES (TIMES) <input type="checkbox"/> NO	NUMBER OF MATS PER ONE PERSON	MATS
DATE OF LAST BCG INNOCULATION		NUMBER OF SICK PERSONS IN THE FAMILY	PERSONS OUT OF
TREATMENT	<input type="checkbox"/> IN HOSPITAL <input type="checkbox"/> TREATED OUTSIDE HOSP. <input type="checkbox"/> VISIT HOSPITAL <input type="checkbox"/> PNEUMOTHORAX <input type="checkbox"/> HEALTH CENTER <input type="checkbox"/> REST IN OWN HOUSE <input type="checkbox"/> LEAVE AS IT IS <input type="checkbox"/> SURGICAL TREAT- MENT <input type="checkbox"/> INJECTION <input type="checkbox"/> DRUGS	INSTITUTION WHERE REPORTER WORKS IN	<input type="checkbox"/> HOSPITAL <input type="checkbox"/> CLINIC <input type="checkbox"/> HEALTH CENTER
MEASURES TAKEN	<input type="checkbox"/> SENT TO HOSPITAL <input type="checkbox"/> VISIT HOSPITAL <input type="checkbox"/> HEALTH CENTER <input type="checkbox"/> AS PRESENT CONDITION	CASE FINDING	<input type="checkbox"/> MASS EXAMINATION <input type="checkbox"/> VISIT FROM OUTSIDE <input type="checkbox"/> OTHER
		LOCATION OF INSTITUTION REPORTER BELONGS TO	

Moore

